

12

Chapter Four
Differentiation

4.1

The Derivative as
a Functions

MATH-110

جمال السعدي
رياضيات - إحصاء

CH. 4.1

The derivative as a function المشتقة كدالة

The derivative of a function f at a fixed number a

is :
$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

If : we replace a by a variable x

We obtain f' as a new function

Called the derivative of f and defined by

equation :
$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

ALSAADI

Example:

If : $f(x) = 3x^2 - 1$

find $f'(x)$?

by def.

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

بالتعريف

$$= \lim_{h \rightarrow 0} \frac{[3(x+h)^2 - 1] - [3x^2 - 1]}{h}$$

$$= \lim_{h \rightarrow 0} \frac{3x^2 + 6xh + 3h^2 - 3x^2}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h(6x + 3h)}{h} = \lim_{h \rightarrow 0} (6x + 3h) = 6x$$

by rule

بقاعده الاشتقاق المباشر

$$y = 3x^2 - 1$$

$$y' = 6x$$

- If $y = f(x)$

المشتقة
رموز
The notations for the derivative are :

$$y' = f'(x) = \frac{dy}{dx} = \frac{df}{dx} = \frac{d}{dx} f(x) = Df(x)$$

- f is differentiable at a if $f'(a)$ exist .
موجوده
قابله للاشتقاق

- f is differentiable on open interval :
فترة مفتوحة
 (a, b) or (a, ∞) or $(-\infty, a)$ or $(-\infty, \infty)$
If it is differentiable at every number
in the interval .

• اذا كانت الداله قابله للاشتقاق عند a فإنها تكون متصله عند a

- Theorem : If f is differentiable at a
قابله للاشتقاق
then f is continuous at a

$$f(x) = |x|$$

* توجد دوال متصله ولكنها غير قابله للاشتقاق مثال:

- There are function that are continuous
but not differentiable .

for example : $f(x) = |x|$ is continuous at $x = 0$


غير قابله للاشتقاق

but not differentiable at $x = 0$

• $f(x) = |x|$ is **continuous** at $x = 0$

because : $\lim_{x \rightarrow 0^+} (x) = \lim_{x \rightarrow 0^-} (-x) = f(0) = 0 \Rightarrow$ **الداله متصله عند $x = 0$**
 قيمه الداله النهايه اليسرى النهايه اليمنى

$$F(x) = \begin{cases} x & \text{if } x > 0 \\ -x & \text{if } x < 0 \end{cases}$$




غير قابله للاشتقاق

• $f(x) = |x|$ is **not differentiable** at $x = 0$

because :

$$f'(x) = \begin{cases} 1 & \text{if } x > 0 \\ -1 & \text{if } x < 0 \end{cases}$$



$f'(0^+) \neq f'(0^-) \Rightarrow$ **الداله غير قابله للاشتقاق عند $x = 0$**
 المشتقة اليسرى \neq المشتقة اليمنى

Notes

• اذا كان المماس أفقي


$$f'(x) = m = 0 \text{ (ثابته)}$$

• اذا كان المماس يصنع زاويه حاده

$$f'(x) = m > 0 \text{ (تزايديه) موجب}$$

• اذا كان المماس يصنع زاويه منفرجه


$$f'(x) = m < 0 \text{ (تناقصيه) سالب}$$




المماس الأفقي

This functions are not differentiable at $x = a$


عند $x = a$
غير قابله للاشتقاق



عند $x = a$
غير قابله للاشتقاق
لانها غير متصله



عند $x = a$
غير قابله للاشتقاق
لوجود مماس رأسي



(a) A corner or Kink (b) A discontinuity (c) A vertical tangent


35-38 The graph of f is given. State with reasons, the numbers at which f is not differentiable

35 $f(x)$ is not differentiable

at $x = -4 \rightarrow$ (corner)

$x = 0 \rightarrow$ (discontinuous (Jump))


$x \approx 2.3 \rightarrow$ (vertical tangent)



36 $f(x)$ is not differentiable

at $x = 0 \rightarrow$ (discontinuous)


$x = 3 \rightarrow$ (corner)



37 $f(x)$ is not differentiable

at $x = -1 \rightarrow$ (vertical tangent)


at $x = 4 \rightarrow$ (corner)



38 $F(x)$ is not differentiable

at $x = -1 \rightarrow$ (discontinuous)

$x = 2 \rightarrow$ (corner)



A
L
S
A
A
D
I

Higher order derivative

المشتقات من الرتب العليا

الدالة

$$y = f(x)$$

المشتقة الأولى

$$y' = f'(x) = \frac{df}{dx}$$

المشتقة الثانية


$$y'' = f''(x) = \frac{d^2 f}{dx^2}$$

المشتقة الثالثة

$$y''' = f'''(x) = \frac{d^3 f}{dx^3}$$

المشتقة الرابعة

$$y^{(4)} = f^{(4)}(x) = \frac{d^4 f}{dx^4}$$



$$y = x^5 - 3x^3 + 4x^2 - 2x + 1 \text{ find } y^{(5)}?$$

$$y' = 5x^4 - 9x^2 + 8x - 2$$

$$y'' = 20x^3 - 18x + 8$$

$$y''' = 60x^2 - 18$$

$$y^{(4)} = 120x$$

$$y^{(5)} = 120$$

ملاحظة


إذا زادت رتبة المشتقة
عن درجة كثيره الحدود
فإن المشتقة = zero

إذا طلب $y^{(6)}$ الناتج مباشره $y^{(6)} = 0$


لأن رتبة المشتقة 6 زادت عن درجة الدالة y وهي 5.

A
L
S
A
A
D
I


رسم الدوال المشهورة ومشتقاتها




المشتقة




المشتقة




المشتقة





المشتقة



المشتقة




المشتقة



ALBAADI


مشتقة دالة من الدرجة n هو دالة من الدرجة $n - 1$

Match graph of each function in (a)–(c) with the graph of its derivative in 1–3. Give reasons for your choices



A
L
S
A
A
D
I

ملحوظة



ملحوظة (2)



الرسم يتكون من فرعين

بينهما زاوية (انكسار)

corner


تكون دوال متصلة ولكنها غير قابلة للاشتقاق
Continuous but not differentiable.

A
L
S
A
A
D
I

Find: $f'(-2)$

عند العدد $x = -2$

المماس لمنحنى الدالة افقى



Horizontal

$$\therefore f'(-2) = 0$$


True or false?

$$f'(2) > 0$$

المماس لأسفل

$$\therefore f'(2) < 0 \quad \text{تناقضيه}$$

$$\therefore f'(2) > 0 \quad \text{false}$$



True or false?


$f(x)$ is differentiable.

at $x = 1$

false

$f(x)$ not diff.

because : there is corner.



The accompanying figure shows the graph of $y = f(x)$


Then $f'(-2) =$

(a) -3

(b) 0

(c) 1

(d) 3



$$f'(-2)$$

عند العدد -2 $x = -$

المماس لمنحنى الدالة افقى

Horizontal


$$\therefore f'(-2) = 0$$

The accompanying figure shows the graph of $y = f(x)$

Then $f'(2) > 0$

(a) True

(b) False



المماس للمنحنى عند $x = 2$

يصنع زاوية حاده مع

الاتجاه الموجب لمحور x

\therefore الدالة تزايديه

$$\therefore f'(2) > 0$$

$$\therefore f'(2) > 0 \text{ True}$$

True or false?

متصله

* $f(x)$ is **continuous** at $x = 0$


There is (Jump) \Rightarrow false.

OR $\lim_{x \rightarrow 0^+} f(x) = 2$

$\lim_{x \rightarrow 0^-} f(x) = 3$

$\therefore f(x)$ is discontinuous at $x = 0$

لأن النهاية اليمنى \neq النهاية اليسرى.



f is differentiable at $x = 1$

(a) True


(b) false

$f(x)$ is differentiable
at $x = 1$

(false)

because : $f(x)$ is not diff. \rightarrow There is corner

at $x = 1$



كل التمنيات بالاجاز والتوفيق

السعدى

13

Chapter Four
Differentiation

4.2

Differentiation
Rules

MATH-110

جمال السعدي
رياضيات - إحصاء



CH 4.2

Derivatives of polynomials and exponential fun.

The product and quotient Rules.

Differentiation Rules

(1) $F(x) = c$ where c is constant. ^{ثابت}
 $F'(x) = 0$ (مشتقة الثابت = zero)

(2) $F(x) = ax$ where a is constant.
 $F'(x) = a$ (أخذ المعامل فقط)

(3) $F(x) = ax^n$
 $F'(x) = n \cdot a x^{n-1}$ (نضرب الأس من المعامل ونقص من الأس 1)

(4) $F(x) = g(x) \cdot h(x)$ * قاعده مشتقة حاصل ضرب والتين
 $F'(x) = g'(x) \cdot h(x) + h'(x) \cdot g(x)$
 الأولى: مشتقة الثانية: الثانية: مشتقة الأولى

(5) $F(x) = \frac{g(x)}{h(x)}$ * قاعده مشتقة خارج قسمة والتين
 $F'(x) = \frac{g' \cdot h - h' \cdot g}{(h)^2} = \frac{\text{أصل} \cdot \text{مشتقة الجاه} - \text{مشتقة الجاه} \cdot \text{مشتقة البسط}}{(\text{المقام})^2}$

$$(6) F(x) = (\quad)^n$$

$$F'(x) = n (\quad)^{n-1} \text{ مشتقة ما بداخل القوسا .}$$

$$(7) F(x) = \sqrt{\quad} \quad (\text{مشتقة الجذر التربيعي})$$

$$F'(x) = \frac{\text{مشتقة ما تحت الجذر}}{2\sqrt{\quad}}$$

$$(8) F(x) = \frac{a}{x^n}$$

$$F'(x) = \frac{-a \cdot n}{x^{n+1}}$$

* (نكس اشارة العدد a ثم ضرب n ثم نقسم على x^{n+1})

$$(9) F(x) = \frac{1}{x} \Rightarrow F(x) = \frac{a}{x}$$

$$F'(x) = \frac{-1}{x^2} \quad F'(x) = \frac{-a}{x^2}$$

(10) معادله المماس (equation of tangent line)

$$y = m(x - x_1) + y_1$$

* حيث (x_1, y_1) النقطة المعطاه * slope m مشتقة الدالة عند النقطة المعطاه

(11) معادله العمودي (equation of normal line)

or perpendicular line

$$y = -\frac{1}{m}(x - x_1) + y_1$$

* اذا كان المقام يتكون من عدد واحد فقط يتم توزيع عدد البسط على نفس المقام (12)
ثم الاختصار ثم الاشارة .

$$(13) \frac{d}{dx} [f \pm g] = \frac{df}{dx} \pm \frac{dg}{dx}$$

المشتقة تتوزع على مجموع وطرح الدوال

$$(14) F(x) = e^{h(x)}$$

مشتقة الدالة الأسية

$$F'(x) = e^{h(x)} \cdot h'(x)$$

\downarrow الدالة كما هي \downarrow مشتقة الأس

Example:

$$F(x) = e^{3x^2 - 2x}$$

$$F'(x) = e^{3x^2 - 2x} \cdot (6x - 2)$$

مشتقة الأس الدالة كما هي

Note:

$$\bullet F(x) = \sqrt{x} \longrightarrow F'(x) = \frac{1}{2\sqrt{x}}$$

$$\bullet F(x) = \frac{1}{x} \longrightarrow F'(x) = \frac{-1}{x^2}$$

$$\bullet F(x) = x\sqrt{x} \longrightarrow F'(x) = \frac{3}{2}\sqrt{x}$$

$$\bullet F(x) = e^x \longrightarrow F'(x) = e^x$$

Differentiate the following functions:

Find y' or f' ?

$$y = \sqrt{5} \rightarrow y' = 0$$

$$y = e^2 \rightarrow y' = 0$$

$$y = \pi^4 \rightarrow y' = 0$$

* مشتقة الثابت
Zero *

$$y = \sqrt{x^2 - 2x} \rightarrow y' = \frac{\text{مشتقة ما تحت الجذر}}{2\sqrt{\quad}} = \frac{2x - 2}{2\sqrt{x^2 - 2x}}$$

$$y' = \frac{2(x-1)}{2\sqrt{x^2 - 2x}} = \frac{x-1}{\sqrt{x^2 - 2x}}$$

$$y = \sqrt[3]{x^2 - 2x}$$

* أي جذر غير التربيعي يحول إلى قوس

$$y = (x^2 - 2x)^{\frac{1}{3}} \rightarrow y' = \frac{1}{3} (x^2 - 2x)^{\frac{1}{3} - 1} \cdot (2x - 2)$$

(2x-2) مشتقة ما بداخل القوس

$$\therefore y' = \frac{1}{3} (x^2 - 2x)^{-\frac{2}{3}} \cdot (2x - 2) = \frac{1 \cdot (2x - 2)}{3 (x^2 - 2x)^{\frac{2}{3}}}$$

$$\Rightarrow y' = \frac{2x - 2}{3 \sqrt[3]{(x^2 - 2x)^2}}$$

$$y = \frac{2x^3 - 6x^4}{2x^2}$$

$$y = \frac{2x^3}{2x^2} - \frac{6x^4}{2x^2}$$

(توزيع)

$$y = x - 3x^2$$

(إختصار)

$$y' = 1 - 6x$$

(اشتقاق)

* المقام يتكون من عدد واحد
توزع عدد وود البسط
على نفس المقام
ثم الاختصار
ثم الاشتقاق .

If: $f(x) = \frac{x^{3/2} + x^{5/2}}{x^{1/2}}$

find $f'(1)$?

* نفس طريقة التبريم السابق .

$$f(x) = \frac{x^{3/2}}{x^{1/2}} + \frac{x^{5/2}}{x^{1/2}}$$

$$f(x) = x + x^2$$

$$f'(x) = 1 + 2x$$

$$\Rightarrow f'(1) = 1 + 2(1) = 1 + 2 = \boxed{3}$$

→ (عند التمهيد نخرج الأسس)

$$* \frac{3}{2} - \frac{1}{2} = \frac{2}{2} = 1$$

$$* \frac{5}{2} - \frac{1}{2} = \frac{4}{2} = 2$$

$$y = \frac{3}{\sqrt[3]{x^2}}$$

* تحويل الجذر إلى صورته أسية
ثم رفعه للبطء بإشارة سالبة
ثم الاشتقاق.


$$y = \frac{3}{x^{2/3}}$$

$$y = 3x^{-2/3} \Rightarrow y' = 3 \cdot \frac{-2}{3} x^{-2/3-1} = -2x^{-5/3} = \frac{-2}{x^{5/3}}$$

$$= \frac{-2}{\sqrt[3]{x^5}} = \frac{-2}{x^3 \sqrt[3]{x^2}}$$

$$y = \frac{4}{x^5}$$

$$y' = \frac{-4 \cdot (5)}{x^{5+1}} = \frac{-20}{x^6}$$

* قاعدة اشتقاق:
 اشتقاق
 نكتب الإشارة العدد ثم نضربها في n

$$\frac{\text{عدد}}{x^n}$$

$$\frac{\text{نفس الإشارة العدد ثم نضربها في n}}{x^{n+1}}$$

$$y = -2x^5 + 3x^{-5} + \frac{1}{x} - \sqrt{x} + e^{3x}$$

من الالة نفسها
من الالة نفسها

$$y' = -10x^4 - 15x^{-6} - \frac{1}{x^2} - \frac{1}{2\sqrt{x}} + 3e^{3x}$$

Page 187

$$(7) \quad g(x) = \frac{3x-1}{2x+1}$$

Find $g'(x)$?

$$\begin{aligned} g'(x) &= \frac{(3)(1) - (2)(-1)}{(2x+1)^2} \\ &= \frac{3+2}{(2x+1)^2} \\ &= \frac{5}{(2x+1)^2} \end{aligned}$$

قاعدة
تستخدم هذه القاعدة
إذا كان البسط والقام
من الدرجة الأولى

$$f(x) = \frac{ax+b}{cx+d}$$

$$f'(x) = \frac{(a \cdot d) - (c \cdot b)}{(cx+d)^2}$$

$$(13) \quad y = \frac{x^3}{1-x^2}$$

$$y' = \frac{\begin{matrix} \frac{1}{3} \\ \downarrow \\ 3x^2 \end{matrix} \cdot \begin{matrix} \text{القام} \\ \downarrow \\ (1-x^2) \end{matrix} - \begin{matrix} \frac{1}{3} \\ \downarrow \\ (-2x) \end{matrix} \cdot \begin{matrix} \text{البسط} \\ \downarrow \\ x^3 \end{matrix}}{(1-x^2)^2}$$

$$= \frac{3x^2 - 3x^4 + 2x^4}{(1-x^2)^2} = \frac{3x^2 - x^4}{(1-x^2)^2}$$

Page 187

$$\textcircled{2} F(x) = \frac{x - 3x\sqrt{x}}{\sqrt{x}}$$

Find $F'(x)$?

$$F(x) = \frac{x}{\sqrt{x}} - \frac{3x\sqrt{x}}{\sqrt{x}}$$

$$= \frac{\sqrt{x} \cdot \sqrt{x}}{\sqrt{x}} - \frac{3x\sqrt{x}}{\sqrt{x}}$$

$$= \sqrt{x} - 3x$$


$$\Rightarrow F'(x) = \frac{1}{2\sqrt{x}} - 3 \quad \text{ممكن توحيده مقامات} = \frac{1 - 6\sqrt{x}}{2\sqrt{x}}$$

المقام يتكون من حد واحد
 = نوزع حدود البسط
 على نفس المقام

$$y = \frac{x^3}{3} + \frac{2}{x^2}$$

$$y' = \frac{3x^2}{3} + \frac{-2 \cdot (2)}{x^3}$$

$$\Rightarrow y' = x^2 - \frac{4}{x^3}$$



• $F(x) = x \cdot (\sqrt{x} + 3)$

Find $F'(x)$? يمكن

$$F(x) = x\sqrt{x} + 3x$$

*) مشتقه حاصل ضرب والتين
*) يمكن فك الأقواس أولاً
ثم الاشتقاق ثانياً
وهذا هو الأسرع

$$\Rightarrow F'(x) = \frac{3}{2}\sqrt{x} + 3$$

• $y = \frac{5}{(5x-1)^3} \Rightarrow y = 5(5x-1)^{-3}$

$$\Rightarrow y' = -15(5x-1)^{-4} \cdot \begin{matrix} 5 \\ \text{مشتقه} \\ \text{ما بداخل} \\ \text{القوس} \end{matrix} = \frac{-75}{(5x-1)^4}$$

• $y = x\sqrt{x}$

$$\Rightarrow y' = \frac{3}{2}\sqrt{x}$$

• $y = \sqrt{x} - 2e^x$

$$y' = \frac{1}{2\sqrt{x}} - 2e^x$$

Page 181

* قاعده : مجرد النظر :
انظر Page 6

$$(16) R(x) = \frac{\sqrt{10}}{x^7} \Rightarrow R'(x) = \frac{-7\sqrt{10}}{x^8}$$

$$(13) V(r) = \frac{4}{3}\pi r^3 \Rightarrow V'(r) = \frac{4}{3}\pi \cdot 3r^2 = \underline{\underline{4\pi r^2}}$$

$$(18) y = \sqrt[3]{x} \Rightarrow y = x^{\frac{1}{3}} \Rightarrow y' = \frac{1}{3}x^{-\frac{2}{3}} = \frac{1}{3x^{\frac{2}{3}}}$$

$$\Rightarrow y' = \frac{1}{3\sqrt[3]{x^2}}$$

$$(20) F(t) = \sqrt{t} - \frac{1}{\sqrt{t}}$$

$$\Rightarrow F(t) = \sqrt{t} - t^{\frac{1}{2}}$$

$$\Rightarrow F'(t) = \frac{1}{2\sqrt{t}} - (-\frac{1}{2})t^{-\frac{3}{2}} = \frac{1}{2\sqrt{t}} + \frac{1}{2t^{\frac{3}{2}}}$$

$$= \frac{1}{2\sqrt{t}} + \frac{1}{2t\sqrt{t}}$$

$$t^{\frac{3}{2}} = t\sqrt{t}$$

$$(28) y = a e^v + \frac{b}{v} + \frac{c}{v^2}$$

$$y' = a e^v + \frac{-b}{v^2} + \frac{-2c}{v^3} = a e^v - \frac{b}{v^2} - \frac{2c}{v^3}$$

$$(31) \quad z = \frac{A}{y^{10}} + B e^y$$

$$z' = \frac{-10A}{y^{11}} + B e^y$$

$$(32) \quad y = e^{x+1} + 1$$

$$y' = e^{x+1} \cdot \frac{1}{\text{الأس}} = e^{x+1}$$

Page 188

Find the equation of the tangent line and the normal line to (33) $y = \underline{2x} \underline{e^x}$ at $(\underset{\downarrow}{0}, \underset{\downarrow}{0})$
 x_1 y_1

$$y' = 2 \cdot e^x + e^x \cdot 2x$$

$$m = 2e^0 + e^0 \cdot 2(0) \Rightarrow m = 2e^0 = 2(1) = \boxed{2}$$

• eq. of tangent line: $y = m(x - x_1) + y_1$

$$y = 2(x - 0) + 0$$

$$\boxed{y = 2x}$$

• eq. of normal line: $y = -\frac{1}{m}(x - x_1) + y_1$

$$y = -\frac{1}{2}(x - 0) + 0$$

$$\Rightarrow \boxed{y = -\frac{1}{2}x}$$

(32) $y = \frac{e^x}{x}$ at $(1, e)$

$y' = \frac{e^x \cdot x - 1 \cdot e^x}{x^2}$

 (Note: e^x derivative is e^x , x derivative is 1)

لايجاد m
عوضه عن x بـ 1
عن y بـ e

$$m = \frac{e \cdot 1 - 1 \cdot e}{1^2} = e - e = 0$$

سادله اعلم
eq. of tangent line : $y = m(x - x_1) + y_1$


النقطة $(1, e)$
 معادله الخط $y = e$
 معادله العمود $x = 1$

$$y = 0(x - 1) + e$$

$$y = e$$

معادله العمود
eq. of normal line

$$x = 1$$



(23) $f(x) = \frac{A}{B + ce^x} \Rightarrow \frac{0 \cdot (B + ce^x) - ce^x \cdot A}{(B + ce^x)^2}$

$$\Rightarrow f'(x) = \frac{-Ace^x}{(B + ce^x)^2}$$

Page 181

(51) find the points on the curve

$$y = 2x^3 + 3x^2 - 12x + 1$$

where the tangent is horizontal.

المماس أفقياً
 أفقياً
 ∴ المماس أفقياً
 ⇒ ∴ $y' = 0$

$$6x^2 + 6x - 12 = 0 \quad \text{حل بواسطة التحليل} \quad (\div 6)$$

$$x + x - 2 = 0$$

$$(x + 2)(x - 1) = 0$$

$$x + 2 = 0$$

$$x = -2$$

لإيجاد y نعوض $x = -2$ في الدالة الأصلية

$$\begin{aligned} y &= 2(-2)^3 + 3(-2)^2 - 12(-2) + 1 \\ &= -16 + 12 + 24 + 1 \\ &= 21 \end{aligned}$$

$$x - 1 = 0$$

$$x = 1$$

لإيجاد y نعوض $x = 1$ في الدالة الأصلية

$$\begin{aligned} y &= 2(1)^3 + 3(1)^2 - 12(1) + 1 \\ &= 2 + 3 - 12 + 1 \\ &= -6 \end{aligned}$$

∴ The tangent is horizontal

at the points: $(-2, 21)$ and $(1, -6)$


Page 181

(50) The equation of motion of a particle is :

$$s = t^3 - 3t$$

where s in meters and t in seconds

(a) Find the velocity and acceleration.



* velocity: $v = \frac{ds}{dt} = 3t^2 - 3$

* acceleration: $a = \frac{dv}{dt} = 6t$

(b) Find the acceleration after 1 second.

↳ $a(1) = 6(1) = 6$

$$x\sqrt{x} = x \cdot x^{\frac{1}{2}} = x^{\frac{3}{2}}$$

$$\Rightarrow \frac{d}{dx} x\sqrt{x} \rightarrow \frac{3}{2} x^{\frac{1}{2}} = \frac{3\sqrt{x}}{2}$$

54) Find an eq. of the tangent line

to the curve $y = x\sqrt{x}$

that is parallel to the line $y = 1 + 3x$

توازي

$$m_1 = m_2$$

→ ميل المستقيم الموازي له = ميل المماس للمنحنى
(مشتقة المستقيم) (مشتقة المنحنى)

$$\frac{3}{2}\sqrt{x} = 3 \quad (\text{بالضرب عن } \frac{2}{3} \text{ للتخلص من معامل الجذر})$$

$$\frac{2}{2} \cdot \frac{2}{2} \sqrt{x} = \frac{2}{2} \cdot 3 \Rightarrow \sqrt{x} = 2 \xrightarrow{\text{التربيع}} x = 4$$

لنعوض من معادله المنحنى للحصول على y

$$\hookrightarrow y = 4\sqrt{4} \rightarrow y = 8$$

∴ نقطة التماس (4 , 8)
الميل (slope) $m = 3$

→ eq. of tangent line :

$$y = m(x - x_1) + y_1$$

$$y = 3(x - 4) + 8$$

$$y = 3x - 12 + 8$$

$$y = 3x - 4$$

* للتأكد من صحة الحل
استخدم النقطة
(4, 8)

من المعادله الأضربه
عوضه بـ x = 4
يكون الناتج y = 8
مما يؤكد صحة الحل.

Page 182

(75) let $f(x) = \begin{cases} x^2 & \text{if } x \leq 2 \\ mx+b & \text{if } x > 2 \end{cases}$

find the values of m and b that make f differentiable every

قابله للارتقاء
معناه لا يوجد
انقطاع شريطة

المشتقة اليسرى = المشتقة اليمنى
عند $x=2$ عند $x=2$

$$m = 2(2)$$

$$\therefore m = 4$$

القيمة عند $x=2$

$$\lim_{x \rightarrow 2^+} (mx+b) = \lim_{x \rightarrow 2} x^2$$

$$4(2) + b = 4$$

$$8 + b = 4$$

$$b = 4 - 8$$

$$\therefore b = -4$$

* (1) If: $y = x^3 + 3(\pi^2 + x^2)$ Find y'' ?

* (2) If: $y = \sin^2 x + \cos^2 x$ Find y' ?

* (3) Find: $\lim_{x \rightarrow 1} \frac{x^{1000} - 1}{x - 1}$

المسألة

أفترض

● Suppose u and v

are differentiable functions where:

$$u(1) = 2 \quad , \quad u'(1) = 0$$

$$v(1) = 5 \quad , \quad v'(1) = -1$$

Find:

$$\textcircled{1} \quad \frac{d}{dx} (uv) = u' \cdot v + v' \cdot u$$

$$\text{at } (x=1) = 0 \cdot (5) + (-1) \cdot (2) = \boxed{-2}$$


$$\textcircled{2} \quad \frac{d}{dx} \left(\frac{u}{v} \right) = \frac{u' \cdot v - v' \cdot u}{v^2}$$

$$\text{at } (x=1) = \frac{(0)(5) - (-1)(2)}{(5)^2} = \boxed{\frac{2}{25}}$$

$$\textcircled{3} \quad \frac{d}{dx} (7v - 2u^2) = 7v' - 4uu'$$

$$\text{at } (x=1) = 7(-1) - 4(2)(0)$$

$$= -7 - 0 = \boxed{-7}$$



$$\bullet y = e^x - 3x^4$$

Find $y^{(5)}$?

* درجة هذا الحد (4 = degree)

رتبه المشتقة (5 = order)

∴ المشتقة الخامسة لهذا الحد zero

* أما e^x فمشتقتها دائماً e^x

لها كما كان عدد مرات الاشتقاق

$$\Rightarrow y^{(5)} = e^x - 0 = e^x$$

$$\bullet y = \frac{e^{3x} + e^{2x}}{e^{2x}}$$

Find y' ?* المقام يتكون من حد واحد فقط
∴ توزيع حدود البسط على نفس المقام ثم الاختصار ثم الاشتقاق

$$y = \frac{e^{3x}}{e^{2x}} + \frac{e^{2x}}{e^{2x}}$$

$$y = e^x + 1$$

$$y' = e^x + 0 \Rightarrow y' = e^x$$

$$\bullet y = x + \frac{1}{x}$$

Find y' ?

$$y' = 1 - \frac{1}{x^2}$$

$$\frac{1}{x} \text{ مشتقة } \\ -\frac{1}{x^2} \text{ هي}$$



THOMAS'
CALCULUS
MEDIA UPGRADE

Chapter 4

Applications of Derivatives

4.1

Extreme Values of Functions

DEFINITIONS Absolute Maximum, Absolute Minimum

Let f be a function with domain D . Then f has an **absolute maximum** value on D at a point c if

$$f(x) \leq f(c) \quad \text{for all } x \text{ in } D$$

and an **absolute minimum** value on D at c if

$$f(x) \geq f(c) \quad \text{for all } x \text{ in } D.$$







FIGURE 4.1 Absolute extrema for the sine and cosine functions on $[-\pi/2, \pi/2]$. These values can depend on the domain of a function.




(a) abs min only



(b) abs max and min



(c) abs max only




(d) no max or min


FIGURE 4.2 Graphs for Example 1.

THEOREM 1 **The Extreme Value Theorem**


If f is continuous on a closed interval $[a, b]$, then f attains both an absolute maximum value M and an absolute minimum value m in $[a, b]$. That is, there are numbers x_1 and x_2 in $[a, b]$ with $f(x_1) = m$, $f(x_2) = M$, and $m \leq f(x) \leq M$ for every other x in $[a, b]$ (Figure 4.3).




Maximum and minimum
at interior points



Maximum and minimum
at endpoints



Maximum at interior point,
minimum at endpoint



Minimum at interior point,
maximum at endpoint

FIGURE 4.3 Some possibilities for a continuous function's maximum and minimum on a closed interval $[a, b]$.




FIGURE 4.4 Even a single point of discontinuity can keep a function from having either a maximum or minimum value on a closed interval. The function

$$y = \begin{cases} x, & 0 \leq x < 1 \\ 0, & x = 1 \end{cases}$$

is continuous at every point of $[0, 1]$ except $x = 1$, yet its graph over $[0, 1]$ does not have a highest point.

DEFINITIONS Local Maximum, Local Minimum

A function f has a **local maximum** value at an interior point c of its domain if

$$f(x) \leq f(c) \quad \text{for all } x \text{ in some open interval containing } c.$$

A function f has a **local minimum** value at an interior point c of its domain if

$$f(x) \geq f(c) \quad \text{for all } x \text{ in some open interval containing } c.$$




FIGURE 4.5 How to classify maxima and minima.

THEOREM 2 The First Derivative Theorem for Local Extreme Values

If f has a local maximum or minimum value at an interior point c of its domain, and if f' is defined at c , then

$$f'(c) = 0.$$




FIGURE 4.6 A curve with a local maximum value. The slope at c , simultaneously the limit of nonpositive numbers and nonnegative numbers, is zero.

DEFINITION **Critical Point**

An interior point of the domain of a function f where f' is zero or undefined is a **critical point** of f .

How to Find the Absolute Extrema of a Continuous Function f on a Finite Closed Interval

1. Evaluate f at all critical points and endpoints.
2. Take the largest and smallest of these values.




FIGURE 4.7 The extreme values of $g(t) = 8t - t^4$ on $[-2, 1]$ (Example 3).




FIGURE 4.8 The extreme values of $f(x) = x^{2/3}$ on $[-2, 3]$ occur at $x = 0$ and $x = 3$ (Example 4).





FIGURE 4.9 Critical points without extreme values. (a) $y' = 3x^2$ is 0 at $x = 0$, but $y = x^3$ has no extremum there. (b) $y' = (1/3)x^{-2/3}$ is undefined at $x = 0$, but $y = x^{1/3}$ has no extremum there.

4.2

The Mean Value Theorem


THEOREM 3 **Rolle's Theorem**

Suppose that $y = f(x)$ is continuous at every point of the closed interval $[a, b]$ and differentiable at every point of its interior (a, b) . If


$$f(a) = f(b),$$

then there is at least one number c in (a, b) at which

$$f'(c) = 0.$$




(a)




(b)


FIGURE 4.10 Rolle's Theorem says that a differentiable curve has at least one horizontal tangent between any two points where it crosses a horizontal line. It may have just one (a), or it may have more (b).



(a) Discontinuous at an endpoint of $[a, b]$



(b) Discontinuous at an interior point of $[a, b]$



(c) Continuous on $[a, b]$ but not differentiable at an interior point

FIGURE 4.11 There may be no horizontal tangent if the hypotheses of Rolle's Theorem do not hold.




FIGURE 4.12 As predicted by Rolle's Theorem, this curve has horizontal tangents between the points where it crosses the x -axis (Example 1).




FIGURE 4.13 The only real zero of the polynomial $y = x^3 + 3x + 1$ is the one shown here where the curve crosses the x -axis between -1 and 0 (Example 2).

THEOREM 4 The Mean Value Theorem

Suppose $y = f(x)$ is continuous on a closed interval $[a, b]$ and differentiable on the interval's interior (a, b) . Then there is at least one point c in (a, b) at which

$$\frac{f(b) - f(a)}{b - a} = f'(c). \quad (1)$$




FIGURE 4.14 Geometrically, the Mean Value Theorem says that somewhere between A and B the curve has at least one tangent parallel to chord AB .




FIGURE 4.15 The graph of f and the chord AB over the interval $[a, b]$.




FIGURE 4.16 The chord AB is the graph of the function $g(x)$. The function $h(x) = f(x) - g(x)$ gives the vertical distance between the graphs of f and g at x .




FIGURE 4.17 The function $f(x) = \sqrt{1 - x^2}$ satisfies the hypotheses (and conclusion) of the Mean Value Theorem on $[-1, 1]$ even though f is not differentiable at -1 and 1 .




FIGURE 4.18 As we find in Example 3, $c = 1$ is where the tangent is parallel to the chord.




FIGURE 4.19 Distance versus elapsed time for the car in Example 4.

COROLLARY 1 **Functions with Zero Derivatives Are Constant**

If $f'(x) = 0$ at each point x of an open interval (a, b) , then $f(x) = C$ for all $x \in (a, b)$, where C is a constant.

COROLLARY 2 **Functions with the Same Derivative Differ by a Constant**

If $f'(x) = g'(x)$ at each point x in an open interval (a, b) , then there exists a constant C such that $f(x) = g(x) + C$ for all $x \in (a, b)$. That is, $f - g$ is a constant on (a, b) .




FIGURE 4.20 From a geometric point of view, Corollary 2 of the Mean Value Theorem says that the graphs of functions with identical derivatives on an interval can differ only by a vertical shift there. The graphs of the functions with derivative $2x$ are the parabolas $y = x^2 + C$, shown here for selected values of C .

4.3

Monotonic Functions and The First Derivative Test

DEFINITIONS Increasing, Decreasing Function

Let f be a function defined on an interval I and let x_1 and x_2 be any two points in I .

1. If $f(x_1) < f(x_2)$ whenever $x_1 < x_2$, then f is said to be **increasing** on I .
2. If $f(x_2) < f(x_1)$ whenever $x_1 < x_2$, then f is said to be **decreasing** on I .

A function that is increasing or decreasing on I is called **monotonic** on I .

COROLLARY 3 **First Derivative Test for Monotonic Functions**

Suppose that f is continuous on $[a, b]$ and differentiable on (a, b) .

If $f'(x) > 0$ at each point $x \in (a, b)$, then f is increasing on $[a, b]$.

If $f'(x) < 0$ at each point $x \in (a, b)$, then f is decreasing on $[a, b]$.




FIGURE 4.21 The function $f(x) = x^2$ is monotonic on the intervals $(-\infty, 0]$ and $[0, \infty)$, but it is not monotonic on $(-\infty, \infty)$.




FIGURE 4.22 The function $f(x) = x^3 - 12x - 5$ is monotonic on three separate intervals (Example 1).

First Derivative Test for Local Extrema

Suppose that c is a critical point of a continuous function f , and that f is differentiable at every point in some interval containing c except possibly at c itself. Moving across c from left to right,

1. if f' changes from negative to positive at c , then f has a local minimum at c ;
2. if f' changes from positive to negative at c , then f has a local maximum at c ;
3. if f' does not change sign at c (that is, f' is positive on both sides of c or negative on both sides), then f has no local extremum at c .




FIGURE 4.23 A function's first derivative tells how the graph rises and falls.




FIGURE 4.24 The function $f(x) = x^{1/3}(x - 4)$ decreases when $x < 1$ and increases when $x > 1$ (Example 2).

4.4

Concavity and Curve Sketching

DEFINITION **Concave Up, Concave Down**

The graph of a differentiable function $y = f(x)$ is

- (a) **concave up** on an open interval I if f' is increasing on I
- (b) **concave down** on an open interval I if f' is decreasing on I .




FIGURE 4.25 The graph of $f(x) = x^3$ is concave down on $(-\infty, 0)$ and concave up on $(0, \infty)$ (Example 1a).

The Second Derivative Test for Concavity

Let $y = f(x)$ be twice-differentiable on an interval I .

1. If $f'' > 0$ on I , the graph of f over I is concave up.
2. If $f'' < 0$ on I , the graph of f over I is concave down.




FIGURE 4.26 The graph of $f(x) = x^2$ is concave up on every interval (Example 1b).

DEFINITION **Point of Inflection**

A point where the graph of a function has a tangent line and where the concavity changes is a **point of inflection**.




FIGURE 4.27 Using the graph of y'' to determine the concavity of y (Example 2).




FIGURE 4.28 The graph of $y = x^4$ has no inflection point at the origin, even though $y'' = 0$ there (Example 3).





FIGURE 4.29 A point where y'' fails to exist can be a point of inflection (Example 4).

THEOREM 5 **Second Derivative Test for Local Extrema**

Suppose f'' is continuous on an open interval that contains $x = c$.

1. If $f'(c) = 0$ and $f''(c) < 0$, then f has a local maximum at $x = c$.
2. If $f'(c) = 0$ and $f''(c) > 0$, then f has a local minimum at $x = c$.
3. If $f'(c) = 0$ and $f''(c) = 0$, then the test fails. The function f may have a local maximum, a local minimum, or neither.



$$f' = 0, f'' < 0$$
$$\Rightarrow \text{local max}$$



$$f' = 0, f'' > 0$$
$$\Rightarrow \text{local min}$$




FIGURE 4.30 The graph of $f(x) = x^4 - 4x^3 + 10$ (Example 6).

Strategy for Graphing $y = f(x)$

1. Identify the domain of f and any symmetries the curve may have.
2. Find y' and y'' .
3. Find the critical points of f , and identify the function's behavior at each one.
4. Find where the curve is increasing and where it is decreasing.
5. Find the points of inflection, if any occur, and determine the concavity of the curve.
6. Identify any asymptotes.
7. Plot key points, such as the intercepts and the points found in Steps 3–5, and sketch the curve.














FIGURE 4.31 The graph of $y = \frac{(x + 1)^2}{1 + x^2}$
(Example 7).


 <p>$y = f(x)$</p> <p>Differentiable \Rightarrow smooth, connected; graph may rise and fall</p>	 <p>$y = f(x)$</p> <p>$y' > 0 \Rightarrow$ rises from left to right; may be wavy</p>	 <p>$y = f(x)$</p> <p>$y' < 0 \Rightarrow$ falls from left to right; may be wavy</p>
 <p>or</p> <p>$y'' > 0 \Rightarrow$ concave up throughout; no waves; graph may rise or fall</p>	 <p>or</p> <p>$y'' < 0 \Rightarrow$ concave down throughout; no waves; graph may rise or fall</p>	 <p>y'' changes sign</p> <p>Inflection point</p>
 <p>or</p> <p>y' changes sign \Rightarrow graph has local maximum or local minimum</p>	 <p>$y' = 0$ and $y'' < 0$ at a point; graph has local maximum</p>	 <p>$y' = 0$ and $y'' > 0$ at a point; graph has local minimum</p>

4.5

Applied Optimization Problems



(a)



(b)

FIGURE 4.32 An open box made by cutting the corners from a square sheet of tin. What size corners maximize the box's volume (Example 1)?




FIGURE 4.33 The volume of the box in Figure 4.32 graphed as a function of x .





FIGURE 4.34 This 1-L can uses the least material when $h = 2r$ (Example 2).



Tall and thin



Short and wide




FIGURE 4.35 The graph of $A = 2\pi r^2 + 2000/r$ is concave up.




FIGURE 4.36 The rectangle inscribed in the semicircle in Example 3.




FIGURE 4.37 A light ray refracted (deflected from its path) as it passes from one medium to a denser medium (Example 4).




FIGURE 4.38 The sign pattern of dt/dx in Example 4.




FIGURE 4.39 The graph of a typical cost function starts concave down and later turns concave up. It crosses the revenue curve at the break-even point B . To the left of B , the company operates at a loss. To the right, the company operates at a profit, with the maximum profit occurring where $c'(x) = r'(x)$. Farther to the right, cost exceeds revenue (perhaps because of a combination of rising labor and material costs and market saturation) and production levels become unprofitable again.




FIGURE 4.40 The cost and revenue curves for Example 5.




FIGURE 4.41 The average daily cost $c(x)$ is the sum of a hyperbola and a linear function (Example 6).

4.6

Indeterminate Forms and L' Hôpital's Rule

THEOREM 6 **L'Hôpital's Rule (First Form)**

Suppose that $f(a) = g(a) = 0$, that $f'(a)$ and $g'(a)$ exist, and that $g'(a) \neq 0$.
Then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{f'(a)}{g'(a)}.$$

THEOREM 7 L'Hôpital's Rule (Stronger Form)

Suppose that $f(a) = g(a) = 0$, that f and g are differentiable on an open interval I containing a , and that $g'(x) \neq 0$ on I if $x \neq a$. Then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)},$$

assuming that the limit on the right side exists.

THEOREM 8 **Cauchy's Mean Value Theorem**

Suppose functions f and g are continuous on $[a, b]$ and differentiable throughout (a, b) and also suppose $g'(x) \neq 0$ throughout (a, b) . Then there exists a number c in (a, b) at which

$$\frac{f'(c)}{g'(c)} = \frac{f(b) - f(a)}{g(b) - g(a)}.$$




FIGURE 4.42 There is at least one value of the parameter $t = c$, $a < c < b$, for which the slope of the tangent to the curve at $(g(c), f(c))$ is the same as the slope of the secant line joining the points $(g(a), f(a))$ and $(g(b), f(b))$.

Using L'Hôpital's Rule

To find

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$$

by l'Hôpital's Rule, continue to differentiate f and g , so long as we still get the form $0/0$ at $x = a$. But as soon as one or the other of these derivatives is different from zero at $x = a$ we stop differentiating. L'Hôpital's Rule does not apply when either the numerator or denominator has a finite nonzero limit.

4.7

Newton's Method

Procedure for Newton's Method

1. Guess a first approximation to a solution of the equation $f(x) = 0$. A graph of $y = f(x)$ may help.
2. Use the first approximation to get a second, the second to get a third, and so on, using the formula

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}, \quad \text{if } f'(x_n) \neq 0 \quad (1)$$




FIGURE 4.43 Newton's method starts with an initial guess x_0 and (under favorable circumstances) improves the guess one step at a time.




FIGURE 4.44 The geometry of the successive steps of Newton's method. From x_n we go up to the curve and follow the tangent line down to find x_{n+1} .




FIGURE 4.45 The graph of $f(x) = x^3 - x - 1$ crosses the x -axis once; this is the root we want to find (Example 2).

TABLE 4.1 The result of applying Newton's method to $f(x) = x^3 - x - 1$ with $x_0 = 1$

n	x_n	$f(x_n)$	$f'(x_n)$	$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$
0	1	-1	2	1.5
1	1.5	0.875	5.75	1.3478 26087
2	1.3478 26087	0.1006 82173	4.4499 05482	1.3252 00399
3	1.3252 00399	0.0020 58362	4.2684 68292	1.3247 18174
4	1.3247 18174	0.0000 00924	4.2646 34722	1.3247 17957
5	1.3247 17957	-1.8672E-13	4.2646 32999	1.3247 17957




FIGURE 4.46 The first three x -values in Table 4.1 (four decimal places).




FIGURE 4.47 Any starting value x_0 to the right of $x = 1/\sqrt{3}$ will lead to the root.




FIGURE 4.48 Newton's method will converge to r from either starting point.




FIGURE 4.49 If $f'(x_n) = 0$, there is no intersection point to define x_{n+1} .




FIGURE 4.50 Newton's method fails to converge. You go from x_0 to x_1 and back to x_0 , never getting any closer to r .





FIGURE 4.51 If you start too far away, Newton's method may miss the root you want.




FIGURE 4.52 (a) Starting values in $(-\infty, -\sqrt{2}/2)$, $(-\sqrt{21}/7, \sqrt{21}/7)$, and $(\sqrt{2}/2, \infty)$ lead respectively to roots A, B, and C. (b) The values $x = \pm\sqrt{21}/7$ lead only to each other. (c) Between $\sqrt{21}/7$ and $\sqrt{2}/2$, there are infinitely many open intervals of points attracted to A alternating with open intervals of points attracted to C. This behavior is mirrored in the interval $(-\sqrt{2}/2, -\sqrt{21}/7)$.




FIGURE 4.53 This computer-generated initial value portrait uses color to show where different points in the complex plane end up when they are used as starting values in applying Newton's method to solve the equation $z^6 - 1 = 0$. Red points go to 1, green points to $(1/2) + (\sqrt{3}/2)i$, dark blue points to $(-1/2) + (\sqrt{3}/2)i$, and so on. Starting values that generate sequences that do not arrive within 0.1 unit of a root after 32 steps are colored black.

4.8

Antiderivatives

DEFINITION **Antiderivative**

A function F is an **antiderivative** of f on an interval I if $F'(x) = f(x)$ for all x in I .

If F is an antiderivative of f on an interval I , then the most general antiderivative of f on I is

$$F(x) + C$$

where C is an arbitrary constant.

TABLE 4.2 Antiderivative formulas

	Function	General antiderivative
1.	x^n	$\frac{x^{n+1}}{n+1} + C, \quad n \neq -1, n \text{ rational}$
2.	$\sin kx$	$-\frac{\cos kx}{k} + C, \quad k \text{ a constant, } k \neq 0$
3.	$\cos kx$	$\frac{\sin kx}{k} + C, \quad k \text{ a constant, } k \neq 0$
4.	$\sec^2 x$	$\tan x + C$
5.	$\csc^2 x$	$-\cot x + C$
6.	$\sec x \tan x$	$\sec x + C$
7.	$\csc x \cot x$	$-\csc x + C$

TABLE 4.3 Antiderivative linearity rules

	Function	General antiderivative
1.	<i>Constant Multiple Rule:</i> $kf(x)$	$kF(x) + C$, k a constant
2.	<i>Negative Rule:</i> $-f(x)$	$-F(x) + C$,
3.	<i>Sum or Difference Rule:</i> $f(x) \pm g(x)$	$F(x) \pm G(x) + C$




FIGURE 4.54 The curves $y = x^3 + C$ fill the coordinate plane without overlapping. In Example 5, we identify the curve $y = x^3 - 2$ as the one that passes through the given point $(1, -1)$.

DEFINITION Indefinite Integral, Integrand

The set of all antiderivatives of f is the **indefinite integral** of f with respect to x , denoted by

$$\int f(x) dx.$$

The symbol \int is an **integral sign**. The function f is the **integrand** of the integral, and x is the **variable of integration**.

Workshop Solutions to Sections 5.1 and 5.2

<p>1) The absolute maximum value of $f(x) = x^3 - 2x^2$ in $[-1, 2]$ is at $x =$</p> <p><u>Solution:</u> Since $f(x)$ is a continuous on $[-1, 2]$, we can use the Closed Interval Method,</p> $f(x) = x^3 - 2x^2$ $f'(x) = 3x^2 - 4x$ <p>Now, we find the critical numbers of $f(x)$ when</p> $f'(x) = 0 \Rightarrow 3x^2 - 4x = 0 \Rightarrow x(3x - 4) = 0$ $\Rightarrow x = 0 \text{ or } x = \frac{4}{3}$ <p>Thus,</p> $f(-1) = (-1)^3 - 2(-1)^2 = -1 - 2 = -3$ $f(2) = (2)^3 - 2(2)^2 = 8 - 8 = 0$ $f(0) = (0)^3 - 2(0)^2 = 0 - 0 = 0$ $f\left(\frac{4}{3}\right) = \left(\frac{4}{3}\right)^3 - 2\left(\frac{4}{3}\right)^2 = \frac{64}{27} - \frac{32}{9} = -\frac{32}{27}$ <p>Hence, we see that the absolute maximum value is 0 at $x = 0$ and $x = 2$</p>	<p>2) The absolute minimum value of $f(x) = x^3 - 3x^2 + 1$ in $\left[-\frac{1}{2}, 4\right]$ is</p> <p><u>Solution:</u> Since $f(x)$ is a continuous on $\left[-\frac{1}{2}, 4\right]$, we can use the Closed Interval Method,</p> $f(x) = x^3 - 3x^2 + 1$ $f'(x) = 3x^2 - 6x$ <p>Now, we find the critical numbers of $f(x)$ when</p> $f'(x) = 0 \Rightarrow 3x^2 - 6x = 0 \Rightarrow 3x(x - 2) = 0$ $\Rightarrow x = 0 \text{ or } x = 2$ <p>Thus,</p> $f\left(-\frac{1}{2}\right) = \left(-\frac{1}{2}\right)^3 - 3\left(-\frac{1}{2}\right)^2 + 1 = -\frac{1}{8} - \frac{3}{4} + 1 = \frac{1}{8}$ $f(4) = (4)^3 - 3(4)^2 + 1 = 64 - 48 + 1 = 17$ $f(0) = (0)^3 - 3(0)^2 + 1 = 0 - 0 + 1 = 1$ $f(2) = (2)^3 - 3(2)^2 + 1 = 8 - 12 + 1 = -3$ <p>Hence, we see that the absolute minimum value is -3 at $x = 2$</p>
<p>3) The absolute maximum point of $f(x) = 3x^2 - 12x + 1$ in $[0, 3]$ is</p> <p><u>Solution:</u> Since $f(x)$ is a continuous on $[0, 3]$, we can use the Closed Interval Method,</p> $f(x) = 3x^2 - 12x + 1$ $f'(x) = 6x - 12$ <p>Now, we find the critical numbers of $f(x)$ when</p> $f'(x) = 0 \Rightarrow 6x - 12 = 0 \Rightarrow 6x = 12$ $\Rightarrow x = 2$ <p>Thus,</p> $f(0) = 3(0)^2 - 12(0) + 1 = 0 - 0 + 1 = 1$ $f(3) = 3(3)^2 - 12(3) + 1 = 27 - 36 + 1 = -8$ $f(2) = 3(2)^2 - 12(2) + 1 = 12 - 24 + 1 = -11$ <p>Hence, we see that the absolute maximum point is $(0, 1)$.</p>	<p>4) The absolute minimum point of $f(x) = 3x^2 - 12x + 1$ in $[0, 3]$ is</p> <p><u>Solution:</u> Since $f(x)$ is a continuous on $[0, 3]$, we can use the Closed Interval Method,</p> $f(x) = 3x^2 - 12x + 1$ $f'(x) = 6x - 12$ <p>Now, we find the critical numbers of $f(x)$ when</p> $f'(x) = 0 \Rightarrow 6x - 12 = 0 \Rightarrow 6x = 12$ $\Rightarrow x = 2$ <p>Thus,</p> $f(0) = 3(0)^2 - 12(0) + 1 = 0 - 0 + 1 = 1$ $f(3) = 3(3)^2 - 12(3) + 1 = 27 - 36 + 1 = -8$ $f(2) = 3(2)^2 - 12(2) + 1 = 12 - 24 + 1 = -11$ <p>Hence, we see that the absolute minimum point is $(2, -11)$.</p>
<p>5) The absolute minimum point of $f(x) = 3x^2 - 12x + 2$ in $[0, 3]$ is</p> <p><u>Solution:</u> Since $f(x)$ is a continuous on $[0, 3]$, we can use the Closed Interval Method,</p> $f(x) = 3x^2 - 12x + 2$ $f'(x) = 6x - 12$ <p>Now, we find the critical numbers of $f(x)$ when</p> $f'(x) = 0 \Rightarrow 6x - 12 = 0 \Rightarrow 6x = 12$ $\Rightarrow x = 2$ <p>Thus,</p> $f(0) = 3(0)^2 - 12(0) + 2 = 0 - 0 + 2 = 2$ $f(3) = 3(3)^2 - 12(3) + 2 = 27 - 36 + 2 = -7$ $f(2) = 3(2)^2 - 12(2) + 2 = 12 - 24 + 2 = -10$ <p>Hence, we see that the absolute minimum point is $(2, -10)$.</p>	<p>6) The values in $(-3, 3)$ which make $f(x) = x^3 - 9x$ satisfy Rolle's Theorem on $[-3, 3]$ are</p> <p><u>Solution:</u></p> <ul style="list-style-type: none"> $\because f(x)$ is a polynomial, then 1- $f(x)$ is a continuous on $[-3, 3]$. 2- $f(x)$ is differentiable on $(-3, 3)$, $f'(x) = 3x^2 - 9$ 3- $f(-3) = (-3)^3 - 9(-3) = -27 + 27 = 0 = f(3)$ <p>Then there is a number $c \in (-3, 3)$ such that</p> $f'(c) = 0 \Rightarrow 3c^2 - 9 = 0 \Rightarrow 3c^2 = 9$ $\Rightarrow c^2 = 3 \Rightarrow c = \pm\sqrt{3}$ <p>Hence, the values are $\pm\sqrt{3} \in (-3, 3)$.</p>

7) The values in $(0,2)$ which make $f(x) = x^3 - 3x^2 + 2x + 5$ satisfy Rolle's Theorem on $[0,2]$ are

Solution:

$\because f(x)$ is a polynomial, then

1- $f(x)$ is a continuous on $[0,2]$.

2- $f(x)$ is differentiable on $(0,2)$,

$$f'(x) = 3x^2 - 6x + 2$$

3- $f(0) = (0)^3 - 3(0)^2 + 2(0) + 5 = 5 = f(2)$

Then there is a number $c \in (0,2)$ such that

$$f'(c) = 0 \Rightarrow 3c^2 - 6c + 2 = 0$$

$$\begin{aligned} \Rightarrow c &= \frac{-(-6) \pm \sqrt{(-6)^2 - 4(3)(2)}}{2(3)} = \frac{6 \pm \sqrt{36 - 24}}{6} \\ &= \frac{6 \pm \sqrt{12}}{6} = \frac{6 \pm \sqrt{3} \times 2}{6} = \frac{6 \pm 2\sqrt{3}}{6} \\ &= \frac{2(3 \pm \sqrt{3})}{6} = \frac{3 \pm \sqrt{3}}{3} = \frac{3}{3} \pm \frac{\sqrt{3}}{3} \\ &= 1 \pm \frac{\sqrt{3}}{3} \end{aligned}$$

Hence, the values are $1 \pm \frac{\sqrt{3}}{3} \in (0,2)$.

8) The value c in $(0,5)$ which makes $f(x) = x^2 - x - 6$ satisfy the Mean Value Theorem on $[0,5]$ is

Solution:

$\because f(x)$ is a polynomial, then

1- $f(x)$ is a continuous on $[0,5]$.

2- $f(x)$ is differentiable on $(0,5)$,

$$f'(x) = 2x - 1$$

Then there is a number $c \in (0,5)$ such that

$$\begin{aligned} f'(c) &= \frac{f(5) - f(0)}{5 - 0} \\ \Rightarrow 2c - 1 &= \frac{[(5)^2 - (5) - 6] - [(0)^2 - (0) - 6]}{5} \\ \Rightarrow 2c - 1 &= \frac{(14) - (-6)}{5} \\ \Rightarrow 2c - 1 &= \frac{14 + 6}{5} \\ \Rightarrow 2c - 1 &= 4 \\ \Rightarrow 2c &= 4 + 1 \\ \Rightarrow c &= \frac{5}{2} \end{aligned}$$

Hence, the value c is $\frac{5}{2} \in (0,5)$.

9) The value c in $(0,2)$ makes $f(x) = x^3 - x$ satisfied the Mean Value Theorem on $[0,2]$ are

Solution:

$\because f(x)$ is a polynomial, then

1- $f(x)$ is a continuous on $[0,2]$.

2- $f(x)$ is differentiable on $(0,2)$,

$$f'(x) = 3x^2 - 1$$

Then there is a number $c \in (0,2)$ such that

$$\begin{aligned} f'(c) &= \frac{f(2) - f(0)}{2 - 0} \\ \Rightarrow 3c^2 - 1 &= \frac{[(2)^3 - (2)] - [(0)^3 - (0)]}{2} \\ \Rightarrow 3c^2 - 1 &= \frac{(6) - (0)}{2} \\ \Rightarrow 3c^2 - 1 &= \frac{6}{2} \\ \Rightarrow 3c^2 - 1 &= 3 \\ \Rightarrow 3c^2 &= 3 + 1 \\ \Rightarrow c^2 &= \frac{4}{3} \\ \Rightarrow c &= \pm \sqrt{\frac{4}{3}} \\ \Rightarrow c &= \pm \frac{2}{\sqrt{3}} \end{aligned}$$

Hence, the value c is $\frac{2}{\sqrt{3}} \in (0,2)$ but $-\frac{2}{\sqrt{3}} \notin (0,2)$.

10) The value in $(0,1)$ which makes $f(x) = 3x^2 + 2x + 5$ satisfy the Mean Value Theorem on $[0,1]$ is

Solution:

$\because f(x)$ is a polynomial, then

1- $f(x)$ is a continuous on $[0,1]$.

2- $f(x)$ is differentiable on $(0,1)$,

$$f'(x) = 6x + 2$$

Then there is a number $c \in (0,1)$ such that

$$\begin{aligned} f'(c) &= \frac{f(1) - f(0)}{1 - 0} \\ \Rightarrow 6c + 2 &= \frac{[3(1)^2 + 2(1) + 5] - [3(0)^2 + 2(0) + 5]}{1} \\ \Rightarrow 6c + 2 &= (3 + 2 + 5) - (0 + 0 + 5) \\ \Rightarrow 6c + 2 &= 10 - 5 \\ \Rightarrow 6c + 2 &= 5 \\ \Rightarrow 6c &= 5 - 2 \\ \Rightarrow 6c &= 3 \\ \Rightarrow c &= \frac{3}{6} \\ \Rightarrow c &= \frac{1}{2} \end{aligned}$$

Hence, the values are $\frac{1}{2} \in (0,1)$.

11) The critical numbers of the function

$$f(x) = x^3 + 3x^2 - 9x + 1 \text{ are}$$

Solution:

$$\begin{aligned} f'(x) &= 3x^2 + 6x - 9 \\ f'(x) = 0 &\Rightarrow 3x^2 + 6x - 9 = 0 \\ &\Rightarrow 3(x^2 + 2x - 3) = 0 \\ &\Rightarrow x^2 + 2x - 3 = 0 \\ &\Rightarrow (x + 3)(x - 1) = 0 \\ &\Rightarrow x = -3 \text{ or } x = 1 \end{aligned}$$

12) The function $f(x) = x^3 + 3x^2 - 9x + 1$ is decreasing on

Solution:

$$\begin{aligned}
 f'(x) &= 3x^2 + 6x - 9 \\
 f'(x) = 0 &\Rightarrow 3x^2 + 6x - 9 = 0 \\
 &\Rightarrow 3(x^2 + 2x - 3) = 0 \\
 &\Rightarrow x^2 + 2x - 3 = 0 \\
 &\Rightarrow (x + 3)(x - 1) = 0 \\
 &\Rightarrow x = -3 \text{ or } x = 1 \\
 &\quad -3 \qquad \qquad 1
 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity
			y

Hence, the function $f(x)$ is decreasing on $(-3, 1)$

13) The function $f(x) = x^3 + 3x^2 - 9x + 1$ is increasing on

Solution:

$$\begin{aligned}
 f'(x) &= 3x^2 + 6x - 9 \\
 f'(x) = 0 &\Rightarrow 3x^2 + 6x - 9 = 0 \\
 &\Rightarrow 3(x^2 + 2x - 3) = 0 \\
 &\Rightarrow x^2 + 2x - 3 = 0 \\
 &\Rightarrow (x + 3)(x - 1) = 0 \\
 &\Rightarrow x = -3 \text{ or } x = 1 \\
 &\quad -3 \qquad \qquad 1
 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity
			y

Hence, the function $f(x)$ is increasing on $(-\infty, -3) \cup (1, \infty)$

14) The function $f(x) = x^3 + 3x^2 - 9x + 1$ has a relative maximum value at the point

Solution:

$$\begin{aligned}
 f'(x) &= 3x^2 + 6x - 9 \\
 f'(x) = 0 &\Rightarrow 3x^2 + 6x - 9 = 0 \\
 &\Rightarrow 3(x^2 + 2x - 3) = 0 \\
 &\Rightarrow x^2 + 2x - 3 = 0 \\
 &\Rightarrow (x + 3)(x - 1) = 0 \\
 &\Rightarrow x = -3 \text{ or } x = 1 \\
 &\quad -3 \qquad \qquad 1
 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity
			y

Hence, the function $f(x)$ has a relative maximum value at the point $(-3, 28)$.

$$\begin{aligned}
 f(-3) &= (-3)^3 + 3(-3)^2 - 9(-3) + 1 \\
 &= -27 + 27 + 27 + 1 = 28
 \end{aligned}$$

15) The function $f(x) = x^3 + 3x^2 - 9x + 1$ has a relative minimum value at the point

Solution:

$$\begin{aligned}
 f'(x) &= 3x^2 + 6x - 9 \\
 f'(x) = 0 &\Rightarrow 3x^2 + 6x - 9 = 0 \\
 &\Rightarrow 3(x^2 + 2x - 3) = 0 \\
 &\Rightarrow x^2 + 2x - 3 = 0 \\
 &\Rightarrow (x + 3)(x - 1) = 0 \\
 &\Rightarrow x = -3 \text{ or } x = 1 \\
 &\quad -3 \qquad \qquad 1
 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity
			y

Hence, the function $f(x)$ has a relative minimum value at the point $(1, -4)$.

$$\begin{aligned}
 f(1) &= (1)^3 + 3(1)^2 - 9(1) + 1 \\
 &= 1 + 3 - 9 + 1 = -4
 \end{aligned}$$

16) The function $f(x) = x^3 + 3x^2 - 9x + 1$ concave upward on

Solution:

$$\begin{aligned}
 f'(x) &= 3x^2 + 6x - 9 \\
 f''(x) &= 6x + 6 \\
 f''(x) = 0 &\Rightarrow 6x + 6 = 0 \\
 &\Rightarrow 6x = -6 \\
 &\Rightarrow x = -\frac{6}{6} \\
 &\Rightarrow x = -1 \\
 &\quad -1
 \end{aligned}$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ is concave upward on $(-1, \infty)$

17) The function $f(x) = x^3 + 3x^2 - 9x + 1$ concave downward on

Solution:

$$\begin{aligned}
 f'(x) &= 3x^2 + 6x - 9 \\
 f''(x) &= 6x + 6 \\
 f''(x) = 0 &\Rightarrow 6x + 6 = 0 \\
 &\Rightarrow 6x = -6 \\
 &\Rightarrow x = -\frac{6}{6} \\
 &\Rightarrow x = -1 \\
 &\quad -1
 \end{aligned}$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ is concave downward on $(-\infty, -1)$

18) The function $f(x) = x^3 + 3x^2 - 9x + 1$ has an inflection point at

Solution:

$$f'(x) = 3x^2 + 6x - 9$$

$$f''(x) = 6x + 6$$

$$f''(x) = 0 \Rightarrow 6x + 6 = 0$$

$$\Rightarrow 6x = -6$$

$$\Rightarrow x = -\frac{6}{6}$$

$$\Rightarrow x = -1$$

$$-1$$

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ has an inflection point at $(-1, 12)$.

$$f(-1) = (-1)^3 + 3(-1)^2 - 9(-1) + 1$$

$$= -1 + 3 + 9 + 1 = 12$$

19) The critical numbers of the function $f(x) = x^3 - 3x^2 - 9x + 1$ are

Solution:

$$f'(x) = 3x^2 - 6x - 9$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x - 9 = 0$$

$$\Rightarrow 3(x^2 - 2x - 3) = 0$$

$$\Rightarrow x^2 - 2x - 3 = 0$$

$$\Rightarrow (x + 1)(x - 3) = 0$$

$$\Rightarrow x = -1 \text{ or } x = 3$$

20) The function $f(x) = x^3 - 3x^2 - 9x + 1$ is decreasing on

Solution:

$$f'(x) = 3x^2 - 6x - 9$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x - 9 = 0$$

$$\Rightarrow 3(x^2 - 2x - 3) = 0$$

$$\Rightarrow x^2 - 2x - 3 = 0$$

$$\Rightarrow (x + 1)(x - 3) = 0$$

$$\Rightarrow x = -1 \text{ or } x = 3$$

$$-1$$

$$3$$

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ is decreasing on $(-1, 3)$

21) The function $f(x) = x^3 - 3x^2 - 9x + 1$ is increasing on

Solution:

$$f'(x) = 3x^2 - 6x - 9$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x - 9 = 0$$

$$\Rightarrow 3(x^2 - 2x - 3) = 0$$

$$\Rightarrow x^2 - 2x - 3 = 0$$

$$\Rightarrow (x + 1)(x - 3) = 0$$

$$\Rightarrow x = -1 \text{ or } x = 3$$

$$-1$$

$$3$$

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ is increasing on $(-\infty, -1) \cup (3, \infty)$

22) The function $f(x) = x^3 - 3x^2 - 9x + 1$ has a relative maximum value at the point

Solution:

$$f'(x) = 3x^2 - 6x - 9$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x - 9 = 0$$

$$\Rightarrow 3(x^2 - 2x - 3) = 0$$

$$\Rightarrow x^2 - 2x - 3 = 0$$

$$\Rightarrow (x + 1)(x - 3) = 0$$

$$\Rightarrow x = -1 \text{ or } x = 3$$

$$-1$$

$$3$$

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ has a relative maximum value at the point $(-1, 6)$.

$$f(-1) = (-1)^3 - 3(-1)^2 - 9(-1) + 1$$

$$= -1 - 3 + 9 + 1 = 6.$$

23) The function $f(x) = x^3 - 3x^2 - 9x + 1$ has a relative minimum value at the point

Solution:

$$f'(x) = 3x^2 - 6x - 9$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x - 9 = 0$$

$$\Rightarrow 3(x^2 - 2x - 3) = 0$$

$$\Rightarrow x^2 - 2x - 3 = 0$$

$$\Rightarrow (x + 1)(x - 3) = 0$$

$$\Rightarrow x = -1 \text{ or } x = 3$$

$$-1$$

$$3$$

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ has a relative minimum value at the point $(3, -26)$.

$$f(3) = (3)^3 - 3(3)^2 - 9(3) + 1$$

$$= 27 - 27 - 27 + 1 = -26.$$

24) The function $f(x) = x^3 - 3x^2 - 9x + 1$ concave upward on

Solution:

$$f'(x) = 3x^2 - 6x - 9$$

$$f''(x) = 6x - 6$$

$$f''(x) = 0 \Rightarrow 6x - 6 = 0$$

$$\Rightarrow 6x = 6$$

$$\Rightarrow x = \frac{6}{6}$$

$$\Rightarrow x = 1$$

1

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ is concave upward on $(1, \infty)$

25) The function $f(x) = x^3 - 3x^2 - 9x + 1$ concave downward on

Solution:

$$f'(x) = 3x^2 - 6x - 9$$

$$f''(x) = 6x - 6$$

$$f''(x) = 0 \Rightarrow 6x - 6 = 0$$

$$\Rightarrow 6x = 6$$

$$\Rightarrow x = \frac{6}{6}$$

$$\Rightarrow x = 1$$

1

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ is concave downward on $(-\infty, 1)$

26) The function $f(x) = x^3 - 3x^2 - 9x + 1$ has an inflection point at

Solution:

$$f'(x) = 3x^2 - 6x - 9$$

$$f''(x) = 6x - 6$$

$$f''(x) = 0 \Rightarrow 6x - 6 = 0$$

$$\Rightarrow 6x = 6$$

$$\Rightarrow x = \frac{6}{6}$$

$$\Rightarrow x = 1$$

1

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ has an inflection point at $(1, -10)$.

$$f(1) = (1)^3 - 3(1)^2 - 9(1) + 1$$

$$= 1 - 3 - 9 + 1 = -10$$

27) The critical numbers of the function $f(x) = x^3 + 3x^2 - 9x + 5$ are

Solution:

$$f'(x) = 3x^2 + 6x - 9$$

$$f'(x) = 0 \Rightarrow 3x^2 + 6x - 9 = 0$$

$$\Rightarrow 3(x^2 + 2x - 3) = 0$$

$$\Rightarrow x^2 + 2x - 3 = 0$$

$$\Rightarrow (x + 3)(x - 1) = 0$$

$$\Rightarrow x = -3 \text{ or } x = 1$$

28) The function $f(x) = x^3 + 3x^2 - 9x + 5$ is decreasing on

Solution:

$$f'(x) = 3x^2 + 6x - 9$$

$$f'(x) = 0 \Rightarrow 3x^2 + 6x - 9 = 0$$

$$\Rightarrow 3(x^2 + 2x - 3) = 0$$

$$\Rightarrow x^2 + 2x - 3 = 0$$

$$\Rightarrow (x + 3)(x - 1) = 0$$

$$\Rightarrow x = -3 \text{ or } x = 1$$

-3

1

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ is decreasing on $(-3, 1)$.

29) The function $f(x) = x^3 + 3x^2 - 9x + 5$ is increasing on

Solution:

$$f'(x) = 3x^2 + 6x - 9$$

$$f'(x) = 0 \Rightarrow 3x^2 + 6x - 9 = 0$$

$$\Rightarrow 3(x^2 + 2x - 3) = 0$$

$$\Rightarrow x^2 + 2x - 3 = 0$$

$$\Rightarrow (x + 3)(x - 1) = 0$$

$$\Rightarrow x = -3 \text{ or } x = 1$$

-3

1

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ is increasing on $(-\infty, -3) \cup (1, \infty)$.

30) The function $f(x) = x^3 + 3x^2 - 9x + 5$ has a relative minimum value at the point

Solution:

$$f'(x) = 3x^2 + 6x - 9$$




$$f'(x) = 0 \Rightarrow 3x^2 + 6x - 9 = 0$$

$$\Rightarrow 3(x^2 + 2x - 3) = 0$$

$$\Rightarrow x^2 + 2x - 3 = 0$$

$$\Rightarrow (x + 3)(x - 1) = 0$$

$$\Rightarrow x = -3 \text{ or } x = 1$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity y

Hence, the function $f(x)$ has a relative minimum value at the point (1,0).

$$f(1) = (1)^3 + 3(1)^2 - 9(1) + 5$$

$$= 1 + 3 - 9 + 5 = 0$$

32) The function $f(x) = x^3 + 3x^2 - 9x + 5$ has an inflection point at

Solution:

$$f'(x) = 3x^2 + 6x - 9$$



$$f''(x) = 6x + 6$$

$$f''(x) = 0 \Rightarrow 6x + 6 = 0$$

$$\Rightarrow 6x = -6$$

$$\Rightarrow x = -\frac{6}{6}$$

$$\Rightarrow x = -1$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ has an inflection point at (-1,16).

$$f(-1) = (-1)^3 + 3(-1)^2 - 9(-1) + 5$$

$$= -1 + 3 + 9 + 5 = 16$$

34) The function $f(x) = x^3 + 3x^2 - 9x + 5$ concave upward on

Solution:

$$f'(x) = 3x^2 + 6x - 9$$



$$f''(x) = 6x + 6$$

$$f''(x) = 0 \Rightarrow 6x + 6 = 0$$

$$\Rightarrow 6x = -6$$

$$\Rightarrow x = -\frac{6}{6}$$

$$\Rightarrow x = -1$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ is concave upward on $(-1, \infty)$.

31) The function $f(x) = x^3 + 3x^2 - 9x + 5$ has a relative maximum value at the point

Solution:

$$f'(x) = 3x^2 + 6x - 9$$




$$f'(x) = 0 \Rightarrow 3x^2 + 6x - 9 = 0$$

$$\Rightarrow 3(x^2 + 2x - 3) = 0$$

$$\Rightarrow x^2 + 2x - 3 = 0$$

$$\Rightarrow (x + 3)(x - 1) = 0$$

$$\Rightarrow x = -3 \text{ or } x = 1$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity y

Hence, the function $f(x)$ has a relative maximum value at the point (-3,32).

$$f(-3) = (-3)^3 + 3(-3)^2 - 9(-3) + 5$$

$$= -27 + 27 + 27 + 5 = 32$$

33) The function $f(x) = x^3 + 3x^2 - 9x + 5$ concave downward on

Solution:

$$f'(x) = 3x^2 + 6x - 9$$



$$f''(x) = 6x + 6$$

$$f''(x) = 0 \Rightarrow 6x + 6 = 0$$

$$\Rightarrow 6x = -6$$

$$\Rightarrow x = -\frac{6}{6}$$

$$\Rightarrow x = -1$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ is concave downward on $(-\infty, -1)$.

35) The critical numbers of the function $f(x) = x^3 - 3x^2 - 9x + 5$ are

Solution:

$$f'(x) = 3x^2 - 6x - 9$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x - 9 = 0$$

$$\Rightarrow 3(x^2 - 2x - 3) = 0$$

$$\Rightarrow x^2 - 2x - 3 = 0$$

$$\Rightarrow (x + 1)(x - 3) = 0$$

$$\Rightarrow x = -1 \text{ or } x = 3$$

36) The function $f(x) = x^3 - 3x^2 - 9x + 5$ is increasing on

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 6x - 9 \\ f'(x) = 0 &\Rightarrow 3x^2 - 6x - 9 = 0 \\ &\Rightarrow 3(x^2 - 2x - 3) = 0 \\ &\Rightarrow x^2 - 2x - 3 = 0 \\ &\Rightarrow (x + 1)(x - 3) = 0 \\ &\Rightarrow x = -1 \text{ or } x = 3 \\ &\quad -1 \qquad \qquad 3 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ is increasing on $(-\infty, -1) \cup (3, \infty)$.

37) The function $f(x) = x^3 - 3x^2 - 9x + 5$ is decreasing on

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 6x - 9 \\ f'(x) = 0 &\Rightarrow 3x^2 - 6x - 9 = 0 \\ &\Rightarrow 3(x^2 - 2x - 3) = 0 \\ &\Rightarrow x^2 - 2x - 3 = 0 \\ &\Rightarrow (x + 1)(x - 3) = 0 \\ &\Rightarrow x = -1 \text{ or } x = 3 \\ &\quad -1 \qquad \qquad 3 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ is decreasing on $(-1, 3)$.

38) The function $f(x) = x^3 - 3x^2 - 9x + 5$ has a relative maximum value at the point

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 6x - 9 \\ f'(x) = 0 &\Rightarrow 3x^2 - 6x - 9 = 0 \\ &\Rightarrow 3(x^2 - 2x - 3) = 0 \\ &\Rightarrow x^2 - 2x - 3 = 0 \\ &\Rightarrow (x + 1)(x - 3) = 0 \\ &\Rightarrow x = -1 \text{ or } x = 3 \\ &\quad -1 \qquad \qquad 3 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative maximum value at the point $(-1, 10)$.

$$\begin{aligned} f(-1) &= (-1)^3 - 3(-1)^2 - 9(-1) + 5 \\ &= -1 - 3 + 9 + 5 = 10. \end{aligned}$$

39) The function $f(x) = x^3 - 3x^2 - 9x + 5$ has a relative minimum value at the point

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 6x - 9 \\ f'(x) = 0 &\Rightarrow 3x^2 - 6x - 9 = 0 \\ &\Rightarrow 3(x^2 - 2x - 3) = 0 \\ &\Rightarrow x^2 - 2x - 3 = 0 \\ &\Rightarrow (x + 1)(x - 3) = 0 \\ &\Rightarrow x = -1 \text{ or } x = 3 \\ &\quad -1 \qquad \qquad 3 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative minimum value at the point $(3, -22)$.

$$\begin{aligned} f(3) &= (3)^3 - 3(3)^2 - 9(3) + 5 \\ &= 27 - 27 - 27 + 5 = -22. \end{aligned}$$

40) The function $f(x) = x^3 - 3x^2 - 9x + 5$ concave upward on

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 6x - 9 \\ f''(x) &= 6x - 6 \\ f''(x) = 0 &\Rightarrow 6x - 6 = 0 \\ &\Rightarrow 6x = 6 \\ &\Rightarrow x = \frac{6}{6} \\ &\Rightarrow x = 1 \\ &\quad 1 \end{aligned}$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ is concave upward on $(1, \infty)$.

41) The function $f(x) = x^3 - 3x^2 - 9x + 5$ concave downward on

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 6x - 9 \\ f''(x) &= 6x - 6 \\ f''(x) = 0 &\Rightarrow 6x - 6 = 0 \\ &\Rightarrow 6x = 6 \\ &\Rightarrow x = \frac{6}{6} \\ &\Rightarrow x = 1 \\ &\quad 1 \end{aligned}$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ is concave downward on $(-\infty, 1)$.

42) The function $f(x) = x^3 - 3x^2 - 9x + 5$ has an inflection point at

Solution:

$$f'(x) = 3x^2 - 6x - 9$$

$$f''(x) = 6x - 6$$

$$f''(x) = 0 \Rightarrow 6x - 6 = 0$$

$$\Rightarrow 6x = 6$$

$$\Rightarrow x = \frac{6}{6}$$

$$\Rightarrow x = 1$$

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ has an inflection point at $(1, -6)$.

$$f(1) = (1)^3 - 3(1)^2 - 9(1) + 5$$

$$= 1 - 3 - 9 + 5 = -6$$

43) The critical numbers of the function

$$f(x) = \frac{1}{3}x^3 - \frac{1}{2}x^2 - 2x + 1$$
 are

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 - 2\left(\frac{1}{2}\right)x - 2 = x^2 - x - 2$$

$$f'(x) = 0 \Rightarrow x^2 - x - 2 = 0$$

$$\Rightarrow (x+1)(x-2) = 0$$

$$\Rightarrow x = -1 \text{ or } x = 2$$

44) The function $f(x) = \frac{1}{3}x^3 - \frac{1}{2}x^2 - 2x + 1$ is increasing on

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 - 2\left(\frac{1}{2}\right)x - 2 = x^2 - x - 2$$

$$f'(x) = 0 \Rightarrow x^2 - x - 2 = 0$$

$$\Rightarrow (x+1)(x-2) = 0$$

$$\Rightarrow x = -1 \text{ or } x = 2$$

$$\begin{matrix} -1 & & 2 \end{matrix}$$

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ is increasing on $(-\infty, -1) \cup (2, \infty)$.

45) The function $f(x) = \frac{1}{3}x^3 - \frac{1}{2}x^2 - 2x + 1$ is decreasing on

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 - 2\left(\frac{1}{2}\right)x - 2 = x^2 - x - 2$$

$$f'(x) = 0 \Rightarrow x^2 - x - 2 = 0$$

$$\Rightarrow (x+1)(x-2) = 0$$

$$\Rightarrow x = -1 \text{ or } x = 2$$

$$\begin{matrix} -1 & & 2 \end{matrix}$$

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ is decreasing on $(-1, 2)$.

46) The function $f(x) = \frac{1}{3}x^3 - \frac{1}{2}x^2 - 2x + 1$ has a relative maximum point

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 - 2\left(\frac{1}{2}\right)x - 2 = x^2 - x - 2$$

$$f'(x) = 0 \Rightarrow x^2 - x - 2 = 0$$

$$\Rightarrow (x+1)(x-2) = 0$$

$$\Rightarrow x = -1 \text{ or } x = 2$$

$$\begin{matrix} -1 & & 2 \end{matrix}$$

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ has a relative maximum point at $(-1, \frac{13}{6})$.

$$f(-1) = \frac{1}{3}(-1)^3 - \frac{1}{2}(-1)^2 - 2(-1) + 1$$

$$= -\frac{1}{3} - \frac{1}{2} + 2 + 1 = \frac{13}{6}$$

47) The function $f(x) = \frac{1}{3}x^3 - \frac{1}{2}x^2 - 2x + 1$ has a relative minimum point

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 - 2\left(\frac{1}{2}\right)x - 2 = x^2 - x - 2$$

$$f'(x) = 0 \Rightarrow x^2 - x - 2 = 0$$

$$\Rightarrow (x+1)(x-2) = 0$$

$$\Rightarrow x = -1 \text{ or } x = 2$$

$$\begin{matrix} -1 & & 2 \end{matrix}$$

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ has a relative minimum point at $(2, -\frac{7}{3})$.

$$f(2) = \frac{1}{3}(2)^3 - \frac{1}{2}(2)^2 - 2(2) + 1$$

$$= \frac{8}{3} - \frac{4}{2} - 4 + 1 = -\frac{7}{3}$$

48) The function $f(x) = \frac{1}{3}x^3 - \frac{1}{2}x^2 - 2x + 1$ concave upward on

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 - 2\left(\frac{1}{2}\right)x - 2 = x^2 - x - 2$$

$$f''(x) = 2x - 1$$

$$f''(x) = 0 \Rightarrow 2x - 1 = 0$$

$$\Rightarrow 2x = 1$$

$$\Rightarrow x = \frac{1}{2}$$

$\frac{1}{2}$

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ is concave upward on $\left(\frac{1}{2}, \infty\right)$.

49) The function $f(x) = \frac{1}{3}x^3 - \frac{1}{2}x^2 - 2x + 1$ concave downward on

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 - 2\left(\frac{1}{2}\right)x - 2 = x^2 - x - 2$$

$$f''(x) = 2x - 1$$

$$f''(x) = 0 \Rightarrow 2x - 1 = 0$$

$$\Rightarrow 2x = 1$$

$$\Rightarrow x = \frac{1}{2}$$

$\frac{1}{2}$

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ is concave downward on $\left(-\infty, \frac{1}{2}\right)$.

50) The function $f(x) = \frac{1}{3}x^3 - \frac{1}{2}x^2 - 2x + 1$ has an inflection point at

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 - 2\left(\frac{1}{2}\right)x - 2 = x^2 - x - 2$$

$$f''(x) = 2x - 1$$

$$f''(x) = 0 \Rightarrow 2x - 1 = 0$$

$$\Rightarrow 2x = 1$$

$$\Rightarrow x = \frac{1}{2}$$

$\frac{1}{2}$

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ has an inflection point at

$\left(\frac{1}{2}, -\frac{1}{12}\right)$.

$$f\left(\frac{1}{2}\right) = \frac{1}{3}\left(\frac{1}{2}\right)^3 - \frac{1}{2}\left(\frac{1}{2}\right)^2 - 2\left(\frac{1}{2}\right) + 1$$

$$= \frac{1}{24} - \frac{1}{8} - 1 + 1 = -\frac{1}{12}$$

51) The critical numbers of the function

$$f(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 - 2x + 1$$
 are

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 + 2\left(\frac{1}{2}\right)x - 2 = x^2 + x - 2$$

$$f'(x) = 0 \Rightarrow x^2 + x - 2 = 0$$

$$\Rightarrow (x+2)(x-1) = 0$$

$$\Rightarrow x = -2 \text{ or } x = 1$$

52) The function $f(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 - 2x + 1$ is increasing on

Solution:




$$f'(x) = 3\left(\frac{1}{3}\right)x^2 + 2\left(\frac{1}{2}\right)x - 2 = x^2 + x - 2$$

$$f'(x) = 0 \Rightarrow x^2 + x - 2 = 0$$

$$\Rightarrow (x+2)(x-1) = 0$$

$$\Rightarrow x = -2 \text{ or } x = 1$$

$$\begin{matrix} -2 & & 1 \end{matrix}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ is increasing on $(-\infty, -2) \cup (1, \infty)$.

53) The function $f(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 - 2x + 1$ is decreasing on

Solution:




$$f'(x) = 3\left(\frac{1}{3}\right)x^2 + 2\left(\frac{1}{2}\right)x - 2 = x^2 + x - 2$$

$$f'(x) = 0 \Rightarrow x^2 + x - 2 = 0$$

$$\Rightarrow (x+2)(x-1) = 0$$

$$\Rightarrow x = -2 \text{ or } x = 1$$

$$\begin{matrix} -2 & & 1 \end{matrix}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity




Hence, the function $f(x)$ is decreasing on $(-2, 1)$.

54) The function $f(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 - 2x + 1$ has a relative maximum point

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 + 2\left(\frac{1}{2}\right)x - 2 = x^2 + x - 2$$

$$\begin{aligned} f'(x) = 0 &\Rightarrow x^2 + x - 2 = 0 \\ &\Rightarrow (x+2)(x-1) = 0 \\ &\Rightarrow x = -2 \text{ or } x = 1 \\ &\quad -2 \qquad \qquad 1 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative maximum point at $\left(-2, \frac{13}{3}\right)$.




$$\begin{aligned} f(-2) &= \frac{1}{3}(-2)^3 + \frac{1}{2}(-2)^2 - 2(-2) + 1 \\ &= -\frac{8}{3} + \frac{4}{2} + 4 + 1 = \frac{13}{3} \end{aligned}$$

55) The function $f(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 - 2x + 1$ has a relative minimum point

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 + 2\left(\frac{1}{2}\right)x - 2 = x^2 + x - 2$$

$$\begin{aligned} f'(x) = 0 &\Rightarrow x^2 + x - 2 = 0 \\ &\Rightarrow (x+2)(x-1) = 0 \\ &\Rightarrow x = -2 \text{ or } x = 1 \\ &\quad -2 \qquad \qquad 1 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative minimum point at $\left(1, -\frac{1}{6}\right)$.

$$\begin{aligned} f(1) &= \frac{1}{3}(1)^3 + \frac{1}{2}(1)^2 - 2(1) + 1 \\ &= \frac{1}{3} + \frac{1}{2} - 2 + 1 = -\frac{1}{6} \end{aligned}$$



56) The function $f(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 - 2x + 1$ concave upward on

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 + 2\left(\frac{1}{2}\right)x - 2 = x^2 + x - 2$$

$$f''(x) = 2x + 1$$

$$\begin{aligned} f''(x) = 0 &\Rightarrow 2x + 1 = 0 \\ &\Rightarrow 2x = -1 \\ &\Rightarrow x = -\frac{1}{2} \\ &\quad -\frac{1}{2} \end{aligned}$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ is concave upward on $\left(-\frac{1}{2}, \infty\right)$.



57) The function $f(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 - 2x + 1$ concave downward on

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 + 2\left(\frac{1}{2}\right)x - 2 = x^2 + x - 2$$

$$f''(x) = 2x + 1$$

$$\begin{aligned} f''(x) = 0 &\Rightarrow 2x + 1 = 0 \\ &\Rightarrow 2x = -1 \\ &\Rightarrow x = -\frac{1}{2} \\ &\quad -\frac{1}{2} \end{aligned}$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ is concave downward on $\left(-\infty, -\frac{1}{2}\right)$.



58) The function $f(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 - 2x + 1$ has an inflection point at

Solution:

$$f'(x) = 3\left(\frac{1}{3}\right)x^2 + 2\left(\frac{1}{2}\right)x - 2 = x^2 + x - 2$$

$$f''(x) = 2x + 1$$

$$\begin{aligned} f''(x) = 0 &\Rightarrow 2x + 1 = 0 \\ &\Rightarrow 2x = -1 \Rightarrow x = -\frac{1}{2} \\ &\quad -\frac{1}{2} \end{aligned}$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ has an inflection point at $\left(-\frac{1}{2}, \frac{25}{12}\right)$.

$$\begin{aligned} f\left(-\frac{1}{2}\right) &= \frac{1}{3}\left(-\frac{1}{2}\right)^3 + \frac{1}{2}\left(-\frac{1}{2}\right)^2 - 2\left(-\frac{1}{2}\right) + 1 \\ &= -\frac{1}{24} + \frac{1}{8} + 1 + 1 = \frac{25}{12} \end{aligned}$$

59) The critical numbers of the function $f(x) = x^3 - 12x + 3$ are

Solution:

$$f'(x) = 3x^2 - 12$$

$$\begin{aligned} f'(x) = 0 &\Rightarrow 3x^2 - 12 = 0 \\ &\Rightarrow 3(x^2 - 4) = 0 \\ &\Rightarrow x^2 - 4 = 0 \\ &\Rightarrow x^2 = 4 \\ &\Rightarrow x = \pm 2 \end{aligned}$$

60) The function $f(x) = x^3 - 12x + 3$ is increasing on

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 12 \\ f'(x) = 0 &\Rightarrow 3x^2 - 12 = 0 \\ &\Rightarrow 3(x^2 - 4) = 0 \\ &\Rightarrow x^2 - 4 = 0 \\ &\Rightarrow x^2 = 4 \\ &\Rightarrow x = \pm 2 \\ &\quad -2 \qquad \qquad 2 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ is increasing on $(-\infty, -2) \cup (2, \infty)$.

61) The function $f(x) = x^3 - 12x + 3$ is decreasing on

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 12 \\ f'(x) = 0 &\Rightarrow 3x^2 - 12 = 0 \\ &\Rightarrow 3(x^2 - 4) = 0 \\ &\Rightarrow x^2 - 4 = 0 \\ &\Rightarrow x^2 = 4 \\ &\Rightarrow x = \pm 2 \\ &\quad -2 \qquad \qquad 2 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ is decreasing on $(-2, 2)$.

62) The function $f(x) = x^3 - 12x + 3$ has a relative maximum point at

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 12 \\ f'(x) = 0 &\Rightarrow 3x^2 - 12 = 0 \\ &\Rightarrow 3(x^2 - 4) = 0 \\ &\Rightarrow x^2 - 4 = 0 \\ &\Rightarrow x^2 = 4 \\ &\Rightarrow x = \pm 2 \\ &\quad -2 \qquad \qquad 2 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative maximum point at $(-2, 19)$.

$$\begin{aligned} f(-2) &= (-2)^3 - 12(-2) + 3 \\ &= -8 + 24 + 3 = 19. \end{aligned}$$

63) The function $f(x) = x^3 - 12x + 3$ has a relative minimum point at

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 12 \\ f'(x) = 0 &\Rightarrow 3x^2 - 12 = 0 \\ &\Rightarrow 3(x^2 - 4) = 0 \\ &\Rightarrow x^2 - 4 = 0 \\ &\Rightarrow x^2 = 4 \\ &\Rightarrow x = \pm 2 \\ &\quad -2 \qquad \qquad 2 \end{aligned}$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative minimum point at $(2, -13)$.

$$\begin{aligned} f(2) &= (2)^3 - 12(2) + 3 \\ &= 8 - 24 + 3 = -13 \end{aligned}$$

64) The function $f(x) = x^3 - 12x + 3$ concave upward on

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 12 \\ f''(x) &= 6x \\ f''(x) = 0 &\Rightarrow 6x = 0 \\ &\Rightarrow x = \frac{0}{6} \\ &\Rightarrow x = 0 \\ &\quad 0 \end{aligned}$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ is concave upward on $(0, \infty)$.

65) The function $f(x) = x^3 - 12x + 3$ concave downward on

Solution:

$$\begin{aligned} f'(x) &= 3x^2 - 12 \\ f''(x) &= 6x \\ f''(x) = 0 &\Rightarrow 6x = 0 \\ &\Rightarrow x = \frac{0}{6} \\ &\Rightarrow x = 0 \\ &\quad 0 \end{aligned}$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ is concave downward on $(-\infty, 0)$.

66) The function $f(x) = x^3 - 12x + 3$ has an inflection point at

Solution:

$$f'(x) = 3x^2 - 12$$

$$f''(x) = 6x$$

$$f''(x) = 0 \Rightarrow 6x = 0$$

$$\Rightarrow x = \frac{0}{6}$$

$$\Rightarrow x = 0$$

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ has an inflection point at $(0,3)$.

$$f(0) = (0)^3 - 12(0)^2 + 3$$

$$= 0 - 0 + 3 = 3$$

67) The critical numbers of the function $f(x) = x^3 - 3x^2 + 1$ are

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x = 0$$

$$\Rightarrow 3(x^2 - 2x) = 0$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x(x - 2) = 0$$

$$\Rightarrow x = 0 \text{ or } x = 2$$

68) The function $f(x) = x^3 - 3x^2 + 1$ is increasing on

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x = 0$$

$$\Rightarrow 3(x^2 - 2x) = 0$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x(x - 2) = 0$$

$$\Rightarrow x = 0 \text{ or } x = 2$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ is increasing on $(-\infty, 0) \cup (2, \infty)$.

69) The function $f(x) = x^3 - 3x^2 + 1$ is decreasing on

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x = 0$$

$$\Rightarrow 3(x^2 - 2x) = 0$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x(x - 2) = 0$$

$$\Rightarrow x = 0 \text{ or } x = 2$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ is decreasing on $(0,2)$.

70) The function $f(x) = x^3 - 3x^2 + 1$ has a relative maximum point at

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x = 0$$

$$\Rightarrow 3(x^2 - 2x) = 0$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x(x - 2) = 0$$

$$\Rightarrow x = 0 \text{ or } x = 2$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative maximum point at $(0,1)$.

$$f(0) = (0)^3 - 3(0)^2 + 1$$

$$= 0 - 0 + 1 = 1.$$

71) The function $f(x) = x^3 - 3x^2 + 1$ has a relative minimum point at

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x = 0$$

$$\Rightarrow 3(x^2 - 2x) = 0$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x(x - 2) = 0$$

$$\Rightarrow x = 0 \text{ or } x = 2$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative minimum point at $(2,-3)$.

$$f(2) = (2)^3 - 3(2)^2 + 1$$

$$= 8 - 12 + 1 = -3.$$

72) The function $f(x) = x^3 - 3x^2 + 1$ concave upward on

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f''(x) = 6x - 6$$

$$f''(x) = 0 \Rightarrow 6x - 6 = 0$$

$$\Rightarrow 6x = 6$$

$$\Rightarrow x = \frac{6}{6}$$

$$\Rightarrow x = 1$$

1

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ is concave upward on $(1, \infty)$.

73) The function $f(x) = x^3 - 3x^2 + 1$ concave downward on

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f''(x) = 6x - 6$$

$$f''(x) = 0 \Rightarrow 6x - 6 = 0$$

$$\Rightarrow 6x = 6$$

$$\Rightarrow x = \frac{6}{6}$$

$$\Rightarrow x = 1$$

1

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ is concave downward on $(-\infty, 1)$.

74) The function $f(x) = x^3 - 3x^2 + 1$ has an inflection point at

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f''(x) = 6x - 6$$

$$f''(x) = 0 \Rightarrow 6x - 6 = 0$$

$$\Rightarrow 6x = 6$$

$$\Rightarrow x = \frac{6}{6}$$

$$\Rightarrow x = 1$$

1

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ has an inflection point at $(1, -1)$.

$$f(1) = (1)^3 - 3(1)^2 + 1$$

$$= 1 - 3 + 1 = -1$$

75) The critical numbers of the function $f(x) = x^3 - 3x^2 + 2$ are

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x = 0$$

$$\Rightarrow 3(x^2 - 2x) = 0$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x(x - 2) = 0$$

$$\Rightarrow x = 0 \text{ or } x = 2$$

76) The function $f(x) = x^3 - 3x^2 + 2$ is increasing on

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x = 0$$

$$\Rightarrow 3(x^2 - 2x) = 0$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x(x - 2) = 0$$

$$\Rightarrow x = 0 \text{ or } x = 2$$

0

2

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ is increasing on $(-\infty, 0) \cup (2, \infty)$.

77) The function $f(x) = x^3 - 3x^2 + 2$ is decreasing on

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x = 0$$

$$\Rightarrow 3(x^2 - 2x) = 0$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x(x - 2) = 0$$

$$\Rightarrow x = 0 \text{ or } x = 2$$

0

2

+	-	+	Sign of $f'(x)$
\nearrow	\searrow	\nearrow	Kind of monotonicity

Hence, the function $f(x)$ is decreasing on $(0, 2)$.

78) The function $f(x) = x^3 - 3x^2 + 2$ has a relative minimum point at

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x = 0$$

$$\Rightarrow 3(x^2 - 2x) = 0$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x(x - 2) = 0$$

$$\Rightarrow x = 0 \text{ or } x = 2$$

+	-	+	Sign of $f'(x)$
↗	↘	↗	Kind of monotonicity

Hence, the function $f(x)$ has a relative minimum point at $(2, -2)$.

$$f(2) = (2)^3 - 3(2)^2 + 2$$

$$= 8 - 12 + 2 = -2.$$

79) The function $f(x) = x^3 - 3x^2 + 2$ has a relative maximum point at

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f'(x) = 0 \Rightarrow 3x^2 - 6x = 0$$

$$\Rightarrow 3(x^2 - 2x) = 0$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x(x - 2) = 0$$

$$\Rightarrow x = 0 \text{ or } x = 2$$

+	-	+	Sign of $f'(x)$
↗	↘	↗	Kind of monotonicity

Hence, the function $f(x)$ has a relative maximum point at $(0, 2)$.

$$f(0) = (0)^3 - 3(0)^2 + 2$$

$$= 0 - 0 + 2 = 2.$$

80) The function $f(x) = x^3 - 3x^2 + 2$ concave downward on

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f''(x) = 6x - 6$$

$$f''(x) = 0 \Rightarrow 6x - 6 = 0$$

$$\Rightarrow 6x = 6$$

$$\Rightarrow x = \frac{6}{6}$$

$$\Rightarrow x = 1$$

-	+	Sign of $f''(x)$
∩	∪	Kind of concavity

Hence, the function $f(x)$ is concave downward on $(-\infty, 1)$.

81) The function $f(x) = x^3 - 3x^2 + 2$ concave upward on

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f''(x) = 6x - 6$$

$$f''(x) = 0 \Rightarrow 6x - 6 = 0$$

$$\Rightarrow 6x = 6$$

$$\Rightarrow x = \frac{6}{6}$$

$$\Rightarrow x = 1$$

-	+	Sign of $f''(x)$
∩	∪	Kind of concavity

Hence, the function $f(x)$ is concave upward on $(1, \infty)$.

82) The function $f(x) = x^3 - 3x^2 + 2$ has an inflection point at

Solution:

$$f'(x) = 3x^2 - 6x$$

$$f''(x) = 6x - 6$$

$$f''(x) = 0 \Rightarrow 6x - 6 = 0$$

$$\Rightarrow 6x = 6$$

$$\Rightarrow x = \frac{6}{6}$$

$$\Rightarrow x = 1$$

-	+	Sign of $f''(x)$
∩	∪	Kind of concavity

Hence, the function $f(x)$ has an inflection point at $(1, 0)$.

$$f(1) = (1)^3 - 3(1)^2 + 2$$

$$= 1 - 3 + 2 = 0$$

83) The critical numbers of the function $f(x) = x^3 - 6x^2 - 36x$ are

Solution:

$$f'(x) = 3x^2 - 12x - 36$$

$$f'(x) = 0 \Rightarrow 3x^2 - 12x - 36 = 0$$

$$\Rightarrow 3(x^2 - 4x - 12) = 0$$

$$\Rightarrow x^2 - 4x - 12 = 0$$

$$\Rightarrow (x + 2)(x - 6) = 0$$

$$\Rightarrow x = -2 \text{ or } x = 6$$

84) The function $f(x) = x^3 - 6x^2 - 36x$ is decreasing on
Solution:

$$f'(x) = 3x^2 - 12x - 36$$

$$f'(x) = 0 \Rightarrow 3x^2 - 12x - 36 = 0$$

$$\Rightarrow 3(x^2 - 4x - 12) = 0$$

$$\Rightarrow x^2 - 4x - 12 = 0$$

$$\Rightarrow (x + 2)(x - 6) = 0$$

$$\Rightarrow x = -2 \text{ or } x = 6$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ is decreasing on $(-2, 6)$.

85) The function $f(x) = x^3 - 6x^2 - 36x$ is increasing on
Solution:

$$f'(x) = 3x^2 - 12x - 36$$

$$f'(x) = 0 \Rightarrow 3x^2 - 12x - 36 = 0$$

$$\Rightarrow 3(x^2 - 4x - 12) = 0$$

$$\Rightarrow x^2 - 4x - 12 = 0$$

$$\Rightarrow (x + 2)(x - 6) = 0$$

$$\Rightarrow x = -2 \text{ or } x = 6$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ is increasing on $(-\infty, -2) \cup (6, \infty)$.

86) The function $f(x) = x^3 - 6x^2 - 36x$ has a relative minimum value at the point
Solution:

$$f'(x) = 3x^2 - 12x - 36$$

$$f'(x) = 0 \Rightarrow 3x^2 - 12x - 36 = 0$$

$$\Rightarrow 3(x^2 - 4x - 12) = 0$$

$$\Rightarrow x^2 - 4x - 12 = 0$$

$$\Rightarrow (x + 2)(x - 6) = 0$$

$$\Rightarrow x = -2 \text{ or } x = 6$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative minimum value at the point $(6, -216)$.

$$f(6) = (6)^3 - 6(6)^2 - 36(6)$$

$$= 216 - 216 - 216 = -216$$

87) The function $f(x) = x^3 - 6x^2 - 36x$ has a relative maximum value at the point
Solution:

$$f'(x) = 3x^2 - 12x - 36$$

$$f'(x) = 0 \Rightarrow 3x^2 - 12x - 36 = 0$$

$$\Rightarrow 3(x^2 - 4x - 12) = 0$$

$$\Rightarrow x^2 - 4x - 12 = 0$$

$$\Rightarrow (x + 2)(x - 6) = 0$$

$$\Rightarrow x = -2 \text{ or } x = 6$$

+	-	+	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative maximum value at the point $(-2, 40)$.

$$f(-2) = (-2)^3 - 6(-2)^2 - 36(-2)$$

$$= -8 - 24 + 72 = 40$$

88) The function $f(x) = x^3 - 6x^2 - 36x$ has an inflection point at
Solution:

$$f'(x) = 3x^2 - 12x - 36$$

$$f''(x) = 6x - 12$$

$$f''(x) = 0 \Rightarrow 6x - 12 = 0$$

$$\Rightarrow 6x = 12$$

$$\Rightarrow x = \frac{12}{6}$$

$$\Rightarrow x = 2$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ has an inflection point at $(2, -88)$.

$$f(2) = (2)^3 - 6(2)^2 - 36(2)$$

$$= 8 - 24 - 72 = -88$$

89) The function $f(x) = x^3 - 6x^2 - 36x$ concave downward on
Solution:

$$f'(x) = 3x^2 - 12x - 36$$

$$f''(x) = 6x - 12$$

$$f''(x) = 0 \Rightarrow 6x - 12 = 0$$

$$\Rightarrow 6x = 12$$

$$\Rightarrow x = \frac{12}{6}$$

$$\Rightarrow x = 2$$

-	+	Sign of $f''(x)$
		Kind of concavity

Hence, the function $f(x)$ is concave downward on $(-\infty, 2)$.

90) The function $f(x) = x^3 - 6x^2 - 36x$ concave upward on

Solution:

$$f'(x) = 3x^2 - 12x - 36$$

$$f''(x) = 6x - 12$$

$$f''(x) = 0 \Rightarrow 6x - 12 = 0$$

$$\Rightarrow 6x = 12$$

$$\Rightarrow x = \frac{12}{6}$$

$$\Rightarrow x = 2$$

2

-	+	Sign of $f''(x)$
\cap	\cup	Kind of concavity

Hence, the function $f(x)$ is concave upward on $(2, \infty)$.

91) The critical numbers of the function $f(x) = -x^3 - 6x^2 - 9x + 1$ are

Solution:

$$f'(x) = -3x^2 - 12x - 9$$

$$f'(x) = 0 \Rightarrow -3x^2 - 12x - 9 = 0$$

$$\Rightarrow -3(x^2 + 4x + 3) = 0$$

$$\Rightarrow x^2 + 4x + 3 = 0$$

$$\Rightarrow (x + 3)(x + 1) = 0$$

$$\Rightarrow x = -3 \text{ or } x = -1$$

92) The function $f(x) = -x^3 - 6x^2 - 9x + 1$ is decreasing on

Solution:

$$f'(x) = -3x^2 - 12x - 9$$

$$f'(x) = 0 \Rightarrow -3x^2 - 12x - 9 = 0$$

$$\Rightarrow -3(x^2 + 4x + 3) = 0$$




$$\Rightarrow x^2 + 4x + 3 = 0$$

$$\Rightarrow (x + 3)(x + 1) = 0$$

$$\Rightarrow x = -3 \text{ or } x = -1$$

-3

-1

-	+	-	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ is decreasing on $(-\infty, -3) \cup (-1, \infty)$.

93) The function $f(x) = -x^3 - 6x^2 - 9x + 1$ is increasing on

Solution:

$$f'(x) = -3x^2 - 12x - 9$$

$$f'(x) = 0 \Rightarrow -3x^2 - 12x - 9 = 0$$

$$\Rightarrow -3(x^2 + 4x + 3) = 0$$




$$\Rightarrow x^2 + 4x + 3 = 0$$

$$\Rightarrow (x + 3)(x + 1) = 0$$

$$\Rightarrow x = -3 \text{ or } x = -1$$

-3

-1

-	+	-	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ is increasing on $(-3, -1)$.

94) The function $f(x) = -x^3 - 6x^2 - 9x + 1$ has a relative minimum value at the point

Solution:

$$f'(x) = -3x^2 - 12x - 9$$

$$f'(x) = 0 \Rightarrow -3x^2 - 12x - 9 = 0$$

$$\Rightarrow -3(x^2 + 4x + 3) = 0$$




$$\Rightarrow x^2 + 4x + 3 = 0$$

$$\Rightarrow (x + 3)(x + 1) = 0$$

$$\Rightarrow x = -3 \text{ or } x = -1$$

-3

-1

-	+	-	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative minimum value at the point $(-3, 1)$.

$$f(-3) = -(-3)^3 - 6(-3)^2 - 9(-3) + 1$$

$$= 27 - 54 + 27 + 1 = 1.$$

95) The function $f(x) = -x^3 - 6x^2 - 9x + 1$ has a relative maximum value at the point

Solution:

$$f'(x) = -3x^2 - 12x - 9$$

$$f'(x) = 0 \Rightarrow -3x^2 - 12x - 9 = 0$$

$$\Rightarrow -3(x^2 + 4x + 3) = 0$$




$$\Rightarrow x^2 + 4x + 3 = 0$$

$$\Rightarrow (x + 3)(x + 1) = 0$$

$$\Rightarrow x = -3 \text{ or } x = -1$$

-3

-1

-	+	-	Sign of $f'(x)$
			Kind of monotonicity

Hence, the function $f(x)$ has a relative maximum value at the point $(-1, 5)$.

$$f(-1) = -(-1)^3 - 6(-1)^2 - 9(-1) + 1$$

$$= 1 - 6 + 9 + 1 = 5.$$

96) The function $f(x) = -x^3 - 6x^2 - 9x + 1$ has an inflection point at

Solution:

$$f'(x) = -3x^2 - 12x - 9$$

$$f''(x) = -6x - 12$$

$$f''(x) = 0 \Rightarrow -6x - 12 = 0$$

$$\Rightarrow -6x = 12$$

$$\Rightarrow x = -\frac{12}{6}$$

$$\Rightarrow x = -2$$

$$-2$$

+	-	Sign of $f''(x)$
U	∩	Kind of concavity

Hence, the function $f(x)$ has an inflection point at $(-2, 3)$.

$$f(-2) = -(-2)^3 - 6(-2)^2 - 9(-2) + 1$$

$$= 8 - 24 + 18 + 1 = 3$$

97) The function $f(x) = -x^3 - 6x^2 - 9x + 1$ concave downward on

Solution:

$$f'(x) = -3x^2 - 12x - 9$$

$$f''(x) = -6x - 12$$

$$f''(x) = 0 \Rightarrow -6x - 12 = 0$$

$$\Rightarrow -6x = 12$$

$$\Rightarrow x = -\frac{12}{6}$$

$$\Rightarrow x = -2$$

$$-2$$

+	-	Sign of $f''(x)$
U	∩	Kind of concavity

Hence, the function $f(x)$ is concave downward on $(-2, \infty)$.

98) The function $f(x) = -x^3 - 6x^2 - 9x + 1$ concave upward on

Solution:

$$f'(x) = -3x^2 - 12x - 9$$

$$f''(x) = -6x - 12$$

$$f''(x) = 0 \Rightarrow -6x - 12 = 0$$

$$\Rightarrow -6x = 12$$

$$\Rightarrow x = -\frac{12}{6}$$

$$\Rightarrow x = -2$$

$$-2$$

+	-	Sign of $f''(x)$
U	∩	Kind of concavity

Hence, the function $f(x)$ is concave upward on $(-\infty, -2)$.

حفظ زمن النظريات والتعريفات التالية :

① Def. (1) \rightarrow absolute max. and absolute min.

② Def. (2) \rightarrow local max. and local min.

③ The Extreme Value Theorem.

f cont. on closed interval $[a, b]$

$\Rightarrow f$ has absolute max. & absolute min.
 \downarrow
 absolute extreme value

④ Def. (6) \rightarrow critical number

$f'(c) = 0$, $f'(c)$ does not exist

⑤ The closed Interval Method

To find an absolute ~~extrem~~ value of cont. f

on $[a, b]$:

(1) Find all critical numbers


(2) Evaluate $f(a)$, $f(b)$, and f at critical numbers

(3) The largest value \rightarrow abs. max.

The smallest value \rightarrow abs. min.

Find absolute max. and absolute min.


from ① graph



f has abs. max at a
 or min at b

ex

$f(a)$ is abs. max.
 $f(b)$ is abs. min.



f has abs. max at a_1, a_2
 or min at b

$f(a_1) = f(a_2)$ is abs. max.
 $f(b)$ is abs. min.

② fun.

مثلاً
 41
 (8)

عدد حرجات

يمكن ان الـ abs. max يكون عند هاهنا
 abs. min عند أكثر من مكان
 (يعني ممكن تأخذها عند أكثر من
 فتبينة عن قيم لا ولكن ههنا الـ abs. min)

4.3

حفظ نص النظريات والتعريفات التالية:

1] Increasing & Decreasing Test
(first derivative)

2] Local max. & local min.
(first derivative)

3] Concavity Test
(second derivative)

4] Def. (Inflection Point)

5] Local-max. & local min.
(second derivative)

First derivative (f')

- 1] increasing
- 2] decreasing
- 3] local max.
- 4] local min.

Second derivative (f'')

- 1] Concave up
- 2] Concave down
- 3] inflection point

5] The graph of the function f
 $f(x) = \frac{-1}{3}x^3 - 4x^2 - 1$ is concave up

on

- (a) $(-\infty, -4)$
- (b) $(-\infty, 4)$
- (c) $(4, \infty)$
- (d) $(-4, \infty)$


6] The inflection point of the function
 $f(x) = x^3 - 6x^2 - 36x$ is

- (a) $(2, f(2))$
- (b) $(-2, f(-2))$
- (c) $(0, f(0))$
- (d) No inflection point.

* كيف إيجاد

① intervals of increasing and decreasing

graph



$(1, 2) \cup (3, 5) \rightarrow$ increasing

$(2, 3) \rightarrow$ decreasing


first derivative
test

Ex (1)



② local max. & local min.

graph




3 \rightarrow local max

5 \rightarrow local min.


no local
max. &
local
min.

first derivative
test

Ex (2)




⑦ The absolute minimum of the function



- (a) $f(2)$
- (b) $f(1)$
- (c) $f(2), f(5)$
- (d) $f(6), f(9)$

⑧



The function f has local maximum at a

- (a) True
- (b) False

1 The function $f(x) = x^3 - 3x$ is decreasing on

- (a) $(-\infty, -1)$
 - (b) $(-1, \infty)$
 - (c) $(-\infty, -1) \cup (1, \infty)$
 - (d) $(-1, 1)$
-

2 If $f''(x) > 0$ for $1 < x < 3$, then the graph of $f(x)$ is concave down on $(1, 3)$

- (a) True
 - (b) False
-

3 The inflection point of the function

$$f(x) = x^3 - 12x + 2 \text{ is}$$

- (a) $(2, -4)$
 - (b) $(0, 12)$
 - (c) $(-2, 28)$
 - (d) f does not have an inflection point.
-

4 The function $f(x) = x^3 + 3x^2$ has

- (a) a local minimum at $x = -2$
- (b) a local maximum at $x = -2$
- (c) a local max. at $x = 0$
- (d) a local max. at $x = 2$

Find critical numbers of the function

$$f(x) = 5x^2 + 4x$$

Critical numbers ۳ خطوات لا يباد


① $D_f = \mathbb{R}$ (Polynomials)

② $f'(x) = 10x + 4$

③

$f'(x) = 0$

f' does not exist



$$\Rightarrow 10x + 4 = 0$$

$$\Rightarrow 10x = -4$$

$$\Rightarrow x = \frac{-4}{10} = -\frac{2}{5}$$

$$\left(x = -\frac{2}{5}\right) \in \mathbb{R} = D_f$$

\therefore Critical number is

$$\left(-\frac{2}{5}\right)$$

① The absolute extreme values of the function $f(x) = x^2 - 4$ on $[-1, 3]$ are

- (a) $f(1), f(0)$
- (b) $f(3), f(1)$
- (c) $f(3), f(-1)$
- (d) $f(3), f(0)$

② The critical numbers of the function $f(x) = x^3 - 3x^2 - 24x$ are

- (a) 2, 4
- (b) -2, -4
- (c) -2, 4
- (d) 2, -4

③ The absolute extreme of the function $f(x) = x^2 - 2x - 5$ on $[0, 3]$ are

	absolute min.	absolute max.
(a)	$f(3)$	$f(0)$
(b)	$f(0)$	$f(1)$
(c)	$f(0)$	$f(3)$
(d)	$f(1)$	$f(3)$

④ The absolute maximum value of $f(x) = 3x^2 - 12x + 1$ on $[1, 3]$ is (are)

- (a) $f(3)$
 - (b) $f(1), f(2)$
 - (c) $f(2)$
 - (d) $f(3), f(1)$
-

⑤ If f is continuous function on a closed interval $[a, b]$, then f attains an absolute maximum value $f(m)$ at some number m in $[a, b]$

- (a) True
 - (b) False.
-

⑥ Let h be a number in the domain D of a function f , then $f(h)$ is the absolute maximum value of f on D if $f(h) \leq f(x)$ for all $x \in D$

- (a) True
- (b) False

Find local max and local min

from ① graph

② Fun $\left(\frac{3}{4}, 2\right)$




ممكن يكون

عندك في أكثر من صنف لعل في
نصف المساحة، وهو ان يكون في
الوسط وليس في طرفي المساحة

local max

local min




لا يكون
الطرف

لا يكون في الطرف

3] Concave up & Concave down.

graph



$(1,3) \cup (4,5) \rightarrow$ Concave down

$(3,4) \cup (5,6) \rightarrow$ Concave up.


second derivative test

مثال (6)

f'' *ثانیه مشتق*

4] Inflection points

graph



3, 4, 5 \rightarrow inflection points

second derivative test

مثال (6)

f'' *ثانیه مشتق*