

# MATH203 Calculus

Dr. Bandar Al-Mohsin

School of Mathematics, KSU

8-12/9/13

# Outline

- Theorem 2.
- Theorem 3 ( $n$ th-term test).
- Theorem 4.
- Theorem 5.
- Theorem 6 (Properties of infinite series).
- Theorem 7.
- Examples.
- Positive term series.
- Theorem 1.
- Theorem 2 (Integral Test).
- Theorem 3 ( $p$ -Series Test).
- Theorem 4 (Basic Comparison Test).
- Theorem 5 (Limit Comparison Test).

### Theorem 2

If a series  $\sum_{n=1}^{\infty} a_n$  is c'gt, then  $\lim_{n \rightarrow \infty} a_n = 0$ .

### Theorem 3 (*n*th-term test)

If  $\lim_{n \rightarrow \infty} a_n \neq 0$ , then the series  $\sum_{n=1}^{\infty} a_n$  is d'gt.

### Theorem 4

If two series  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  are such that  $a_i = b_i$  for every  $i > k$ , where  $k$  is a positive interger, then both series converge or diverge together.

### Theorem 5

If we delete first  $k$  terms of a series

$\sum_{n=1}^{\infty} a_n = a_1 + a_2 + \cdots + a_k + \cdots + a_n + \dots$  then its behaviour does not change.

### Theorem 6 (properties)

Let  $\sum_{n=1}^{\infty} a_n = A$  and  $\sum_{n=1}^{\infty} b_n = B$  and  $C$  is a real number, then

- $\sum_{n=1}^{\infty} C a_n = C \sum_{n=1}^{\infty} a_n$
- $\sum_{n=1}^{\infty} (a_n \pm b_n) = A \pm B.$

### Theorem 7

If  $\sum_{n=1}^{\infty} a_n$  is convergent, and  $\sum_{n=1}^{\infty} b_n$  is divergent, then  $\sum_{n=1}^{\infty} (a_n + b_n)$  is divergent.

### Examples

**In page (26)** (i):  $3 + \frac{3}{4} + \cdots + \frac{3}{(4)^{n-1}} + \cdots$

(ii):  $\sum_{n=1}^{\infty} (\sqrt{2})^{n-1}$

**Solution:**

# Examples

**In page (27) Q25:** 
$$\sum_{n=1}^{\infty} a_n = \frac{1}{4 * 5} + \frac{1}{5 * 6} + \dots + \frac{1}{(n+3)(n+4)} + \dots$$

**Q28:** 
$$\sum_{n=1}^{\infty} a_n = \frac{-1}{1 * 2} + \frac{-1}{2 * 3} + \dots + \frac{-1}{n(n+1)} + \dots$$

**Solution:**

# Examples

**In page (28) Q1:** 
$$\sum_{n=1}^{\infty} \frac{3n}{(5n-1)}$$

Q2: 
$$\sum_{n=1}^{\infty} \frac{1}{\sqrt[n]{e}}$$

Q4: 
$$\sum_{n=1}^{\infty} \frac{n}{\ln(n+1)}$$

**Solution:**

## Def of Positive Term Series

a series  $\sum_{n=1}^{\infty} a_n$  such that  $a_n > 0$  for every  $n$

## Theorem 1

If  $\lim_{n \rightarrow \infty} a_n$  is a positive term series and if there exists a number  $M$  such that  $S_n = a_1 + a_2 + \cdots + a_n < M$  for every  $n$ , then the series is c'gt and has sum  $S \leq M$ . If no such  $M$  exists, then the series is d'gt.



## Theorem 2 (Integral Test)

Let  $\sum_{n=1}^{\infty} a_n$  be a positive term series. Suppose also

- $f$  is a positive continuous function for  $x \geq 1$  such that
- $f(n) = a_n$ , for  $n = 1, 2, 3, \dots$
- $f$  is a decreasing function of interval  $[1, \infty)$

then,  $\sum_{n=1}^{\infty} a_n$  is c'gt if  $\int_1^{\infty} f(x)dx$  is c'gt

and  $\sum_{n=1}^{\infty} a_n$  is d'gt if  $\int_1^{\infty} f(x)dx$  is d'gt

### Theorem 3 ( $p$ -Series Test)

The  $p$ -series is given by  $\sum_{n=1}^{\infty} \frac{1}{n^p} = \frac{1}{1^p} + \frac{1}{2^p} + \dots$ , where  $p > 0$  by definition.

- If  $p > 1$ , then the series converges.
- If  $0 < p \leq 1$ , then the series diverges.

### Theorem 4 (Basic Comparison Test)

Let  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  be two positive term series. If  $0 \leq a_n \leq b_n$  for all  $n$ , then the following rules apply:

- If  $\sum_{n=1}^{\infty} b_n$  converges, then  $\sum_{n=1}^{\infty} a_n$  converges.
- If  $\sum_{n=1}^{\infty} a_n$  diverges, then  $\sum_{n=1}^{\infty} b_n$  diverges.

### Theorem 5 (Limit Comparison Test)

Let  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  be two positive term series. If  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$ , where  $c > 0$ , then both series converge or diverge together.

# Examples

**In page (34) Q2:**  $\sum_{n=1}^{\infty} \frac{n^2}{e^{n^3}}$

Q3:  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$

Q4:  $\sum_{n=1}^{\infty} \frac{\arctan n}{1 + n^2}$

**Solution:**

# Examples

**In page (38)** (i):  $\sum_{n=1}^{\infty} \frac{1}{5 + 6^n}$  (hint: using direct CT)

(ii):  $\sum_{n=1}^{\infty} \frac{1}{\sqrt[4]{n} + 1}$  (hint: using direct CT)

**In page (39)** (i):  $\sum_{n=1}^{\infty} \frac{1}{1 + e^{2n}}$  (hint: using Limit CT)

(ii):  $\sum_{n=1}^{\infty} \frac{n^2 + \sqrt{n}}{6 + n^2 + n^{7/2}}$  (hint: using Limit CT)

**Solution:**

## The Ratio Test

Let  $\sum_{n=1}^{\infty} a_n$  be a positive term series and suppose that  $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = L$ , then

- the series  $\sum_{n=1}^{\infty} a_n$  converges if  $L < 1$ .
- the series  $\sum_{n=1}^{\infty} a_n$  diverges if  $L > 1$ .
- If  $L = 1$  (fails), the series may converge or diverge.

$$(i): \sum_{n=1}^{\infty} n!$$

$$(ii): \sum_{n=1}^{\infty} \frac{1}{(n+1)!}$$

## The Root Test

Let  $\sum_{n=1}^{\infty} a_n$  be a positive term series and suppose that  $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = L$ , then

- the series  $\sum_{n=1}^{\infty} a_n$  converges if  $L < 1$ .
- the series  $\sum_{n=1}^{\infty} a_n$  diverges if  $L > 1$ .
- If  $L = 1$  (fails), the series may converge or diverge.

$$(i): \sum_{n=1}^{\infty} \frac{5^n}{n^n}$$

$$(ii): \sum_{n=1}^{\infty} \left( \frac{8n^2 - 7}{n + 1} \right)^n$$