

المملكة العربية السعودية

وزارة التعليم

MINISTRY OF EDUCATION



لكل المهتمين و المهتمات
بدروس و مراجع الجامعية

هام

مدونة المناهج السعودية eduschool40.blog

Test bank chapter (1)

Choose the correct answer

1. The SI unit of time is the

- a) hour
b) **second**
c) minute
d) ampere

وحدة الوقت "time"
هي "second" الثانية

2. The diameter of an atom is approximately 1×10^{-7} mm. What is this diameter when expressed in nanometers?

- a) 1×10^{-18} nm
b) 1×10^{-15} nm
c) 1×10^{-9} nm
d) **1×10^{-1} nm**

من كبير (متر) إلى صغير (متر)
from mm to nm
 $1 \times 10^{-7} \times 10^6 = 1 \times 10^{-1} \text{ mm}$

3. 6.0 km is how many micrometers?

- a) 6.0×10^6 μm
b) 1.7×10^7 μm
c) **6.0×10^9 μm**
d) 1.7×10^4 μm

$6 \text{ km} \rightarrow \mu \text{ micrometer}$
كبير \times صغير
 $6 \times 10^3 = 6 \times 10^9 \mu\text{m}$

4. The SI prefixes giga and micro represent, respectively:

- a) 10^9 and 10^{-6} .
b) 10^6 and 10^{-3} .
c) 10^3 and 10^{-3} .
d) **10^9 and 10^{-6} .**

giga, micro
(G), (μ)
 $10^9, 10^{-6}$

جواب 5: مع صغير كبير \div
a) $2 \times 10^2 \text{ mg} \xrightarrow{\div 10^3} \text{g}$

$$2 \times 10^2 \div 10^3 = 0.2 \text{ g} = 2 \times 10^{-1} \text{ g}$$

b) $0.0010 \text{ kg} \xrightarrow{\times 10^3} \text{g}$ مع كبير صغير

$$0.0010 \times 10^3 = 1 \text{ g}$$

c) $1 \times 10^5 \text{ g} \xrightarrow{\div 10^6} \text{g}$ مع صغير كبير

$$1 \times 10^5 \div 10^6 = 1 \times 10^{-1} \text{ g} = 0.1 \text{ g}$$

d) $2 \times 10^2 \text{ cg} \xrightarrow{\div 10^2} \text{g}$ مع كبير كبير

$$2 \times 10^2 \div 10^2 = 2 \text{ g}$$

5. Which of these quantities represents the largest mass?

- a) 2.0×10^2 mg
b) 0.0010 kg
c) 1.0×10^5 μg
d) **2.0×10^2 cg**

مع الكبير كتلة

6. How many cubic centimeters are there in exactly one cubic meter?

- a) 1×10^6 cm^3
b) 1×10^3 cm^3
c) 1×10^2 cm^3
d) **1×10^6 cm^3**

from 1 m^3 to cm^3
كبير \rightarrow صغير
 $(1 \text{ m})^3 = (1 \times 10^2 \text{ cm})^3$
 $\text{m}^3 = 1 \times 10^6 \text{ cm}^3$

7. Ammonia boils at -33.4°C . What temperature is this in $^{\circ}\text{F}$?

- a) -60.1°F
- b) -92.1°F
- c) -28.1°F
- d) $+13.5^{\circ}\text{F}$

2 $^{\circ}\text{F} = \left(\frac{9}{5}\right) \cdot ^{\circ}\text{C} + 32$ القانون

$$^{\circ}\text{F} = \left(\frac{9}{5}\right)(-33.4) + 32$$

$$= -28.12 \approx -28.1^{\circ}\text{F}$$

8. Which of the following is not an SI base unit?

- a) **Kilometer** \times \rightarrow Prefix مشتقة
- b) Kilogram \checkmark كتلة
- c) Second \checkmark للوقت
- d) Kelvin \checkmark للحرارة

9. Which of the following SI base units is not commonly used in chemistry?

- a) kilogram
- b) kelvin
- c) **candela**
- d) mole

10. Which of the following prefixes means $1/1000$?

- a) kilo
- b) deci
- c) centi
- d) milli

$$1/1000 = 1 \times 10^{-3}$$

ألف درازي ال ميلي

11. Which of the following prefixes means 1000 ?

- a) kilo
- b) deci
- c) centi
- d) milli

$$1000 = 10^3$$

ألف درازي ال كيلو

12. Convert -77°F to kalvin ?

- a) 212.6 K
- b) -212.6 K
- c) -28.1 K
- d) $+13.5\text{ K}$

جاء ما في علاقة بين $^{\circ}\text{F}$ و $^{\circ}\text{C}$ وال K --
 مآحول من $^{\circ}\text{F}$ ل $^{\circ}\text{C}$ ومن ثم من $^{\circ}\text{C}$ ل K

$$^{\circ}\text{C} = \left(\frac{5}{9}\right)(^{\circ}\text{F} - 32)$$

$$= \left(\frac{5}{9}\right)(-77 - 32) = -60.55^{\circ}\text{C}$$

$$\text{K} = ^{\circ}\text{C} + 273.15$$

$$= -60.55 + 273.15$$

$$= 212.6\text{ K}$$

13. The number 0.0005678 expressed in scientific notation is:

- a) 5.678×10^4
- b) 5.67×10^{-7}
- c) 5.678×10^{-4}
- d) 5.678×10^{-3}

0.0005678
1 2 3 4
 5.678×10^{-4}

Explanation: Since this number is less than one start moving the decimal point to the right until there is ONE non-zero number to the left of the decimal point. Write the rest of the number as is. Write the exponent as the number of places the decimal point was moved.

14. Which of the following is the smallest distance? من أصغر المسافات
- a) 21 m
 - b) 2.1×10^2 m $\div 10^2 = 2.1$ m
 - c) 21 mm $\div 10^3 = 0.021$ m
 - d) 2.1×10^4 pm $\div 10^{12} = 2.1 \times 10^{-8}$ m

Explanation: Even though 2.1×10^4 is the largest number in this question, the units of pm (picometers) are the smallest units here, making it the smallest distance.

15. What temperature is 95°F when converted to degrees Celsius? القانون: $^\circ\text{C} = \left(\frac{5}{9}\right)(^\circ\text{F} - 32)$
- a) 63°C
 - b) 35°C
 - c) 127°C
 - d) 15°C
- $= \left(\frac{5}{9}\right)(95 - 32)$
 $= 35^\circ\text{C}$

16. What temperature is 37°C when converted to kelvin? القانون: $\text{K} = ^\circ\text{C} + 273.15$
- a) **310.15**
 - b) 99 k
 - c) 236 k
 - d) 67.15
- $\text{K} = 37 + 273.15$
 $= 310.15 \text{ K}$

17. What temperature is 77 K when converted to degrees Celsius? القانون: $\text{K} = ^\circ\text{C} + 273.15$
- a) -296°C
 - b) 105°C
 - c) -196°C
 - d) 25°C
- $77 = ^\circ\text{C} + 273.15$
 $^\circ\text{C} = 77 - 273.15$
 $= -196.15 \approx -196^\circ\text{C}$

18. Express 75 Tg as pg
- a) 0.75 pg
 - b) **75×10^{24} pg**
 - c) 0.75 pg
 - d) 75×10^{-24} pg



$75 \text{ Tg} \rightarrow \text{Pg}$
 $75 \times 10^{24} \text{ pg}$

19. The SI prefixes *Tera* and *nano* represent, respectively:

- a) 10^9 and 10^{-6}
- b) 10^6 and 10^{-3}
- c) 10^3 and 10^{-3}
- d) 10^{12} and 10^{-9}

Tera 10^{12}
Nano 10^{-9}

20. Which of these quantities represents the smallest mass?

- a) 2.0×10^2 mg
- b) 0.0010 kg
- c) 1×10^5 μ g
- d) 2.0×10^2 cg

أصغر كتلة (ج)

a) $2 \times 10^2 \text{ mg} \xrightarrow{\div 10^3} 0.2 = 2 \times 10^{-1} \text{ g}$ من صغرا كبر

b) $0.0010 \text{ kg} \xrightarrow{\times 10^3} 1 \text{ g}$ من كبر اصغر

c) $1 \times 10^5 \mu\text{g} \xrightarrow{\div 10^6} 0.1 \text{ g} \approx 1 \times 10^{-1} \text{ g}$ من صغرا كبر

d) $2 \times 10^2 \text{ cg} \xrightarrow{\div 10^2} 2 \text{ g}$ من صغرا كبر

21. Express 7.5 ng as Tg

- a) 7.5×10^{-21} Tg
- b) 75×10^{24} Tg
- c) 0.75 Tg
- d) 7.5×10^{21} Tg

$7.5 \text{ ng} \xrightarrow{\text{صغير كبر}} \text{Tg}$

$7.5 \div 10^{21} = 7.5 \times 10^{-21} \text{ Tg}$

الفرق بين 10^{-9} و 10^{12} هو $9+12=21$

28. At what temperature does the numerical reading on a Fahrenheit thermometer equal that on a Celsius thermometer?

- a) 0°F
- b) -40°F
- c) 100°F
- d) -32°F

اول طريقة: باي اعداد يكون درجة (مختلطة)
ثاني طريقة: املهازي انفسه

Explanation: since the temperature reading is the same so that mean $^\circ\text{F} = ^\circ\text{C}$

ثاني طريقة:

$^\circ\text{F} = [^\circ\text{C} \times 9/5] + 32^\circ\text{F}$

اول طريقة:

$^\circ\text{F} = ((\frac{9}{5})^\circ\text{C}) + 32$

ان اعداد درجات الحرارة هما نفس الارقام

Let temperature = t

$^\circ\text{C} = (\frac{5}{9})(40 - 32) = -40^\circ$

$t = (t(\frac{9}{5})) + 32$

$t = [t \times 9/5] + 32^\circ\text{F}$

انقله باليمين

$t - 9/5 t = 32^\circ\text{F}$

$t - (\frac{9}{5})t = 32$

$-4/5 t = 32^\circ\text{F}$

نضرب الطرفين في $\frac{5}{4}$ لكي نحصل على t وحده

$(-\frac{4}{4}) - \frac{4}{5}t = 32(-\frac{5}{4})$

$t = -40^\circ\text{F} = -40^\circ\text{C}$

$t = -40$

$t = -40^\circ\text{F}$

Test bank chapter (2)

Choose the correct answer

NOTE: A periodic table is required to work many of the problems in this chapter.

1. Which of these elements is most likely to be a good conductor of electricity?

- a) N *nonmetal*
 b) S *nonmetal*
 c) He *nonmetal*
 d) Fe *metal* ✓

2. An atom of the isotope sulfur-32 consists of how many protons, neutrons, and electrons?
 (p = proton, n = neutron, e = electron)

- a) 15 p, 16 n, 15 e
 b) 16 p, 15 n, 16 e
 c) 16 p, 31 n, 16 e
 d) 32 p, 31 n, 32 e

$$A \rightarrow 32$$


$$Z \rightarrow 16$$

$$P = 16, e = 16$$

$$n = 32 - 16 = 16$$

3. A magnesium ion, Mg^{2+} , has

- a) 12 protons and 13 electrons.
 b) 24 protons and 26 electrons.
 c) 12 protons and 10 electrons.
 d) 24 protons and 22 electrons.



$$p = 12$$

$$e = 12 - 2 = 10e$$

4. Which of these pairs of elements would be most likely to form an ionic compound?

- a) P and Br *non, non X*
 b) Cu and K *metal, metal X*
 c) C and O *non, non X*
 d) O and Zn *metal, nonmetal* ✓


nonmetal & metal

5. The elements in a column of the periodic table are known as

- a) metalloids. *ص.ع*
 b) a period.
 c) noble gases.
 d) a group.


6. Which is the correct formula for copper (II) phosphate?

- a) Cu_2PO_4
 b) $\text{Cu}_3(\text{PO}_4)_2$
 c) Cu_2PO_3
 d) $\text{Cu}(\text{PO}_4)_2$




7. The correct name for NH_4NO_3 is

- a) ammonium nitrate.
 b) ammonium nitrogen trioxide.
 c) ammonia nitrogen oxide.
 d) hydrogen nitrogen oxide.




8. What is the formula for the ionic compound formed by calcium ions and nitrate ions?

- a) Ca_3N_2
 b) $\text{Ca}(\text{NO}_3)_2$
 c) Ca_2NO_3
 d) Ca_2NO_2



9. The Stock system name for Mn_2O_7 is

- a) dimanganese heptaoxide.
 b) magnesium oxide.
 c) manganese(VII) oxide.
 d) manganese(II) oxide.



مanganese(VII) oxide
 (is not) Mn

10. Which of these elements is chemically similar to oxygen?


- a) sulfur
 b) calcium
 c) iron
 d) nickel

سلفور (is not) Mn

11. The formula of stannic oxide is SnO_2 . The valence of Sn is:

- a) +1
 b) +2
 c) +3
 d) +4

$$\begin{array}{l} X + (-2)(2) = 0 \\ X - 4 = 0 \\ X = +4 \end{array}$$



Explanation: to know the charge on Sn atom, make this simple calculation remember that the charge on oxygen atom is -2, let X is the charge on Sn atom

$$X + (-2 (\text{charge on O}) \times 2 (\text{number of O atoms})) = 0 \text{ (equal zero because the compound is neutral)}$$

$$X - 4 = 0 \gg \gg \gg \gg \gg x = +4$$

12. Which pair of atoms constitutes a pair of isotopes of the same element?

- (a) ${}_{6}^{14}\text{X}$ ${}_{7}^{14}\text{X}$ نفس العنصر
- (b) ${}_{6}^{14}\text{X}$ ${}_{6}^{12}\text{X}$ $\overset{A}{Z}\text{X}$
- (c) ${}_{9}^{17}\text{X}$ ${}_{8}^{17}\text{X}$
- (d) ${}_{10}^{19}\text{X}$ ${}_{9}^{19}\text{X}$

Explanation: Isotopes of an element are atoms of the same element with same number of protons but different number of neutrons. Only choice (b) has 2 atoms of X with 6 protons and 8 and 6 neutrons respectively.

13. Elements in Group 8A are known as the _____.

- a) chalcogens
- b) alkali metals
- c) **noble gases**
- d) alkaline earth metals

8A: Nobel gas

14. _____ typically forms ions with a 2+ charge.

- a) Transition metals
- b) Halogens
- c) **Alkaline earth metals**
- d) Alkali metals

A2 \Rightarrow +2 شحنة
Alkaline earth metal) الفلزات القلوية الأرضية

Explanation: The alkaline earth metals are in group 2A of the periodic table and lose 2 electrons to form cations with 2 positive charges.

15. An anion is defined as

- a) **a charged atom or group of atoms with a net negative charge.**
- b) a stable atom.
- c) a group of stable atoms.
- d) an atom or group of atoms with a net positive charge.

16. A cation is defined as

- a) a charged atom or group of atoms with a net negative charge.
- b) a stable atom.
- c) a group of stable atoms.
- d) **an atom or group of atoms with a net positive charge.**

17. Atoms of the same element with different mass numbers (or number of neutrons) are called

- a) ions.
- b) neutrons.
- c) chemical families.
- d) **isotopes.**

18. How many neutrons are there in an atom of lead ${}_{82}^{208}\text{Pb}$ whose mass number A is 208?

- a) 82
- b) **126** $n = A - Z = 208 - 82 = 126$
- c) 208
- d) 290

19. Molecules consist of the same element with different numbers of atoms and chemical structure are called ...


- a) Ions
- b) Neutrons
- c) **Allotropes**
- d) Isotopes

20. An atom of the isotope ${}^{16}\text{S}$ -31 consists of how many protons, neutrons, and electrons?

- a) 15 p, 16 n, 15 e
- b) **16 p, 15 n, 16 e**
- c) 16 p, 31 n, 16 e
- d) 32 p, 31 n, 32 e


21. A magnesium ion, ${}_{20}\text{Ca}^{2+}$, has

- a) 20 protons and 22 electrons.
- b) 20 protons and 20 electrons.
- c) **20 protons and 18 electrons.**
- d) 22 protons and 20 electrons.



22. A sulfide ion, ${}_{16}\text{S}^{2-}$, has:

- a) 16 protons and 16 electrons
- b) 32 protons and 16 electrons
- c) 16 protons and 14 electrons
- d) **16 protons and 18 electrons**




23. Which of these pairs of elements would be most likely to form a molecular compound?

- a) Na and Br (X) فلز ولا فلز
- b) Ca and O (X) فلز وفلز
- c) **C and O** $\text{كلاهما فلز ولا فلز}$
- d) Zn and O (X) فلز وفلز

لا فلز ولا فلز
شبه فلز مع لا فلز

24. What is the formula for the ionic compound formed by calcium ions and nitrate ions?

- a) Ca_3N_2
- b) **$\text{Ca}(\text{NO}_3)_2$**
- c) Ca_2NO_3
- d) Ca_2NO_2



25. Which is the correct formula for copper(II) phosphate?

- a) Cu_2PO_4
- b) $\text{Cu}_3(\text{PO}_4)_2$
- c) Cu_2PO_3
- d) $\text{Cu}(\text{PO}_4)_2$

26. The correct name for NH_4NO_3 is

- a) **ammonium nitrate.**
- b) ammonium nitrogen trioxide.
- c) ammonia nitrogen oxide.
- d) hydrogen nitrogen oxide.

27. The correct name for PCl_5 is

- a) monophosphate pentachloride
- b) phosphorus chloride
- c) monophosphate tetrachloride
- d) **phosphorus pentachloride**

PCl_5
Phosphorus pentachloride

28. Which of the following expressions represents two molecules of water?

- a) H_2O
- b) H_2O_2
- c) **$2\text{H}_2\text{O}$**
- d) 2HO_2

$2\text{H}_2\text{O}$

29. The empirical formula of a compound with molecules containing 12 carbon atoms, 14 hydrogen atoms, and 6 oxygen atoms is _____.

- a) $\text{C}_{12}\text{H}_{14}\text{O}_6$
- b) $\text{C}_2\text{H}_4\text{O}$
- c) CH_2O
- d) **$\text{C}_6\text{H}_7\text{O}_3$**

$\frac{\text{C}_{12}}{2} \frac{\text{H}_{14}}{2} \frac{\text{O}_6}{2}$ كبريتا - هيدروجين
 $\text{C}_6\text{H}_7\text{O}_3$

Explanation: The empirical formula is always the simplest possible whole number ratio between the atoms of the molecules.

30. The charge on the manganese in the salt MnF_3 is _____.


- a) +1
- b) -1
- c) **+3**
- d) -2

$$X + (3(-1)) = 0$$

$$X - 3 = 0$$

$$X = +3$$


(ج)



Explanation: Since every F has one negative charge, the Mn can have only 3 positive charges.

30. Magnesium reacts with a certain element to form a compound with the general formula MgX . What would the most likely formula be for the compound formed between potassium and element X?

- a) KX
 b) K_2X_2
 c) K_2X_3
 d) **None of the above**



Explanation: In the compound MgX , X must have 2 negative charges since Mg will always have 2 positive charges. The element K will always form an ion with 1 positive charge and hence the only combination of K and X could be K_2X , which is not one of the options.

31. Barium forms an ion with a charge of _____.

- a) +1
 b) -2
 c) +3
 d) **None of the above.**



Explanation: Barium is in group 2A of the periodic table and forms ions with only 2 positive charges.

31. Aluminum forms an ion with a charge of _____.

- a) +2
 b) -3
 c) **+3**
 d) +1




32. Iodine forms an ion with a charge of _____.

- a) -7
 b) +1
 c) **-1**
 d) +2



33. The chemical symbol for the ion with 11 protons and 10 electrons.

- a) Na
 b) F^-
 c) Ne
 d) **Na^+**



34. Which of these compounds is a binary compound? ثنائي

- a) **NaCl** → ثنائي ✓
- b) MgSO₄ } ❌
- c) NaOH } ❌
- d) HCN } ❌

35. Atoms with the same number of electrons and number of protons are called...

- a) ions
- b) isotopes
- c) **neutral atoms**
- d) different atoms

36. Atoms which have different number of electrons are called...

- a) **ions**
- b) isotopes
- c) neutral atoms
- d) different atoms

37. Use the following table and choose which of the species are positively charged? موجب

Atom or ion element	I	II	III	IV	V	VI
Atom or ion electrons (e)	6	10	18	10	28	7
Atom or ion protons (p)	6	8	17	11	30	7
Atom or ion neutrons (n)	6	8	18	11	36	6

- A. III and V
- B. **IV and V**
- C. II and III
- D. I and VI

38. Which isotope has 45 neutrons?

- (a). ⁸⁰₃₆Kr 80 - 36 = 44 ❌
- (b). ⁷⁸₃₄Se 78 - 34 = 44 ❌
- (c). ⁸⁰₃₅Br Ⓞ 80 - 35 = 45 ✓
- (d). ³⁴₁₇Cl 34 - 17 = 17 ❌

39. In the periodic table, the elements are arranged in _____.

- a) alphabetical order
- b) **order of increasing atomic number** مرتبة بزيادة العدد الذري
- c) order of increasing metallic properties
- d) order of increasing neutron content

الزاوية العلوية اليمنى

40. An element in the upper right corner of the periodic table is _____.

- a) either a metal or metalloid
- b) definitely a metal
- c) **definitely a non-metal**
- d) either a metalloid or a non-metal

الزاوية السفلية اليسرى

41. An element that appears in the lower left corner of a periodic table is _____.

- a) either a metal or metalloid
- b) **definitely a metal**
- c) either a metalloid or a non-metal
- d) definitely a non-metal

42. A molecular formula always indicates _____.

- a) **how many of each atom are in a molecule**
- b) the simplest whole-number ratio of different atoms in a compound
- c) which atoms are attached to which in a molecule
- d) the isotope of each element in a compound
- e)

43. An empirical formula always indicates _____.

- a) which atoms are attached to which in a molecule
- b) how many of each atom are in a molecule
- c) **the simplest whole-number ratio of different atoms in a compound**
- d) the geometry of a molecule

44. There are _____ protons, _____ neutrons, and _____ electrons in $^{131}\text{I}^-$.

- a) 131, 53, and 54
- b) 131, 53 and 52
- c) **53, 78, and 54**
- d) 53, 131, and 52

131 I⁻ اكتسب إلكترون

$$P = 53, e = 53 + 1 = 54, n = 78 \leftarrow 131 - 53$$

45. Which species has 48 electrons?

- (a) $^{118}_{50}\text{Sn}^{+2}$ ← 2 إلكترون ← 50 - 2 = 48 إلكترون ✓
- (b) $^{116}_{50}\text{Sn}^{+4}$ ← 4 إلكترون ← 50 - 4 = 46 إلكترون ✗
- (c) $^{112}_{48}\text{Cd}^{+2}$ ← 2 إلكترون ← 48 - 2 = 46 إلكترون ✗
- (d) $^{68}_{31}\text{Ga}$ ← 31 إلكترون ✗

Test bank chapter (3)

Choose the correct answer

1. What is the mass, in grams, of one copper atom?

- a) 1.055 10^{-22} g
- b) 63.55 g
- c) 1 amu
- d) 1.66 10^{-24} g

$$n = \frac{m}{M} \Rightarrow n = \frac{N}{N_A}$$

$$m = \frac{63.55}{6.022 \times 10^{23}} = 1.055 \times 10^{-22} \text{ g}$$

2. Determine the number of moles of aluminum in 96.7 g of Al.

- a) 0.279 mol
- b) 3.58 mol
- c) 7.43 mol
- d) 4.21 mol

$$n = \frac{m}{M} = \frac{96.7}{26.98} = 3.58 \text{ mol}$$

3. Which of the following samples contains the greatest number of atoms?

- a) 100 g of Pb
- b) 2.0 mole of Ar
- c) mole of Fe
- d) 5 g of He

$a) \frac{m}{M} = \frac{N}{N_A} \Rightarrow \frac{100}{207.2} = \frac{N}{6.022 \times 10^{23}} \Rightarrow N = 2.906 \times 10^{23} \text{ atoms}$
 $b) n = \frac{N}{N_A} \Rightarrow 2 = \frac{N}{6.022 \times 10^{23}} \Rightarrow N = 1.2044 \times 10^{24} \text{ atoms}$
 $c) n = \frac{N}{N_A} \Rightarrow 1 = \frac{N}{6.022 \times 10^{23}} \Rightarrow N = 6.022 \times 10^{23} \text{ atoms}$
 $d) \frac{m}{M} = \frac{N}{N_A} \Rightarrow \frac{5}{4.003} = \frac{N}{6.022 \times 10^{23}} \Rightarrow N = 7.522 \times 10^{23} \text{ atoms}$

4. Formaldehyde has the formula CH_2O . How many molecules are there in 0.11 g of formaldehyde?

- a) 6.1×10^{-27}
- b) 3.7×10^{-3}
- c) 4
- d) 2.2×10^{21}

$$\frac{m}{M} = \frac{N}{N_A} \Rightarrow \frac{0.11}{30.027} = \frac{N}{6.022 \times 10^{23}}$$

$$N = \frac{0.11 \times 6.022 \times 10^{23}}{30.027} = 2.2 \times 10^{21}$$

5. How many sulfur atoms are present in 25.6 g of $\text{Al}_2(\text{S}_2\text{O}_3)_3$?

- a) 0.393
- b) 6
- c) 3.95×10^{22}
- d) 2.37×10^{23}

$\frac{m}{M} = \frac{N}{N_A} \Rightarrow \frac{25.6}{390.399} = \frac{N}{6.022 \times 10^{23}}$
 $N = 3.949 \times 10^{22} \text{ molecules}$
 1 molecule $\rightarrow 6$ S atoms
 $3.949 \times 10^{22} \text{ molecules} \rightarrow 2.37 \times 10^{23} \text{ S atoms}$


6. The percent composition by mass of a compound is 76.0% C, 12.8% H, and 11.2% O. The molar mass of this compound is 284.5 g/mol. What is the molecular formula of the compound?

- a) $\text{C}_{10}\text{H}_6\text{O}$
- b) $\text{C}_9\text{H}_{18}\text{O}$
- c) $\text{C}_{16}\text{H}_{28}\text{O}_4$
- d) $\text{C}_{18}\text{H}_{36}\text{O}_2$


	C	H	O
	76g	12.8g	11.2g
$n = \frac{m}{M}$	$\frac{76}{12.01}$	$\frac{12.8}{1.008}$	$\frac{11.2}{16}$
	≈ 6.327	≈ 12.70	≈ 0.7
	$\frac{6.327}{0.7}$	$\frac{12.70}{0.7}$	$\frac{0.7}{0.7}$
	≈ 9	≈ 18.14	$= 1$
		≈ 18	2012

(1) 11% O
 (2) 12.8% H
 (3) 76% C
 (4) لا لازم يكون عدد صحيح


Ratios $\frac{284.5}{142.243} \approx 2$




7. What is the coefficient of H_2O when the following equation is properly balanced with the smallest set of whole numbers?



- a) 3
b) 4
c) 6
d) 12



8. When 22.0 g NaCl and 21.0 g H_2SO_4 are mixed and react according to the equation below, which is the limiting reagent? $2NaCl + H_2SO_4 \rightarrow Na_2SO_4 + 2HCl$



9. When the following equation is balanced, the coefficients are _____.




(a) 1, 1, 1, 1

(b) 2, 3, 2, 3


(c) 4, 7, 4, 6

(d) 1, 3, 1, 2



10. How many moles of carbon atoms are in 4 mol of dimethylsulfoxide (C_2H_6SO)?


- a) 2
b) 6
c) 8
d) 4



Explanation: This is based on reading the formula and correctly extracting information from it. The formula C_2H_6SO indicates that every mole of this compound has 2 moles of carbon atoms in it. Thus 4 moles of the compound would have $4 \times 2 = 8$ moles of C atoms.

11. There are _____ sulfur atoms in 25 molecules of $C_4H_4S_2$.

- a) 1.5×10^{25}
b) 4.8×10^{25}
c) 3.0×10^{23}
d) 50



Explanation: The molecular formula indicates that every molecule of $C_4H_4S_2$ has 2 sulfur atoms per molecule and hence 25 molecules of this compound will have $25 \times 2 = 50$ atoms of sulfur.

12. There are _____ hydrogen atoms in 25 molecules of $C_4H_4S_2$.

- a) 25
b) 3.8×10^{24}
c) 6.0×10^{25}
d) **100**

$$\begin{aligned} 1 \text{ molecule of } C_4H_4S_2 &\rightarrow 4 \text{ atoms of H} \\ 25 \text{ molecules of } C_4H_4S_2 &\rightarrow x \text{ atoms of H} \\ H &= 4 \times 25 = 100 \text{ atoms} \end{aligned}$$

Explanation: The formula of $C_4H_4S_2$ indicates that there are 4 hydrogen atoms per molecule and hence 100 hydrogen atoms in 25 molecules of $C_4H_4S_2$.

13. How many grams of oxygen are in 65.0 g of $C_2H_2O_2$?

- a) 18
b) 29
c) 9.5
d) **35.8**

$$\begin{aligned} \text{المطلوب} \{ 65 \text{ g of } C_2H_2O_2 &\Rightarrow x \text{ g of } O_2 \\ \text{النتيجة} \{ 58 \text{ g of } C_2H_2O_2 &\Rightarrow 32 \text{ g of } O_2 \\ x &= \frac{32 \times 65}{58} = 35.862 \text{ g} \end{aligned}$$

Explanation: This question uses the mole to mole ratio between oxygen and $C_2H_2O_2$ and needs the following

$$\text{steps. } \frac{65.0 \text{ g } C_2H_2O_2}{58.0 \text{ g} \cdot \text{mol}^{-1}} \times \frac{2 \text{ moles O}}{1 \text{ mole } C_2H_2O_2} \times \frac{15.99 \text{ g O}}{1 \text{ mole of O}} = 35.8 \text{ g of O}$$

17. How many moles of carbon dioxide are there in 52.06 g of carbon dioxide?

- a) 0.8452
b) **1.183**
c) 1.183×10^{23}
d) 8.648×10^2

$$\begin{aligned} n &= \frac{m}{MM} \\ n &= \frac{52.06}{44.01} = 1.183 \text{ mol} \end{aligned}$$

Explanation: This is a straight-forward conversion from grams to moles of CO_2 which is done as follows:

$$52.06 \text{ g } CO_2 \times \frac{1 \text{ mole } CO_2}{44.01 \text{ g } CO_2} = 1.183 \text{ moles of } CO_2$$

18. How many moles of the compound magnesium nitrate, $Mg(NO_3)_2$, are in a 2.35 g sample of this compound?

- a) 38.4
b) 65.8
c) **0.0158**
d) 0.0261

$$n = \frac{m}{MM} = \frac{2.35}{148.324} = 0.0158 \text{ moles}$$

Explanation: This is a straight-forward conversion from grams to moles of $Mg(NO_3)_2$ which is done as

follows:

$$2.35 \text{ g Mg(NO}_3)_2 \times \frac{1 \text{ mole Mg(NO}_3)_2}{148.3148 \text{ g}} = 0.0158 \text{ moles} \quad \text{C } \checkmark$$


19. A 25.5-g sample of ammonium carbonate contains _____ mol of ammonium ions.

- a) 0.468
 b) 0.288
 c) 0.531
 d) 2.00
- $\text{NH}_4^+ \text{CO}_3^{2-}$
 $(\text{NH}_4)_2\text{CO}_3$
 $n = \frac{25.5}{96.19} = 0.265 \text{ mol}$
- $1 \text{ mol of } (\text{NH}_4)_2\text{CO}_3 \rightarrow 2 \text{ mol of } \text{NH}_4$
 $0.265 \text{ mol of } (\text{NH}_4)_2\text{CO}_3 \rightarrow 0.531 \text{ mol of } \text{NH}_4$
 $\text{NH}_4 = 2 \times 0.2656 = 0.531$

Explanation: Realize that the formula for ammonium carbonate is $(\text{NH}_4)_2\text{CO}_3$ and calculate the molar mass (96.0856 g/mol). Convert grams to moles and then using the stoichiometric ratio find the # of moles of ammonium ions.

$$25.5 \text{ g } (\text{NH}_4)_2\text{CO}_3 \times \frac{1 \text{ mol } (\text{NH}_4)_2\text{CO}_3}{96.0856 \text{ g}} \times \frac{2 \text{ moles } \text{NH}_4^+}{1 \text{ mol } (\text{NH}_4)_2\text{CO}_3} = 0.531 \text{ moles} \quad \text{C } \checkmark$$

20. Magnesium and nitrogen react in a combination reaction to produce magnesium nitride:



In a particular experiment, a 5.47-g sample of N_2 reacts completely. How many grams of Mg are needed for

- a) 14.2 g
 b) 24.1 g
 c) 16.1 g
 d) 0.92 g
- $3\text{Mg} + \text{N}_2 \rightarrow \text{Mg}_3\text{N}_2$ this reaction?
 $x = \frac{5.47 \times 79.93}{28.014} = 14.24 \text{ g}$
 5.47 g
 79.93
 28.014 g

Explanation: Ensure that the equation is balanced. The grams of N_2 must be converted to moles of N_2 and then using the stoichiometric ratio between the Mg and N_2 , the grams of Mg can be calculated.

$$5.47 \text{ g } \text{N}_2 \times \frac{1 \text{ mole } \text{N}_2}{28.0134 \text{ g}} \times \frac{3 \text{ mole Mg}}{1 \text{ mole } \text{N}_2} \times \frac{24.3050 \text{ g Mg}}{1 \text{ mole Mg}} = 14.2 \text{ g Mg}$$

21. What information would you need to calculate the average atomic mass of an element?

- a) The number of neutrons in the element.
 b) The atomic number of the element.
 c) The mass and abundance of each isotope of the element.
 d) The position in the periodic table of the element.

22. The atomic masses of Cl (75.53 %) and Cl (24.47 %) are 34.968 amu and 36.956 amu, respectively. Calculate the average atomic mass of chlorine.

- a) 35.96 amu
 b) 35.45 amu
 c) 36.47 amu
 d) 71.92 amu

$$= \left(\frac{75.53}{100} \right) (34.968) + \left(\frac{24.47}{100} \right) (36.956)$$

$$= 35.45 \text{ amu}$$

23. How many atoms are there in 5.10 moles of sulfur ($16S=32$ amu)?

- a) 3.07×10^{24}
 b) 9.59×10^{22}
 c) 6.02×10^{23}
 d) 9.82×10^{25}

$$n = \frac{N}{N_A}$$

$$5.10 \times \frac{N}{6.022 \times 10^{23}} \Rightarrow N = 3.07 \times 10^{24}$$

24. Iodine has two isotopes ^{126}I and ^{127}I , with the equal abundance. Calculate the average atomic mass of Iodine (^{53}I).

- a) 126.5 amu
 b) 35.45 amu
 c) 1.265 amu
 d) 71.92 amu

$$= \frac{50}{100} (126) + \frac{50}{100} (127)$$

$$= 126.5 \text{ amu}$$

25. The atomic masses of ^6Li and ^7Li are 6.0151 amu and 7.0160 amu, respectively. Calculate the natural abundance of these two isotopes. The average atomic mass of Lithium ($\text{Li}=6.941$ amu).

- a) $^6\text{Li}=7.49\%$, $^7\text{Li}=92.51\%$
 b) $^7\text{Li}=7.49\%$, $^6\text{Li}=92.51\%$
 c) $^6\text{Li}=8.49\%$, $^7\text{Li}=95.51\%$
 d) $^7\text{Li}=7.22\%$, $^6\text{Li}=82.51\%$

$$\text{Average Atomic Mass} = \frac{\text{abundance}}{100} (\text{mass}) + \frac{\text{abundance}}{100} (\text{mass}) + \dots$$

$$6.941 = \frac{6.0151 \times X_1 + (100 - X_1)(7.0160)}{100}$$

$$694.1 = 6.0151X_1 + 701.6 - 7.0160X_1$$

$$\frac{-7.5}{-1.0009} = \frac{-1.0009X}{-1.0009} \Rightarrow X = 7.49\%$$

$$(100 - X) = (100 - 7.49)$$

$$= 92.51\%$$

26. How many atoms are present in 3.14 g of copper (Cu)?

- a) 2.98×10^{22}
 b) 1.92×10^{23}
 c) 1.89×10^{24}
 d) 6.02×10^{23}


$$n = \frac{m}{M} = \frac{N}{N_A}$$

$$\frac{3.14}{63.546} = \frac{N}{6.022 \times 10^{23}}$$

$$N = \frac{3.14 \times 6.022 \times 10^{23}}{63.546}$$

$$\approx 2.98 \times 10^{22} \text{ atoms}$$


27. Nitric oxide (NO) reacts with oxygen gas to form nitrogen dioxide (NO₂), a dark-brown gas:



In one experiment 0.886 mole of NO is mixed with 0.503 mole of O₂. Calculate the number of moles of NO₂ produced (note: first determine which is the limiting reagent).

a) 0.886 mol $2NO + O_2 \rightarrow 2NO_2$ { $2NO + O_2 \rightarrow 2NO_2$ }
 b) 0.503 mol $\frac{0.886}{2}$ } $\frac{0.503}{1}$ } $\frac{0.886}{2}$ } $X \text{ mol?}$ ← صف السؤال
 c) 1.01 mol $\frac{0.886}{2} = 0.443 \text{ mol} < 0.503 \text{ mol}$ } $\frac{0.886}{2}$ } $X \text{ mol?}$ ← صف المعطيات
 d) 1.77 mol $\frac{0.886}{2} = 0.443 \text{ mol}$ } $\frac{0.886}{2}$ } $X = \frac{0.886 \times 2}{2} = 0.886 \text{ moles}$


28. The fertilizer ammonium sulfate [(NH₄)₂SO₄] is prepared by the reaction between ammonia (NH₃) and sulfuric acid:



How many kilograms of NH₃ are needed to produce 1.00 × 10⁵ kg of (NH₄)₂SO₄?

a) 1.70 × 10⁴ kg $2NH_3 + H_2SO_4 \rightarrow (NH_4)_2SO_4$ } $X = \frac{34.06 \times 10^3}{132.08} = 25787401.57g \div 10^3$
 b) 3.22 × 10³ kg $\frac{1 \text{ kg?}}{X \text{ g?}}$ } $\frac{1 \times 10^5 \text{ kg}}{10^3 \text{ g}}$ } $= 25787.40$ (تحول 10³ بقسمة 10³)
 c) 2.58 × 10⁴ kg \downarrow } \downarrow } $= 2.58 \times 10^4 \text{ kg}$
 d) 7.42 × 10⁴ kg $34.06g$ } $132.08g$ }


29. Consider the combustion of carbon monoxide (CO) in oxygen gas:



Starting with 3.60 moles of CO, calculate the number of moles of CO₂ produced if there is enough oxygen gas to react with all of the CO.

a) 7.20 mol $2 \text{ mol of CO} \rightarrow 2 \text{ mol of CO}_2$
 b) 44.0 mol $3.60 \text{ mol of CO} \rightarrow X \text{ mol of CO}_2$
 c) 3.60 mol $\frac{2 \times X \text{ mole of CO}_2}{2} = \frac{2 \times 3.60}{2} \Rightarrow \text{mol of CO}_2 = 3.6 \text{ mol of CO}_2$
 d) 1.80 mol


30. Nitrous oxide (N₂O) is also called "laughing gas." It can be prepared by the thermal decomposition of ammonium Nitrate (NH₄NO₃). The other product is H₂O. The balanced equation for this reaction is:




How many grams of N₂O are formed if 0.46 mole of NH₄NO₃ is used in the reaction?

a) 2.0 g $NH_4NO_3 \rightarrow N_2O$ } $X = 0.46 \times 44.014$
 b) 3.7 × 10¹ g $\frac{0.46 \text{ mol}}{1 \text{ mol}}$ } $\frac{X \text{ g?}}{44.014}$ } $= 20.246$
 c) 2.0 × 10¹ g \downarrow } \downarrow } $= 2.0 \times 10^1 \text{ g}$
 d) 4.6 × 10⁻¹ g


31. What is the theoretical yield of chromium that can be produced by the reaction of 40.0 g of Cr₂O₃ with 8.00 g of aluminum according to the chemical equation below?




32. Hydrogen fluoride is used in the manufacture of Freons (which destroy ozone in the stratosphere) and in the production of aluminum metal. It is prepared by the reaction




33. When 22.0 g NaCl and 21.0 g H₂SO₄ are mixed and react according to the equation below, which is the limiting reagent?



34. Hydrochloric acid can be prepared by the following reaction:



How many grams of HCl can be prepared from 2.00 mol H₂SO₄ and 150 g NaCl?




Chem.110

Limiting reagent = 1.28 < 2 mol

2012

Dr. Laila Al-Harbi



35. Calculate the molar mass of Li_2CO_3 .

- a) 73.89 g
 b) 66.95 g
 c) 41.89 g
 d) 96.02 g

$$2(6.941) + 12.011 + 3(16)$$

$$M = 73.893 \text{ g/mol}$$

36. How many molecules of ethane (C_2H_6) are present in 0.334 g of C_2H_6 ?

- a) 2.01×10^{23}
 b) 6.69×10^{21}
 c) 4.96×10^{22}
 d) 8.89×10^{20}

$$n = \frac{m}{M} = \frac{N}{N_A}$$

$$\frac{0.334}{30.07} = \frac{N}{6.022 \times 10^{23}}$$

$$N = \frac{0.334 \times 6.022 \times 10^{23}}{30.07} = 6.689 \times 10^{21} \approx 6.69 \times 10^{21} \text{ molecules}$$

37. All of the substances listed below are fertilizers that contribute nitrogen to the soil. Which of these is the richest source of nitrogen on a mass percentage basis?

- a) Urea, $(\text{NH}_2)_2\text{CO}$
 b) Ammonium nitrate, NH_4NO_3
 c) Guanidine, $\text{HNC}(\text{NH}_2)_2$
 d) Ammonia, NH_3

$$\text{(a) } N\% = \frac{(2 \times 14)}{46} \times 100 = 70\%$$

$$\text{(b) } N\% = \frac{(2 \times 14)}{80} \times 100 = 35\%$$

$$\text{(c) } N\% = \frac{(3 \times 14)}{59} \times 100 = 71\%$$

$$\text{(d) } N\% = \frac{14}{17} \times 100 = 82\% \checkmark$$

38. Allicin is the compound responsible for the characteristic smell of garlic. An analysis of the compound gives the following percent composition by mass: C: 44.4 percent; H: 6.21 percent; S: 39.5 percent; O: 9.86 percent. What is its molecular formula given that its molar mass is about 162 g?


- a) $\text{C}_{12}\text{H}_{20}\text{S}_4\text{O}_2$
 b) $\text{C}_7\text{H}_{14}\text{SO}$
 c) $\text{C}_6\text{H}_{10}\text{S}_2\text{O}$
 d) $\text{C}_5\text{H}_{12}\text{S}_2\text{O}_2$

	C	H	S	O
a)	44.4	6.21	39.5	9.86
b)	12.011	1.008	32.066	16
c)	3.6966	6.1607	1.2318	0.61625
d)	5.9726	9.9721	1.9922	1


empirical formula $\text{C}_6\text{H}_{10}\text{S}_2\text{O}$
 Ratio = $\frac{162}{162.278} = 0.998 \approx 1$
 molecular formula

39. The formula for rust can be represented by Fe_2O_3 . How many moles of Fe are present in 24.6 g of the compound?

- a) 2.13 mol
 b) 0.456 mol
 c) 0.154 mol
 d) 0.308 mol



Chem.110



Dr. Laila Al-Harbi

$$x = 0.308 \text{ mol}$$

40. What is the mass, in grams, of one copper atom? $n = \frac{m}{M} = \frac{N}{N_A}$


- a) 1.055×10^{-22} g
 b) 63.55 g
 c) 1 amu
 d) 1.66×10^{-24} g

$$\frac{m}{63.546} \times \frac{1}{6.022 \times 10^{23}}$$


$$m = \frac{63.546}{6.022 \times 10^{23}} = 1.055 \times 10^{-22} \text{ g}$$

41. How many grams of sulfur (S) are needed to react completely with 246 g of mercury (Hg) to form HgS?

- a) 39.3 g
 b) 24.6 g
 c) 9.66×10^3 g
 d) 201 g




$$X = \frac{246 \times 32.066}{200.59} = 39.3\text{g}$$



42. Tin(II) fluoride (SnF_2) is often added to toothpaste as an ingredient to prevent tooth decay. What is the mass of F in grams in 24.6 g of the compound?

- a) 18.6 g
 b) 24.3 g
 c) 5.97 g
 d) 75.7 g




$$X = \frac{24.6 \times 37.996}{156.706} = 5.965 \approx 5.97\text{g}$$

43. What is the empirical formula of the compound with the following composition? 2.1 percent H, 65.3 percent O, 32.6 percent S.

- a) H_2SO_4
 b) H_2SO_3
 c) $\text{H}_2\text{S}_2\text{O}_3$
 d) HSO_3

H	O	S
$n = \frac{2.1}{1.008}$	$= \frac{65.3}{16}$	$= \frac{32.6}{32.066}$
$= 2.083$	$= 4.081$	$= 1.017$
≈ 2	≈ 4	$= 1$



(1) نصاب النسب
 (2) نطلع بحالات
 (3) نقسم على الصغرى

44. Which of the following samples contains the greatest number of atoms?

- a) 100 g of Pb
 b) 2.0 mole of Ar
 c) mole of Fe
 d) 5 g of He

45. Formaldehyde has the formula CH_2O . How many molecules are there in 0.11 g of formaldehyde?

- a) 6.1×10^{-27} molecule
 b) 3.7×10^{-3} molecule
 c) 4×10^{21} molecule
 d) 2.2×10^{21} molecule

$$\textcircled{1} n = \frac{m}{M} = \frac{0.11}{30.027} = 3.66 \times 10^{-3} \text{ mol}$$

$$\textcircled{2} n = \frac{N}{N_A} \quad 3.66 \times 10^{-3} = \frac{N}{6.022 \times 10^{23}}$$

$$N = 2.2 \times 10^{21} \text{ molecules}$$

46. Determine the number of moles of aluminum in 96.7 g of Al.

- a) 0.279 mol
- b) 3.58 mol
- c) 7.43 mol
- d) 4.21 mol

47. How many sulfur atoms are present in 25.6 g of $\text{Al}_2(\text{S}_2\text{O}_3)_3$?








- a) 0.393
- b) 3.95×10^{22}
- c) 7.90×10^{22}
- d) 2.37×10^{23}

48. What is the coefficient of H_2O when the following equation is properly balanced with the smallest set of whole numbers?



- a) 3
- b) 4
- c) 6
- d) 12

50. Which of the following equations is balanced?



51. Determine the number of moles of aluminum in 96.7 g of Al

- a) 0.279 mol
- b) 3.58 mol
- c) 7.43 mol
- d) 4.21 mol

Test bank chapter (4)

Choose the correct answer

1. A 50.0 mL sample of 0.436 M NH₄NO₃ is diluted with water to a total volume of 250.0 mL. What is the ammonium nitrate concentration in the resulting solution?

- a) 21.8 M
- b) 0.459 M
- c) 2.18 x 10⁻² M
- d) 8.72 x 10⁻² M

$M_i V_i = M_f V_f$ القانون

$$M_f = \frac{50 \times 0.436}{250}$$

$$M_f = 0.0872 = 8.72 \times 10^{-2} M$$

2. How many milliliters would you need to prepare 60.0 mL of 0.200 M HNO₃ from a stock solution of 4.00 M HNO₃?

- a) 3 mL
- b) 240 mL
- c) 24 mL
- d) 1000 mL

$M_i V_i = M_f V_f$

$$V_i = \frac{0.200 \times 60}{4}$$

$$V_i = 3 mL$$

3. What is the concentration (M) of KCl in a solution made by mixing 25.0 mL of 0.100 M KCl with 50.0 mL of 0.100 M KCl?

- a) 0.0500
- b) 0.100
- c) 0.0333
- d) 0.0250

$n_1 = M \times V = 0.100 \times 0.025 = 2.5 \times 10^{-3} \text{ mol}$

$n_2 = M \times V = 0.100 \times 0.05 = 5 \times 10^{-3} \text{ mol}$

$\text{Total of } n = 2.5 \times 10^{-3} + 5 \times 10^{-3} \text{ mol} = 7.5 \times 10^{-3} \text{ mol}$

$\text{Total of } V = V_1 + V_2 = 0.025 + 0.05 = 0.075 L$

$M = \frac{n}{V} = \frac{7.5 \times 10^{-3}}{0.075} = 0.1 M$

4. What is the concentration (M) of CH₃OH in a solution prepared by dissolving 11.7 g of CH₃OH in sufficient water to give exactly 230 mL of solution?

- a) 11.7
- b) 2.30 x 10⁻²
- c) 0.0841
- d) 1.59

$n = \frac{m}{MM}$

$$= \frac{11.7}{32.042} = 0.365 \text{ mol}$$

$M = \frac{n}{V}$

$$= \frac{0.365}{0.23} = 1.59$$

Explanation: Need to convert the grams of CH₃OH to moles and then find the molarity of the solution by using the molarity formula. Do not forget to convert the ml to L.

$$11.7 \text{ g CH}_3\text{OH} \times \frac{1 \text{ mole CH}_3\text{OH}}{32.042 \text{ g}} \times \frac{1}{0.230 \text{ L}} = 1.59 \text{ M}$$

$M_1 V_1 + M_2 V_2 = M_f V_f$

$$\frac{(0.1 \times 25) + (0.1 \times 50)}{75} = \frac{M_f (75)}{75}$$

$$M_f = \frac{7.5}{75} = 0.1 M$$

5. How many grams of H_3PO_4 are in 35.1 mL of a 2.75 M solution of H_3PO_4 ?

- a) 0.61
b) 9.46
c) 20
d) 4.9

$$M = \frac{n}{V}$$

$$2.75 = \frac{n}{0.0351} \Rightarrow n = 0.0965 \text{ mol}$$

$$n = \frac{m}{MM}$$

$$0.0965 = \frac{m}{97.998} \Rightarrow m = 9.46 \text{ g}$$

Explanation: Need to convert the ml of H_3PO_4 to liters and then find the # of moles of phosphoric acid. The moles of phosphoric acid can then be converted to grams of phosphoric acid.

$$35.1 \text{ mL} \times \frac{1 \text{ L}}{1000 \text{ mL}} \times 2.75 \text{ M} \times \frac{97.99 \text{ g } \text{H}_3\text{PO}_4}{1 \text{ mole}} = 9.46 \text{ g } \text{H}_3\text{PO}_4$$

7. What is the concentration (M) of a Na_2SO_4 solution prepared by dissolving 5.35 g of Na_2SO_4 in sufficient water to give 330 mL of solution?

- a) 1.14×10^2
b) 0.016
c) 61.7
d) 0.114

$$M = \frac{n}{V}$$

$$M = \frac{0.0377}{0.33}$$

$$= 0.114$$

$$n = \frac{m}{MM} = \frac{5.35}{142.042}$$

$$= 0.0377$$

Explanation: Convert grams of Na_2SO_4 to moles of Na_2SO_4 , ml of water to liters of water and then find the molarity of the solution by using the molarity formula.

$$5.35 \text{ g } \text{Na}_2\text{SO}_4 \times \frac{1 \text{ mole } \text{Na}_2\text{SO}_4}{142.035 \text{ g } \text{Na}_2\text{SO}_4} \times \frac{1}{0.330 \text{ L}} = 0.114 \text{ M } \text{Na}_2\text{SO}_4$$

8. How many grams of LiOH are there in 750.0 mL of a 0.0158 M LiOH solution?

- a) 2.11×10^{-5}
b) 11.3
c) 0.284
d) 3.50

$$M = \frac{n}{V}$$

$$n = 0.75 \times 0.0158$$

$$n = 0.01185 \text{ mol}$$

$$n = \frac{m}{MM}$$

$$m = 23.949 \times 0.01185$$

$$= 0.2838 \approx 0.284 \text{ g}$$

Explanation: Calculate the number of moles of LiOH present in this solution using the molarity formula and then convert the number of moles to grams of LiOH.

$$7.50 \times 10^{-1} \text{ L} \times 0.0158 \text{ M} \times \frac{23.948 \text{ g}}{1 \text{ mole LiOH}} = 0.284 \text{ g LiOH}$$

10. A 50.0 mL sample of 0.436 M NH_4NO_3 is diluted with water to a total volume of 250.0 mL. What is the ammonium nitrate concentration in the resulting solution?

- a) 21.8 M
b) 0.459 M
c) 2.18×10^{-2} M
d) 8.72×10^{-2} M

12. A 3.682 g sample of potassium chlorate KClO_3 is dissolved in enough water to give 375 mL of solution. What is the chlorate ion concentration in this solution?

- a) $3.00 \times 10^{-2} \text{ M}$
- b) $4.41 \times 10^{-2} \text{ M}$
- c) 0.118 M
- d) $8.01 \times 10^{-2} \text{ M}$

$$\frac{m}{MM} = M \times V$$

الحل باستخدام القانون:

$$\frac{3.682}{122.551} = M \times 0.375$$

$$\frac{0.030}{0.375} = \frac{0.375 M}{0.375}$$

$$M = 0.8801 \text{ M}$$

$$= 8.01 \times 10^{-2} \text{ M}$$

Test bank chapter (5)

Choose the most correct answer

1. A sample of oxygen occupies $\frac{V_1}{P_1}$ 47.2 liters under a pressure of $\frac{P_1}{P_2}$ 1240 torr at 25°C. What volume would it occupy at 25°C if the pressure were decreased to 730 torr?

- a) 27.8 L
- b) 29.3 L
- c) 32.3 L
- d) **80.2 L**

$$P_1 V_1 = P_2 V_2$$

$$\frac{1240 \times 47.2}{730} = \frac{730 V_2}{730}$$

$$V_2 = 80.2 \text{ L}$$

2. Under conditions of fixed temperature and amount of gas, Boyle's law requires that

- I. $P_1 V_1 = P_2 V_2$
- II. $PV = \text{constant}$
- III. $P_1/P_2 = V_2/V_1$

- a) I only
- b) II only
- c) III only
- d) **I, II, and III**

$$P_1 V_1 = \text{Constant}, P_2 V_2 = \text{Constant}$$

$$P_1 V_1 = P_2 V_2 \leftarrow \frac{P_1}{P_2} \times \frac{V_1}{V_2}$$

3. The volume of a sample of nitrogen is $\frac{V_1}{T_1}$ 6.00 liters at $\frac{P_1}{P_2}$ 35°C and 740 torr. What volume will it occupy at (STP)?

- a) 6.59 L
- b) 5.46 L
- c) 6.95 L
- d) **5.18 L**

$$35 + 273.15 = 308.15 \text{ K}$$

$$\frac{740}{760} = 0.97 \text{ atm}$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \Rightarrow \frac{0.97 \times 6}{308.15} = \frac{1 \times V_2}{273.15}$$

$$\Rightarrow \frac{308.15 V_2}{308.15} = \frac{1589.73}{308.15} = 5.16 \text{ L}$$

4. The density of chlorine gas at STP, in grams per liter, is approximately:

- a) 6.2
- b) **3.2**
- c) 3.9
- d) 4.5

$$d = \frac{PM}{RT} = \frac{1 \times 70.91}{0.0821 \times 273.15} = 3.2$$

Explanation: $d = \text{molar mass} \times p / RT = 70 \times 1 / 0.082 \times 273 = 3.17 \text{ g/L}$

5. What pressure (in atm) would be exerted by 76 g of fluorine gas in a 1.50 liter vessel at -37°C?

- a) **26 atm**
- b) 4.1 atm
- c) 19,600 atm
- d) 84 atm

$$PV = \frac{m}{M} RT$$

$$1.5P = \frac{76}{2 \times 16} \times 0.0821 \times 236.15 \Rightarrow 1.5P = 38.78 \Rightarrow P = 25.85 \approx 26 \text{ atm}$$

6. What is the density of ammonia gas at 2.00 atm pressure and a temperature of 25.0°C?

- a) 0.720 g/L
- b) 0.980 g/L
- c) **1.39 g/L**
- d) 16.6 g/L

$$d = \frac{PM}{RT} \Rightarrow d = \frac{2 \times 17.031}{0.0821 \times 298.15}$$

$$d = 1.39 \text{ g/L}$$

7. Convert 2.0 atm to mmHg

- a) 150 mmHg
- b) 0.27 mmHg
- c) 150 mmHg
- d) **1520 mmHg**

$$1 \text{ atm} \rightarrow 760 \text{ mmHg}$$

$$2 \text{ atm} \rightarrow X$$

$$X = 1520 \text{ mmHg}$$

8. A container with volume 71.9 mL contains water vapor at a pressure of 10.4 atm and a temperature of 465°C. How many grams of the gas are in the container?

- a) 0.421 g $PV = \frac{m}{MM} RT$
 b) **0.222 g**
 c) 0.183 g $10.4 \times 0.0719 = \frac{m}{18.016} \times 0.0821 \times 738.15$
 d) 0.129 g
- $0.74776 = \frac{60.53m}{18.016}$
 $0.74776 = 3.36m$
 $m = 0.222g$

Explanation: $n = PV/RT = 0.0719 \times 10.4 = 0.0821 \times (465 + 273) = 0.012$ mole

Mass = n × molar mass = 0.012 × 18 = 0.222 g

9. What is the molar mass of a pure gaseous compound having a density of 4.95 g/L at -35°C and 1020 torr?

- a) 24 g/mole $d = \frac{PM}{RT} \Rightarrow M = \frac{dRT}{P}$
 b) 11 g/mole
 c) **72 g/mole**
 d) 120 g/mole $MM = \frac{4.95 \times 0.0821 \times 238.15}{1.34} = 72.14 g/mole \approx 72$

10. A 0.580 g sample of a compound containing only carbon and hydrogen contains 0.480 g of carbon and 0.100 g of hydrogen. At STP, 33.6 mL of the gas has a mass of 0.087 g. What is the molecular (true) formula for the compound?

- a) CH₃
 b) C₂H₆
 c) C₂H₅
 d) **C₄H₁₀**
- $\frac{0.480}{12.011} = 0.040$
 $\frac{0.100}{1.008} = 0.099$
 $\frac{0.040}{0.040} = 1$
 $\frac{0.099}{0.040} = 2.5 \times 2$
 empirical formula: C₂H₅
 $MM = 29.062$
 $d = \frac{m}{V} = 2.59$
 $MM = \frac{dRT}{P} = 57.9956 \approx 58$
 Ratio = $\frac{58}{29.062} \approx 2$
 (C₂H₅)₂ = C₄H₁₀

11. Gas occupy 6L at 37°C what will be its volume when its temperature is doubled?

- a) **12 L**
 b) 6L
 c) 3.2 L
 d) 2L
- $\frac{V_1}{T_1} = \frac{V_2}{T_2}$
 $\frac{6}{310.15} = \frac{V_2}{620.3} \Rightarrow V_2 = \frac{6 \times 620.3}{310.15} = 12L$

12. A mixture of 90.0 grams of CH₄ and 10.0 grams of argon has a pressure of 250 torr under conditions of constant temperature and volume. The partial pressure of CH₄ in torr is:

- a) 143
 b) 100
 c) 10.7
 d) **239**
- $n_{CH_4} = \frac{90}{16.04} = 5.61 \text{ mol}$
 $n_{Ar} = \frac{10}{40} = 0.25 \text{ mol}$
 $X_{CH_4} = \frac{5.61}{5.61 + 0.25} = 0.96$
 $P_{CH_4} = 0.96 \times 250 = 239 \text{ torr}$

Explanation: from Dalton law $P_{CH_4} = X_{CH_4} P_{total}$, $n_{CH_4} = 90/16 = 5.625$ mole, $n_{Ar} = 10/39.95 = 0.250$ mole

$X_{CH_4} = n_{CH_4} / (n_{CH_4} + n_{Ar}) = 5.625 / (5.625 + 0.250) = 0.96$
 $P_{CH_4} = 0.96 \times 250 = 239.3$ torr

13. What pressure (in atm) would be exerted by a mixture of 1.4 g of nitrogen gas and 4.8 g of oxygen gas in a 200 mL container at 57°C?

- a) 4.7
 b) 34
 c) 47
 d) **27**
- $PV = n_{total} RT$
 $P = \frac{(0.05 + 0.15) \times 0.0821 \times 330.15}{0.2} \Rightarrow P = 27.07 \approx 27 \text{ atm}$

Explanation: $P = n_{total} RT/V$, $n_{N_2} = 1.4/28 = 0.05$ mole, $n_{O_2} = 4.8/32 = 0.15$ mole
 $P = (0.05 + 0.15) \times 0.0821 \times (57 + 273) / 0.2 = 27$ atm

$n_{N_2} = \frac{1.4}{28} = 0.05 \text{ mol}$
 Disalac
 calson JI

$n_{O_2} = \frac{4.8}{32} = 0.15 \text{ mol}$

14. A sample of hydrogen gas collected by displacement of water occupied 30.0 mL at 24°C and pressure 736 torr. What volume would the hydrogen occupy if it were dry and at STP? The vapor pressure of water at 24.0°C is 22.4 torr.

- a) 32.4 mL
 b) 21.6 mL
 c) 36.8 mL
 d) **25.9 mL**

$$P_T = P_{H_2} + P_{H_2O} \Rightarrow P_{H_2} = P_T - P_{H_2O}$$

$$P_{H_2} = 736 - 22.4 = 713.6 \text{ torr} \Rightarrow 0.939 \text{ atm}$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

Explanation: from Dalton law $\gg P_{H_2} = P_{\text{total}} - P_{H_2O}$, $P_{H_2} = 736 - 22.4 = 713.6 \text{ torr}$

$$n = PV/RT \gg n = (713.6/760) \times 0.03 / 0.0821 \times (24+273) = 0.00115 \text{ mole}$$


$$\text{at STP} \gg V = nRT/P = 0.00115 \times 0.0821 \times 273 / 1 = 0.026 \text{ L} \times 1000 = 25.89 \text{ mL}$$

$$\frac{1 \times V_1}{273.15} = \frac{0.030 \times 0.939}{297.15}$$

$$297.15 V_1 = 7.69$$

$$V_1 = 0.0259 \text{ L} \Rightarrow V = 25.9 \text{ mL}$$

15. Ammonia burns in oxygen gas to form nitric oxide (NO) and water vapor. How many volumes of NO are obtained from one volume of ammonia at the same temperature and pressure?



- a) **One**
 b) (b) Two
 c) (c) Three
 d) (d) Four

$$\frac{V_1}{n_1} = \frac{V_2}{n_2} \Rightarrow \frac{4 V_2}{4} = \frac{4}{4} \Rightarrow V_2 = 1$$

$$\frac{1}{4} = \frac{V_2}{4}$$

16. The pressure of 6.0 L of an ideal gas in a flexible container is decreased to one-third of its original value, and its absolute temperature is decreased by one-half. What is the final volume of the gas?

- a) **9.0 L**
 b) 6.0 L
 c) 4.0 L
 d) 1 L

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \Rightarrow \frac{6P}{T_1} = \frac{2P}{\frac{1}{2}T_2} \Rightarrow V_2 = \frac{6 \times 3}{2} = \frac{18}{2} = 9 \text{ L}$$

$$\frac{6P}{T_1} = \frac{1}{2} \frac{P_2 V_2}{T_2} \Rightarrow \frac{6RT_1}{T_1} = \frac{1}{2} \frac{P_2 V_2 T_1}{\frac{1}{2} T_2} \Rightarrow \frac{6RT_1}{T_1} = \frac{P_2 V_2 T_1}{T_2}$$

Explanation: let $V_1 = 6$ & $V_2 = ?$, $T_1 = T$ & $T_2 = \frac{1}{2}T$, $P_1 = P$ & $P_2 = \frac{1}{3}P$

$$\text{From combined gas law } P_1 V_1 / T_1 = P_2 V_2 / T_2 \gg \frac{P \times 6}{T} = \frac{\frac{1}{3}P \times V_2}{\frac{1}{2}T} \gg V_2 = \frac{P \times 6 \times T \times 3}{T \times 2 \times P} = 9 \text{ L}$$

17. Gas A is at 30°C and gas B is at 20°C. Both gases are at 1 atmosphere. What is the ratio of the volume of 1 mole gas A to 1 mole of gas B

- a) 606:303
 b) 3 : 2
 c) 2 : 3
 d) **303 : 293**

$$V_A = \frac{1}{303.15}, V_B = \frac{1}{293.15} \Rightarrow \frac{V_A}{303.15} = \frac{V_B}{293.15}$$

$$\text{Explanation: } \frac{V_A}{T_A} = \frac{V_B}{T_B} \gg \frac{V_A}{30+273} = \frac{V_B}{20+273} \gg \frac{V_A}{303} = \frac{V_B}{293}$$

18. The sample of argon occupies 50L at standard temperature. Assuming constant pressure, what volume with the gas occupy if the temperature is doubled.


- a) 25L
 b) 50L
 c) **100L**
 d) 100 mL

علاقة طردية بين الحجم والحرارة
 $T \uparrow, V \uparrow$
 بما انه T تتدبل يعني ال V تتدبل
 $50 \times 2 = 100 \text{ L}$

$$\frac{50}{273.15} = \frac{V_2}{273.15 \times 2} \Rightarrow V_2 = \frac{50 \times 546.3}{273.15} = 100 \text{ L}$$


او الحد المثلثون:

19. What total gas volume (in liters) at 520°C and 880 torr would result from the decomposition of 33 g of potassium bicarbonate according to the equation:



- (a) 56 L
 (b) 37 L
 (c) 10 L
 (d) 19 L
- $PV = nRT$
 $1.16 \times V = \frac{33}{100.117} \times 0.082 \times 793.15 \Rightarrow V = \frac{21.44}{1.16} = 18.5 \approx 19 \text{ L}$

20. Calculate the weight of KClO_3 that would be required to produce 29.5 L of oxygen measured at 127°C and 760 torr.



21. The ideal gas law predicts that the molar volume (volume of one mole) of gas equals:

- (a) mRT/PV
 (b) $(MM)P/RT$
 (c) $1/2ms^{-2}$
 (d) RT/P
- molar volume $(\frac{V}{n})$
 $PV = nRT \Rightarrow \frac{V}{n} = \frac{RT}{P}$

22. For a gas, which pair of variables are inversely proportional to each other (if all other conditions remain constant)?

- a) $P, V \rightarrow$ inverse
 b) $V, T \rightarrow$ direct
 c) $n, V \rightarrow$ direct
 d) n, P

23. Convert 562 mmHg to atm

- a) 0.739 atm
 b) 4.27×10^5 atm
 c) 1.05 atm
 d) 0.562 atm
- $x = \frac{562}{760} = 0.739 \text{ atm}$

24. What is the volume of one mole of an ideal gas at STP?

- a) 24.5 L
 b) 22.4 L
 c) 1.0 L
 d) 10.0 L

25. What are standard temperature and pressure (STP)?

- a) 0 C, 1 torr
 b) 25 C, 1 torr
 c) 0 C, 1 atm
 d) 25 °C, 1 atm
- $\begin{matrix} T & P \\ \downarrow & \downarrow \\ 0^\circ\text{C} & 1 \text{ atm} \end{matrix}$

26. What is the unit of mole fraction

- a) mol
 b) mol^{-1}
 c) unitless
- x_i

27. Refer to Dalton's law of partial pressures and explain what mole fraction is
- The number of moles of one component
 - The ratio of the number of moles of one component to the number of moles of all components present.**
 - The number of moles of one component divided by 100
 - The ratio of the number of moles of all components present to the number of moles of one component.
28. Write the ideal gas equation. Give the units for each term in the equation
- $PV = nRT$; P in torr, V in L, n in mol, R in Latm/Kmol, T in $^{\circ}\text{C}$.
 - $PV = nRT$; P in torr, V in L, n in mol, R in Latm/Kmol, T in K.
 - (c) $PV = nRT$; P in atm, V in L, n in mol, R in Latm/Kmol, T in K.**
 - $PV = nRT$; P in atm, V in L, n in mol, R in Latm/Kmol, T in $^{\circ}\text{C}$.
29. What is the difference between a gas and a vapor?
- A gas is a substance normally in the gaseous state at normal atmospheric conditions (25C, 1 atm); a vapor is the gaseous form of any substance that is a liquid or a solid at normal temperatures and pressures.**
 - A gas is the gaseous form of any substance; a vapor refers to a gas over a water surface.
 - A gas is a substance normally in the gaseous state at normal atmospheric conditions (25C, 1 atm); a vapor is a gas over a water surface.
 - A gas and a vapor are two interchangeable nomenclatures; they are identical.
30. What volume is occupied by 19.6 g of methane (CH_4) at 27°C and 1.59 atm?
- 1.71 L
 - 18.9 L**
 - 27.7 L
 - 302 L

31. A 4.37 gram sample of a certain diatomic gas occupies a volume of 3.00 L at 1.00 atm and a temperature of 45°C . Identify this gas.

$d = \frac{m}{V} = \frac{4.37}{3} = 1.46$ $PV = \frac{m}{M} RT$
 $d = \frac{PMM}{RT} \Rightarrow MM = \frac{dRT}{P} = 38.1$ $M = \frac{mRT}{PV}$
 $MM = \frac{4.37 \times 0.082 \times 318.15}{1 \times 3} \Rightarrow MM = 38 \text{ g/mol}$

28 g/mol ← a) F_2
 28 g/mol ← b) N_2
 2 g/mol ← c) H_2
 32 g/mol ← d) O_2

Explanation: $MM = mRT/PV \gggg MM = 4.37 \times 0.0821 \times (45+273) / 1 \times 3 = 37.77/2 = 18.88 \text{ g/mole- } \text{F}_2$

32. A sample of hydrogen gas was collected over water at 21°C and 685 mmHg. The volume of the container was 7.80 L. Calculate the mass of H_2 (g) collected. (Vapor pressure of water = 18.6 mmHg at 21°C .)

$P_T = P_{\text{H}_2} + P_{\text{H}_2\text{O}}$
 $P_{\text{H}_2} = 685 - 18.6$
 $P_{\text{H}_2} = 666.4 \text{ mmHg} \approx 0.88 \text{ atm}$

$PV = \frac{m}{MM} RT$
 $m = \frac{PVMM}{RT} = \frac{0.88 \times 7.80 \times 2.016}{0.082 \times 294.15} = 0.573 \text{ g}$

- 0.283 g
- 0.572 g**
- 0.589 g
- 7.14

33. Which of the following is/are characteristic(s) of gases?

- High compressibility
- Relatively large distances between molecules
- Formation of homogeneous mixtures regardless of the nature of gases
- High compressibility, relatively large distances between molecules AND formation of homogeneous mixtures regardless of the nature of gases**

34. A small bubble rises from the bottom of a lake, where the temperature and pressure are 4°C and 3.0 atm , to the water's surface, where the temperature is 25°C and the pressure is 0.95 atm . Calculate the final volume of the bubble if its initial volume was 2.1 mL .

- a) 0.72 mL
 b) 6.2 mL
 c) 41.4 mL
 d) 7.1 mL

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$\frac{3 \times 2.1}{277.15} = \frac{0.95 \times V_2}{298.15}$$

$$0.0227 = 3.17 \times 10^{-3} V_2$$

$$V_2 = 7.1\text{ mL}$$

35. Calculate the mass, in grams, of 2.74 L of CO gas measured at 33°C and 945 mmHg .

- a) 0.263 g
 b) 2.46 g
 c) 3.80 g
 d) 35.2 g

$$PV = nRT$$

$$PV = \frac{m}{MM} RT$$

$$m = \frac{PVMM}{RT}$$

$$m = \frac{1.24 \times 2.74 \times 28.01}{0.082 \times 306.15}$$

$$m = 3.79\text{ g} \approx 3.8\text{ g}$$

36. Which of the following gases will have the greatest density at the same specified temperature and pressure?

- a) H_2
 b) CCl_4
 c) CO_2
 d) C_2H_6

$$D = \frac{PM}{RT} \Rightarrow P, T \text{ constant} \Rightarrow d = \frac{M}{R}$$

a) $D_{\text{H}_2} = \frac{1.008 \times 2}{0.082} = 24.59\text{ g/L}$

b) $D_{\text{CCl}_4} = \frac{154.46}{0.082} = 1273.9\text{ g/L}$

c) $D_{\text{CO}_2} = \frac{44.01}{0.082} = 536.72$

d) $D_{\text{C}_2\text{H}_6} = \frac{30.07}{0.082} = 366.71$

37. Determine the molar mass of chloroform gas if a sample weighing 0.389 g is collected in a flask with a volume of 102 cm^3 at 97°C . The pressure of the chloroform is 728 mmHg .

$1\text{ L} = 1000\text{ mL}$
 $1\text{ mL} = 1\text{ cm}^3$
 $102 \times 10^{-3} = 0.102\text{ L}$

- a) 187 g/mol
 b) 121 g/mol
 c) 112 g/mol
 d) 31.6 g/mol

$$PV = nRT$$

$$PV = \frac{m}{MM} RT$$

$$MM = \frac{mRT}{PV}$$

$$MM = \frac{0.389 \times 0.082 \times 370.15}{0.102 \times 0.96} = 120.58 \approx 121\text{ g/mol}$$

38. What is the molar mass of Freon-11 gas if its density is 6.13 g/L at STP ?

- a) 0.274 g/mol
 b) 3.64 g/mol
 c) 78.2 g/mol
 d) 137 g/mol

$$D = \frac{PM}{RT}$$

$$MM = \frac{DRT}{P}$$

$$MM = \frac{6.13 \times 0.082 \times 273.15}{1} = 137.3 \approx 137\text{ g/mol}$$

40. A mixture of three gases has a total pressure of $1,380\text{ mmHg}$ at 298 K . The mixture is analyzed and is found to contain 1.27 mol CO_2 , 3.04 mol CO , and 1.50 mol Ar . What is the partial pressure of Ar ?

- a) 0.258 atm
 b) 301 mmHg
 c) 356 mmHg
 d) $5,345\text{ mmHg}$

$$P_{\text{Ar}} = (X_{\text{Ar}}) \cdot P_T$$

$$X_{\text{Ar}} = \frac{1.50}{1.50 + 1.27 + 3.04} = 0.258$$

$$P_{\text{Ar}} = 0.258 \times 1380 = 356.04 \approx 356\text{ mmHg}$$

41. A sample of hydrogen gas was collected over water at 21°C and 685 mmHg. The volume of the container was 7.80 L. Calculate the mass of H₂(g) collected. (Vapor pressure of water = 18.6 mmHg at 21°C.)

- a) 0.283 g
- b) 0.572 g
- c) 0.589 g
- d) 7.14 g

42. A 0.271 g sample of an unknown vapor occupies 294 mL at 140°C and 847 mmHg. The empirical formula of the compound is CH₂. What is the molecular formula of the compound?

MM = 14.027 g/mol


- a) CH₂
- b) C₂H₄
- c) C₃H₆
- d) C₄H₈

$PV = \frac{m}{MM} RT$
 $MM = \frac{mRT}{PV}$


$MM = \frac{0.271 \times 0.082 \times 413.15}{0.294 \times 0.294}$
 $MM = 28.13 \text{ g/mol}$

Ratio = $\frac{28.13}{14.027} \approx 2$
 $(CH_2)_2 \Rightarrow C_2H_4$

43. How many liters of chlorine gas at 25°C and 0.950 atm can be produced by the reaction of 12.0 g of MnO₂?



- a) 5.36×10^{-3} L
- b) 0.138 L
- c) 0.282 L
- d) 3.55 L



$X = 0.138 \text{ mol}$

$PV = nRT \Rightarrow V = \frac{nRT}{P}$
 $= \frac{0.138 \times 0.082 \times 298.15}{0.950} = 3.55 \text{ L}$

Choose the most correct answer

1. The lowest energy state of an atom is referred to as its

- a) bottom state.
- b) **ground state.**
- c) fundamental state.
- d) original state.

2. All s orbitals are

- a) shaped like four-leaf clovers.
- b) dumbbell-shaped.
- c) **spherical.**
- d) triangular.

3. $[\text{He}]2s^2 2p^2$ is the electron configuration of which element?

- a) Beryllium Be
- b) Boron B
- c) **carbon C**
- d) nitrogen N

$2s^2 2p^2 = 4$ المجموعة
الدورة الثانية

4. What are the valence electrons of vanadium (V)?

- a) $4s^2$
- b) $3d^3$
- c) **$4s^2 3d^3$**
- d) $3d^3$

$V^{23}: 1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^3$

5. What are the valence electrons of gallium Ga?

- a) $4s^2$
- b) $3d^3$
- c) **$4s^2 4p^1$**
- d) $3d^3$

$Ga^{31}: 1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^1$
 $ns^2 np^1$

6. The electron configuration of a neutral atom is $[\text{Ne}] 3s^1 3p^1$. The four quantum numbers of the last electron are:

- a) (2, 1, -1, +1/2)
- b) (3, 3, -1, +1/2)
- c) (3, 0, -1, +1/2)
- d) (3, 1, -1, +1/2)



$n=3, l=1, m_l=-1, m_s=+1/2$

$(3, 1, -1, +1/2)$

7. How many unpaired electrons does chromium (Cr) have?

- a) 0
- b) 2
- c) 4
- d) **6**


$Cr^{24}: [\text{Ar}] 4s^2 3d^4$ less stable
 $[\text{Ar}] 4s^1 3d^5$ more stable



8. How many unpaired electrons does selenium (Se) have?

- a) 0
- b) **2**
- c) 4
- d) 6

$Se: [\text{Ar}] 4s^2 4p^4$



9. What is the maximum number of orbitals described by the quantum numbers: $n=3$ $l=2$ عدد
شكل
d
- a) 1
b) 3
c) **5**
d) 9
- $ml = 2l + 1$
 $ml = 2(2) + 1 = 5$

10. What is the maximum number of orbitals described by the quantum numbers: $n=4$ اسطوان
n
- a) 7
b) 14
c) **16**
d) 48
- $n^2 = 4^2 = 16$

11. The maximum number of electrons that can occupy an energy level described by the principal quantum number, n , is
- a) $n + 1$
b) $2n$
c) **$2n^2$**
d) n^2
- $2n^2$

12. A possible set of quantum numbers for the last electron added to complete an atom of sodium (Na) in its ground state is
- a) $n=3, l=1, m_l=0, m_s=1/2$
b) $n=3, l=0, m_l=0, m_s=1/2$
c) $n=2, l=1, m_l=-1, m_s=1/2$
d) $n=2, l=0, m_l=-1, m_s=1/2$
- $Na: 1s^2 2s^2 2p^6 3s^1$ last e
 $(3, 0, 0, +1/2)$
- $n=3$
 $l=0$
 $m_l=0$
 $m_s=+1/2$

13. The ground-state electron configuration of a calcium atom is
- a) $[Ne]3s^2$
b) $[Ne]3s^2 3p^6$
c) $[Ar]4s^1 3d^1$
d) **$[Ar]4s^2$**
- $Ca^{20}: [Ar] 4s^2$

14. Which one of the following sets of quantum numbers is not possible?

	n	l	m_l	m_s
Row 1	4	3	-2	+1/2
Row 2	3	2	-3	-1/2
Row 3	3	0	0	+1/2
Row 4	4	1	1	-1/2
Row 5	2	0	0	+1/2

$d(2, 0, -3)$ → $-2 -1 0 +1 +2$

- a) Row 1
b) **Row 2**
c) Row 3
d) Row 4

15. The number of orbitals in a d subshell is


- a) 1
b) 3
c) **5**
d) 7

$l=2$ بالاقواس → $ml = 2l + 1$
 $= 2(2) + 1 = 5$

عدد الفوت 5

□ □ □ □ □

16. Which ground-state atom has an electron configuration described by the following orbital diagram?




- a) phosphorus
- b) germanium
- c) selenium
- d) tellurium

بالذرة الرابعة
والجموعه ال 6 سادسه
 $4s^2 4p^4 = 6$

17. A ground-state atom of nickel has ___ unpaired electrons and is ___.

- a) 0, diamagnetic
- b) 6, diamagnetic
- c) 3, paramagnetic
- d) 2, paramagnetic




18. What is the frequency (s^{-1}) of electromagnetic radiation that has a wavelength of 0.53 m?

- a) 5.7×10^8
 - b) 1.8×10^{-9}
 - c) 1.6×10^8
 - d) 1.3×10^{-33}
- $c = \lambda \nu$
 $\nu = \frac{c}{\lambda} \Rightarrow \nu = \frac{3 \times 10^8}{0.53} = 566037735.8 \text{ (د) } 5.7 \times 10^8$

Explanation: The frequency and wavelength of electromagnetic radiation are related by the equation $c = \lambda \nu$, where c is the speed of light ($=3.00 \times 10^8 \text{ m/s}$), λ is the wavelength in m and ν is the frequency is s^{-1} or Hz. The frequency can be calculated by rearranging the above formula to get $\nu = c/\lambda = 3 \times 10^8 / 0.53 = 5.7 \times 10^8 s^{-1}$

19. The energy of a photon of light is ___ proportional to its frequency and ___ proportional to its wavelength.

- a) directly, directly
- b) inversely, inversely
- c) inversely, directly
- d) directly, inversely



20. The wavelength of a photon of energy $5.25 \times 10^{-19} \text{ J}$ is ___ m.

- a) 2.64×10^6
- b) 3.79×10^{-7}
- c) 2.38×10^{23}
- d) 4.21×10^{-24}

$E = h \frac{c}{\lambda}$

$5.25 \times 10^{-19} = \frac{6.63 \times 10^{-34} \times 3 \times 10^8}{\lambda}$

$\lambda = \frac{6.63 \times 10^{-34} \times 3 \times 10^8}{5.25 \times 10^{-19}}$

$\lambda = 3.79 \times 10^{-7} \text{ m}$

Explanation: The wavelength and energy are related by the formula $E = hc/\lambda$, where h ($6.626 \times 10^{-34} \text{ Js}$) is Planck's constant, c is the speed of light ($3.00 \times 10^8 \text{ m/s}$) and λ is the wavelength in meters. The wavelength can then be calculated by rearranging the above formula as follows: $\lambda = hc/E = 6.63 \times 10^{-34} \times 3 \times 10^8 / 5.25 \times 10^{-19} = 3.79 \times 10^{-7} \text{ m}$

21. What is the frequency (s^{-1}) of a photon of energy $4.38 \times 10^{-18} J$?

- a) 438
- b) 1.45×10^{-16}
- c) **6.61×10^{15}**
- d) 2.30×10^7

$$E = h\nu$$

$$\nu = \frac{E}{h} \Rightarrow \nu = \frac{4.38 \times 10^{-18}}{6.63 \times 10^{-34}} = 6.61 \times 10^{15} \text{ Hz} \Rightarrow \frac{1}{s} \Rightarrow s^{-1}$$

Explanation: The frequency ν of this photon can be calculated by rearranging the equation $E = h\nu$ where E is the energy, h = Planck's constant and ν = frequency in s^{-1} . $\nu = E/h = 4.38 \times 10^{-18} / 6.63 \times 10^{-34} = 6.61 \times 10^{15}$

22. An electron in a Bohr hydrogen atom has energy of $-1.362 \times 10^{-19} J$. The value of n for this electron is _____.

- a) 1
- b) 2
- c) 3
- d) **4**

$$E_n = -R_H \left(\frac{1}{n^2} \right) \left(\frac{n^2}{1} \right) \Rightarrow \sqrt{n^2} = \sqrt{\frac{-R_H}{E_n}}$$

$$n = \sqrt{\frac{-2.18 \times 10^{-18} J}{-1.362 \times 10^{-19} J}} \Rightarrow n \approx 4$$

Explanation: The energy of an electron in a particular energy state in the hydrogen atom can be calculated by using the formula $E = (-2.18 \times 10^{-18} J)/n^2$, where n is the principal quantum number for the energy state. The value of n can be found by rearranging the above formula as follows:

$$n = \sqrt{\frac{-2.18 \times 10^{-18} J}{-1.362 \times 10^{-19} J}} = 4$$

19. The $n = 2$ to $n = 6$ transition in the Bohr hydrogen atom corresponds to the _____ of a photon with a wavelength of _____ nm.

- a) emission, 411
- b) **absorption, 411**
- c) absorption, 657
- d) emission, 389

$$\Delta E = R_H \left(\frac{1}{n_i^2} - \frac{1}{n_f^2} \right)$$

$$\Delta E = 2.18 \times 10^{-18} \left(\frac{1}{4} - \frac{1}{36} \right)$$

$$= 2.18 \times 10^{-18} (0.22) = 4.84 \times 10^{-19} J$$

$$E = h \frac{c}{\lambda}$$

$$4.84 \times 10^{-19} = 6.63 \times 10^{-34} \times \frac{3 \times 10^8}{\lambda}$$

$$\lambda = 4.109 \times 10^{-7} m = 4.11 \times 10^{-7} \times 10^9 = 411 \text{ nm}$$

Explanation: There are 2 parts to this question. Since the electron is moving from a smaller value of n (n_i) to a larger value of n (n_f), it must be absorbing energy. The wavelength responsible for this transition can be calculated by using the formula: $E = R_H (1/n_i^2 - 1/n_f^2)$ & $E = hc/\lambda$

20. How many quantum numbers are necessary to designate a particular electron in an atom _____?

- a) 3
- b) **4** $\rightarrow s, p, d$
- c) 2
- d) 1 n, l, m, m_s

21. The _____ quantum number defines the shape of an orbital.

- a) spin
- b) magnetic
- c) principal
- d) **angular** $\rightarrow l, p$

22. There are _____ orbitals in the ³ ⁿ third shell $n=3$


$$n^2 = 3^2 = 9$$

- a) 25
- b) 4
- c) 9
- d) 16

Explanation: The number of orbitals in a shell is easily calculated by the formula # of orbitals = n^2 where n = principal quantum number, which is 3 in this case.

23. The angular quantum number is 3 in _____ orbitals.

$$l=3 \Rightarrow f$$



- a) s
- b) p
- c) ~~d~~
- d) f ✓

24. The $n = 1$ shell contains _____ p orbitals. All the other shells contain _____ p orbitals.

- a) 3, 6
- b) 0, 3
- c) 6, 2
- d) 3, 3

Explanation: If $n = 1$, then the only possible value of l is 0 which means that $n = 1$ can contain only s orbitals. When $n > 1$, the value of $l = 1$ is possible making the existence of 3 p orbitals possible.

25. The principal quantum number of the first d subshell is _____.


- a) 1
- b) 2
- c) 3
- d) 4

26. The total number of orbitals in a shell is given by _____.

- a) L^2
- b) n^2
- c) $2n$
- d) $2n + 1$

28. Each p-subshell can accommodate a maximum of _____ electrons.

- a) 6
- b) 2
- c) 10
- d) 3



Explanation: There are 3 different p orbitals: p_x , p_y and p_z . Each of these can contain 2 electrons leading to the maximum number of electrons as 6.


29. Each p-subshell can accommodate a maximum of _____ electrons.

- a) 6
- b) 2
- c) 10
- d) 3

Explanation: There are 3 different p orbitals: p_x , p_y , and p_z . Each of these can contain 2 electrons leading to the maximum number of electrons as 6.

30. The $3p$ subshell in the ground state of atomic xenon contains _____ electrons.


- a) 2
- b) 6
- c) 36
- d) 10



Explanation: Since Xe is a noble gas, its subshells will be completely filled regardless of their principal quantum number. Thus the $3p$ subshell will contain 6 electrons.

31. $[\text{Ar}]4s^2 3d^{10} 4p^3$ is the electron configuration of a(n) _____ atom.


- a) As
- b) V
- c) P
- d) Sb



Explanation: The easiest way to answer this question is to count the total number of electrons and find which element that number corresponds to. The total number of electrons is = 18 (for the Ar) + 2 + 10 + 3 = 33 which corresponds to As.

32. The principal quantum number for the outermost electrons in a Br atom in the ground state is _____.

- a) 2
- b) 3
- c) 4
- d) 5



Explanation: The electronic configuration of bromine is $[\text{Ar}]3d^{10} 4s^2 4p^5$ shows that the outermost electrons are in the s and p orbitals in the 4th energy level making the principal quantum number = 4.

33. All of the _____ have a valence shell electron configuration ns^1 .

- a) noble gases
- b) halogens
- c) chalcogens
- d) alkali metals

34. Which one of the following is correct?

- a) $v + \lambda = c$
- b) $v/\lambda = c$
- c) $\lambda = cv$
- d) $v\lambda = c$


$$c = \lambda \times \nu$$

35. In the Bohr model of the atom, _____.

- a) electrons travel in circular paths called orbitals
- b) electrons can have any energy
- c) **electron energies are quantized**
- d) electron paths are controlled by probability

36. Which one of the following is not a valid value for the magnetic quantum number of an electron in a 5d subshell?


- a) 2
- b) 3
- c) 0
- d) 1



Explanation: For an electron in the 5d subshell the value of $\ell = 2$ and the magnetic quantum number $m\ell$ can have values from $-1, \dots, 0, \dots, +1$, meaning $m\ell$ could not have a value $\neq 3$.

37. Which of the subshells below do not exist due to the constraints upon the angular quantum number?


- a) 2s
- b) **2d**
- c) 2p
- d) none of the above



Explanation: The values of the azimuthal quantum number "l" are decided by the values of the principal quantum number "n". The values of l will only be from $0 \dots n - 1$. Thus for $n = 2$, only the values of 0 and 1 will be possible for l, which means that only the 2s and 2p orbitals will be possible.

38. An electron cannot have the quantum numbers $n =$ _____, $l =$ _____, $m\ell =$ _____.

- a) 2, 0, 0
- b) 2, 1, -1
- c) 3, 1, -1
- d) **1, 1, 1**







Explanation: The values of 1, 1, 1 would be impossible since if $n = 1$, the only value of ℓ would be $= 0$.

39. Which quantum number determines the energy of an electron in a hydrogen atom?

- a) **n**
- b) n and ℓ
- c) $m\ell$
- d) ℓ

39. Which electron configuration represents a violation of the Pauli exclusion principle?



Explanation: According to the Pauli Exclusion Principle no two electrons in an atom cannot have the same 4 quantum numbers. The 2 electrons in the 2s orbital have the same value for their m_s which is not allowed. (d)

40. Which of the following is a valid set of four quantum numbers? (n, ℓ, m_ℓ, m_s)

- a) 2, 0, 0, $+\frac{1}{2}$ ✓
- b) 2, 2, 1, $-\frac{1}{2}$ ✗
- c) 1, 0, 1, $+\frac{1}{2}$ ✗
- d) 2, 1, +2, $+\frac{1}{2}$ ✗

Explanation: Here is why only option (a) is the correct answer: In option (b), $\ell=2$ which is not allowed, in (c) $m_\ell \neq 1$ since $l=0$ and in (d) $m_\ell > 1$ which are all not allowed.

41. Which of the following is not a valid set of four quantum numbers? (n, ℓ, m_ℓ, m_s)

- a) 2, 0, 0, $+\frac{1}{2}$ ✓
- b) 2, 1, 0, $-\frac{1}{2}$ ✓
- c) 1, 1, 0, $+\frac{1}{2}$ ✗ $n-1 \text{ } \ell \text{ } m_\ell = \ell \text{ } \text{etc}$
- d) 1, 0, 0, $+\frac{1}{2}$ ✓

Explanation: Since n can never be equal to ℓ , option c is the only set that is not valid.

Test bank chapter (8)

Choose the most correct answer

1. Elements in the modern version of the periodic table are arranged in order of increasing _____.
- oxidation number
 - atomic mass
 - average atomic mass
 - atomic number**

Explanation: The older version of the periodic table had the elements arranged in order of increasing atomic mass, but the modern version of the periodic table is based on the increasing order of atomic number.

2. The first ionization energies of the elements _____ as you go from left to right across a period of the periodic table, and _____ as you go from the bottom to the top of a group in the table. Shaped like four-leaf clovers.

- increase, increase**
- increase, decrease
- decrease, increase
- decrease, decrease



Explanation: The ionization energies (IE s) of elements increase to the right in a row since larger amounts of energy need to be supplied to remove an electron. The elements become more non-metallic making it harder to remove an electron.



3. The _____ have the most negative electron affinities

- alkaline earth metals
- alkali metals
- halogens**
- transition metals

Explanation: The electron affinity of an element is defined as the energy change that occurs when an electron is added to a gaseous atom. The halogens have the most negative electron affinities indicating that they are most comfortable accepting an electron. The formation of an anion essentially gives the halogen atom the electron configuration of the nearest noble gas. The negative sign here indicates that the addition of an electron to the halogens results in energy being released by the halogen atom.

4. Na reacts with element X to form an ionic compound with the formula Na_3X . Ca will react with X to form _____.


- CaX_2
- CaX
- Ca_2X_3
- Ca_3X_2**



Explanation: The element X must have 3 negative charges for it to form the compound Na_3X , since each Ca has 2 positive charges, the formula of the compound formed by the reaction of Ca and X would have to be Ca_3X_2 .

5. 16. Atomic radius generally increases as we move _____.

- down a group and from right to left across a period**
- up a group and from left to right across a period
- down a group and from left to right across a period
- there is no trend



6. 18. The atomic radius of main-group elements generally increases down a group because _____.

- a) effective nuclear charge increases down a group
- b) effective nuclear charge decreases down a group
- c) both effective nuclear charge increases down a group and the principal quantum number of the valence orbitals increases
- d) **the principal quantum number of the valence orbitals increases**

7. Which of the following correctly lists the five atoms in order of increasing size (smallest to largest)?

- a) $O < F < S < Mg < Ba$
- b) **$F < O < S < Mg < Ba$**
- c) $F < O < S < Ba < Mg$
- d) $O < F < S < Ba < Mg$

in ↓ de →



Explanation: Fluorine and oxygen are in the same period (#2) and next to each other with F being the smallest of these 5 atoms. Ba is in group 2A and in row 6 (farthest "down" a group) and is the largest of the atoms. Mg is in group 2A and in the third period and hence will be bigger than F, O and S. Even though S is in the same period as Mg it is in group 6A making it smaller than Mg.

8. Which of the following correctly lists the five atoms in order of increasing size (smallest to largest)?

- a) $F < K < Ge < Br < Rb$
- b) $F < Ge < Br < K < Rb$
- c) $F < K < Br < Ge < Rb$
- d) **$F < Br < Ge < K < Rb$**

in ↓ de →




Explanation: Fluorine is in group 7A and period 2 making it the smallest of the 5 atoms here. Br is also in group 7A but is in period 4 making it larger than F, Ge is in group 4A and also in period 4 but is to the left of Br making it larger than Br. K and Rb are both in group 1A but K is in period 4 and Rb is in period 5, making the Rb atom the largest of all the 5 atoms.

9. Of the following atoms, which has the largest first ionization energy?

- a) Br
- b) **O**
- c) C
- d) P

in ↓ de →



Explanation: The ionization energy (IE) typically increases from left to right in a period and decreases from top to bottom in a group. Thus for C and O which are in the same period, O will have the larger IE. Br is in period #4 and will have the lowest IE out of these 4 elements.

10. Of the following elements, which has the largest first ionization energy?


- a) Na
- b) Al
- c) Se
- d) **Cl**

in ↓ de →

Explanation: The ionization energy (IE) typically increases from left to right in a period and decreases from top to bottom in a group. Na, Al and Cl are all in period 3 with the chlorine atom to the farthest right and will have the highest IE.

11. Which ion below has the largest radius?


- a) Cl^-
- b) K^+
- c) Br^-
- d) F^-



Explanation: Typically cations are smaller than their parent atoms while anions are larger than the parent atoms. Of the atoms here, the Br atom would be the largest as it is farthest down the group and hence its anion also will be the largest ion.

12. The ion with the smallest radius is _____.

- a) Br^-
- b) Cl^-
- c) O^{2-}
- d) F^-



Explanation: Typically anions are larger than the parent atoms. Of the atoms here, the F atom would be the smallest as it is farthest down the group and hence its anion also will be the smallest ion.

13. Which of the following is an isoelectronic series?


- a) $\text{B}^{5-}, \text{Si}^{4-}, \text{As}^{3-}, \text{Te}^{2-}$
- b) $\text{O}^{2-}, \text{F}^-, \text{Ne}, \text{Na}^+$
- c) S, Cl, Ar, K
- d) None of the above



Explanation: Isoelectronic series contain a combination of atoms and ions or only ions with the same number of electrons. Here the series containing the O^{2-} , F^- , Ne and Na^+ is the only one where all atoms/ions contain 10 electrons.

14. _____ is isoelectronic with argon and _____ is isoelectronic with neon.

- a) Cl^-, F^-
- b) Cl^-, Cl^+
- c) F^+, F^-
- d) Ne, Kr^+



Explanation: The Cl^- ion has 18 electrons and is isoelectronic with argon while the F^- ion has 10 electrons making it isoelectronic with neon.

15. Chlorine is much more apt to exist as an anion than is sodium. This is because _____.

- a) chlorine is bigger than sodium
- b) chlorine has a greater ionization energy than sodium does
- c) **chlorine has a greater electron affinity than sodium does**
- d) chlorine is a gas and sodium is a solid




16. The alkaline earth metals are found in _____ of the periodic table.

- a) Group 1A
- b) **Group 2A**
- c) Group 7A
- d) Group 8A

17. How many *valence electrons* does a boron atom (B) have?

- a) 1
- b) 3
- c) 5
- d) 7




18. Which ion is *isoelectronic* with Ar?

- a) $Ni^{2+} \rightarrow [Ar]_{18} 4s^0 3d^8$
- b) $F^- \rightarrow [Ne]$
- c) $Br^- \rightarrow [Kr]$
- d) $K^+ \rightarrow [Ar]$ ✓

19. Which of these choices is the electron configuration of the iron (III) ion (Fe^{3+})?

- a) $[Ar]4s^2 3d^5$
- b) $[Ar]4s^1 3d^5$
- c) $[Ar]3d^5$
- d) $[Ar]3d^6$




20. In what group of the periodic table is the element with the electron configuration $[Ar]4s^2 3d^{10} 4p^1$?

- a) 1A
- b) 2A
- c) 3A
- d) 4A

group = $2+1=3A$

21. How many *valence electrons* does a tin (Sn) atom have?

- a) 2
- b) 4
- c) 14
- d) 36



22. Which of these ground-state ions has unpaired electrons?

- a) $p^{3-} \rightarrow [Ar]$
- b) $V^{5+} \rightarrow [Ar]$
- c) $S^{2-} \rightarrow [Ar]$
- d) $Sc^{2+} \rightarrow 8[Ar]_{18} 4s^0 3d^1 \rightarrow$ unpaired

23. Consider the element with the electron configuration $[Xe]6s^2 4f^7$. This element is

- a) a representative element.
- b) a lanthanide element.
- c) a nonmetal.
- d) an actinide element

Period: 4
block: f


24. If the radius of atom X is greater than the radius of atom Y, then it is also likely that

- a) X has a larger electron affinity than Y does.
- b) X has a larger first ionization energy than Y does.
- c) X has greater metallic character than Y does.

25. Arrange these ions in order of increasing ionic radius: K^+ , P^{3-} , S^{2-} , Cl^- .

Increasing radius \rightarrow

- Row 1 $K^+ < Cl^- < S^{2-} < P^{3-}$
Row 2 $K^+ < P^{3-} < S^{2-} < Cl^-$
Row 3 $P^{3-} < S^{2-} < Cl^- < K^+$
Row 4 $Cl^- < S^{2-} < P^{3-} < K^+$
Row 5 $Cl^- < S^{2-} < K^+ < P^{3-}$



- a) Row 1
b) Row 2
c) Row 3
d) Row 4

26. - Selenium (${}_{34}Se$) element is

- a) a nonmetal
b) found in group 6A
c) found in period 2
d) **both a and b**

28. The outer electron configuration of the noble gases is

- a) $ns^2 np^6$
b) $ns^2 nd^{10}$
c) $ns^2 np^4$
d) $ns^2 np^8$

29. Which of the following species is isoelectronic with $Cl^- \rightarrow [Ar]_{18}$

- a) $K^+ \rightarrow [Ar]_{18}$
b) $Na^+ \rightarrow [Ne]_6$
c) O^{2-}
d) 2^-

30. Gallium (Ga) element is found in the periodic table in


- a) period 3, group 1B
b) period 3A, group 4
c) period 4, group 1A
d) **period 4, group 3A**

31. Titanium (Ti) element is found in the periodic table in

- a) s-block
b) p-block
c) **d-block**
d) f-block

32. Write the electronic configuration for Co^{+2}

- a) $[\text{Ar}] 4s^2 3d^5$
- b) $[\text{Ar}] 4s^2 3d^7$
- c) $[\text{Ar}] 4s^1 3d^6$
- d) $[\text{Ar}] 4s^1 3d^5$



33. Select the correct order of radius of the two ions

- a) $A^+ > A^-$ ✗
- b) $A^- < A^+$ ✗
- c) $A^{2+} > A^+$ ✗
- d) $A^{2+} < A^+$ ✓

لأن الأيونات
كلما فقدت إلكترونات كلما كبرت

34. Two ions are referred to as isoelectronic if they have the same number of



- a) **electrons.**
- b) protons.
- c) atoms.
- d) neutrons.

35. The energy required to remove an electron from an atom in its ground state is known as the

- a) potential energy.
- b) activation energy.
- c) electron affinity.
- d) **ionization energy.**

36. Which will have the highest ionization energy?

- a) C
- b) N
- c) O
- d) B





Exception $6A < 5A$

37. Order the following (N^{3-} , Li^+ , C , O^{2-}) according to increasing atomic/ionic radius.

- a) $\text{C} < \text{Li}^+ < \text{O}^{2-} < \text{N}^{3-}$
- b) $\text{N}^{3-} < \text{O}^{2-} < \text{C} < \text{Li}^+$
- c) $\text{Li}^+ < \text{C} < \text{N}^{3-} < \text{O}^{2-}$
- d) **$\text{Li}^+ < \text{C} < \text{O}^{2-} < \text{N}^{3-}$**

كلما فقدت إلكترونات كلما كبرت
 كلما اكتسبت إلكترونات كلما صغر



Test bank chapter (9)

س١٥
س١٢
NO₃⁻ صحیح جواب ہے
Satisfied نامعلوم جواب ہے

Choose the most correct answer

1. The two types of chemical bonds commonly found in compounds are:

- a) ionic and covalent.
- b) ionic and electrolytic.
- c) **ionic and covalent.**
- d) electrolytic and compound.

انواع الرابطة الكيميائية
↓ ↓
تساهلية أيونية

2. The electrons used by atoms to form chemical bonds are the:

- a) core electrons.
- b) **valence electrons.**
- c) lone pair electrons.
- d) unpaired electrons.

3. "Atoms tend to gain, lose, or share electrons until they are surrounded by eight valence electrons" is a statement of:

- a) the rule of octaves.
- b) the double quartet rule.
- c) the eight electron rule.
- d) **the octet rule.**


4. When a transition metal atom becomes a +1 ion, the electron lost usually comes from what type of orbital?

- a) p
- b) f
- c) d
- d) **s**

$ns(n-1)d$
↑
تسحب من s اول

5. A molecule of CS₂ contains

- a) two single bonds.
- b) **two double bonds.**
- c) one single bond and one double bond.
- d) one single bond and one triple bond.



6. An atom in the ground state has atomic number $Z=5$. Choose the correct electron-dot structure which represents this atom? **ANS. B**

- (A) $\cdot \ddot{X} \cdot$
- (B) $\ddot{X} \cdot$
- (C) $\cdot \ddot{X} \cdot$
- (D) $\cdot \ddot{X} \cdot$

$$1s^2 2s^2 2p^1 \Rightarrow 2+1=3$$



7. Which compound below contains an atom that is surrounded by more than an octet of electrons?

- a) PF₅ له 5 ذرات نسوي الكثر من ال octet
- b) CH₄
- c) NBr₃
- d) OF₂

8. Which choice below correctly lists the elements in order of increasing electronegativity?

- a) C < N < O < F
 - b) N < C < O < F
 - c) N < C < F < O
 - d) C < N < F < O
- C N O F
↑
in

9. Which atom sometimes violates the octet rule?

- a) C
- b) N
- c) O
- d) S

10. How many resonance structures can be drawn for NO₃⁻?

- a) 1
 - b) 2
 - c) 3
 - d) 4
- $A = 3 \times 6 + 5 + 1 = 24 e^-$
- $\text{:}\ddot{\text{O}}-\text{N}=\ddot{\text{O}}\text{:} \leftrightarrow \ddot{\text{O}}=\text{N}-\ddot{\text{O}}\text{:} \leftrightarrow \text{:}\ddot{\text{O}}-\text{N}-\ddot{\text{O}}\text{:}$

11. Considering formal charge, what is the preferred Lewis structure of NCO? **ANS. 1**

- 1. $\text{:}\ddot{\text{N}}\equiv\text{C}-\ddot{\text{O}}\text{:}$ 4. $\text{:}\ddot{\text{N}}-\text{C}-\ddot{\text{O}}\text{:}$
- 2. $\text{:}\ddot{\text{N}}=\text{C}=\ddot{\text{O}}\text{:}$ 5. $\text{:}\ddot{\text{N}}=\text{C}::\ddot{\text{O}}\text{:}$
- 3. $\text{:}\ddot{\text{N}}-\text{C}::\ddot{\text{O}}\text{:}$

12. In Lewis structure of (SO₄)²⁻ structure the correct formal charge on sulfur (S) is:

- a) ~~+2~~
 - b) -2
 - c) +1
 - d) 0
- $\text{O}=\text{S}=\text{O}$
 $\text{O}=\text{S}-\text{O}$
- Satisfies لانه ما قال Satisfies \Rightarrow فيه ذرات بس \Rightarrow متسوية بس لونها Satisfies \Rightarrow متسوية بس لونها octet rule \Rightarrow متسوية بس لونها octet rule

13. Which of these pairs of elements would be most likely to form an ionic compound?

- a) Cl and I \rightarrow non, non
 - b) Al and K \rightarrow metal, metal
 - c) **Cl and Mg** \rightarrow metal, nonmetal
 - d) C and S \rightarrow non, non
- metal, non metal

قوله اذ ان الاكثر كبر السالبه
وهو الايونيك

14. Which of these covalent bonds is the most polar (i.e., highest percent ionic character)?

- a) $\overset{\text{metal}}{\text{Al}}-\text{I}$ اقول
 - b) $\text{Si}-\text{I}$ مستقر
 - c) $\text{Al}-\text{Cl}$ اكبر
 - d) $\text{Si}-\text{Cl}$ مستقر
- \downarrow metals

15. The Lewis structure for CS₂ is: ANS. c

- a) $\ddot{\text{C}}=\ddot{\text{S}}-\ddot{\text{S}}$
 - b) $\ddot{\text{S}}-\ddot{\text{C}}-\ddot{\text{S}}$ more stable
 - c) $\boxed{\ddot{\text{S}}=\text{C}=\ddot{\text{S}}}$ استقرار اكثر
 - d) $\ddot{\text{S}}-\text{C}=\ddot{\text{S}}$
- $S = 6 - (4 + 2) = 0$
 $C = 4 - (4 + 0) = 0$

16. The number of lone electron pairs in the N₂ molecule is ____.

- a) 1
 - b) 2
 - c) 3
 - d) 4
- $:\text{N} \equiv \text{N}:$

17. Classify the $\text{O}-\text{H}$ bond in CH₃OH as ionic, polar covalent, or nonpolar covalent.


- a) Ionic
- b) polar covalent
- c) nonpolar covalent
- d) none of the above

18. The Lewis structure for a chlorate ion, ClO₃⁻, should show ____ single bond(s), ____ double bond(s), and ____ lone pair(s).

- a) 2, 1, 10
 - b) 3, 0, 9
 - c) 2, 1, 8
 - d) 3, 0, 10
- ClO_3^-
- A) $(1 \times 7) + (3 \times 6) + 1 = 26e$
- B) $= (1 \times 8) + (3 \times 8) = 32e$
- C) $B - A = 32 - 26 = \frac{6e}{2} = 3 \text{ bond}$
- $\text{O} = \text{Cl} - \text{O}$
- O
- D) $A - C = 26 - 6 = 20e$
non bonding e or lone pair


19. The number of resonance structures for the sulfur dioxide (SO₂) molecule that satisfy the octet rule is

- a) 1
 - b) 2
 - c) 3
 - d) None of these.
- SO_2
- A) $= 6 + 2 \times 6 = 18e$




20. What is the formal charge on the oxygen atom in N_2O (the atomic order is N-N-O)?

- a) 0
- b) +1
- c) -1
- d) -2



21. Which of these substances will display an incomplete octet in its Lewis structure?

- a) CO_2
- b) Cl_2
- c) ICl
- d) NO



22. There are 2 paired and 3 unpaired electrons in the Lewis symbol for a phosphorus atom (P).


- a) 4, 2
- b) 2, 4
- c) 4, 3
- d) 2, 3

Explanation: Read the question carefully here, you are being asked for how many valence electrons are paired and how many are unpaired. The abbreviated electron configuration of the P atom is given by $[\text{Ne}] 3s^2 3p^3$. The outermost electrons would be arranged as 2 electrons paired and 3 electrons unpaired as shown below:



23. Based on the octet rule, magnesium (Mg) most likely forms a _____ ion.


- a) Mg^{2-}
- b) Mg^{2+}
- c) Mg^{6-}
- d) Mg^{6+}



Explanation: According to the octet rule the Mg atom will achieve an octet by losing its 2 outermost electrons and thus gaining 2+ charges. Since Mg is located in the alkali metal group it will lose electrons rather than gaining them.

24. Based on the octet rule, phosphorus (P) most likely forms a P^{-3} ion.

- a) P^{3+}
- b) P^{5-}
- c) P^{5+}
- d) P^{3-}



Explanation: According to the octet rule the phosphorus atom should gain 3 electrons, thus gaining 3 negative charges and forming the phosphide ion.

25- The only noble gas without eight valence electrons is _____.

- a) Ar
- b) Ne
- c) He
- d) Kr

الغضيلوم لهما الذرية الالكترونية
موزون اذواتو

Explanation: The noble gases are characterized by the presence of eight electrons in their outermost shell with one notable exception of Helium. Since He has only 2 electrons it can never have 8 in its outermost shell.

26- What is the maximum number of double bonds that a hydrogen atom (H) can form?

- a) 0
- b) 1
- c) 2
- d) 3

بسي يشارك لاد 2e
بسي Single بس

Explanation: Each hydrogen atom has a single electron in its valence shell and as a result can form only one bond. It cannot form a double bond as it does not have the necessary electrons to share.

28. What is the maximum number of double bonds that a carbon atom (C) can form?

- a) 4
- b) 1
- c) 2
- d) 0

=C= ≡C-
واحد مستقل ، واحد تويل ، 2 ديل

Explanation: Each carbon atom has 4 valence electrons that it can share with other atoms. Since each double bond corresponds to a pair of electrons, the carbon atom can form only 2 double bonds.

29. Given the electronegativities below, which covalent single bond is most polar?

Atom	H	C	N	O
Electronegativity	2.1	2.5	3.0	3.5

- a) C-H $2.5 - 2.1 = 0.4$ Polar \Rightarrow Non
- b) N-H $3 - 2.1 = 0.9$ Polar
- c) O-H $3.5 - 2.1 = 1.4$ Polar
- d) O-N $3.5 - 3 = 0.5$ Polar

Explanation: Bond polarity can be judged based on the differences between the electronegativities of the atoms involved. Of the available choices, the bond between O and H will have the largest electronegativity difference making it the most polar bond in this group.

30. The ion ICl_4^- has _____ valence electrons.

- a) 34
- b) 36
- c) 35
- d) 28

ICl_4^-
 $(1 \times 7) + (4 \times 7) + 1 = 36e$

Explanation: valence electrons A = $(7 \times 1) + (7 \times 1) + 1 = 36$


31- Electronegativity _____ from left to right within a period and _____ from top to bottom within a group.

- a) decreases, increases
- b) increases, increases
- c) stays the same, increases
- d) **increases, decreases**

Explanation: Atomic size decreases from the left to the right in a period thus making it easier for the nuclei to attract electrons towards themselves resulting in an increase in the electronegativity. On the other hand atomic size increases down a group making it harder for the nuclei to attract the valence electrons towards themselves resulting in a decrease in electronegativity.

32. The Lewis structure of PF_3 shows that the central phosphorus atom has ___ nonbonding and ___ bonding electron pairs.

- a) 2, 2
- b) 1, 3
- c) 3, 1
- d) 1, 2



$$A = (1 \times 5) + (3 \times 7) = 26e$$


$$B = (1 \times 8) + (3 \times 8) = 32e$$

$$C \quad B - A = 32 - 26 = \frac{6e}{2} = 3 \text{ bonds}$$

$$D \quad A - C = 26 - 6 = 20 \text{ nonbonding } e$$

33. Which of the following molecules contains both ionic and covalent bonds?

- a) C_3H_{12} non, non
- b) **$NaClO_4$**
- c) $CaCl_2$ non, metal
- d) H_2O non, non




34. The ability of an atom in a molecule to attract electron density to itself is termed

- a) **Electronegativity**
- b) Electron affinity
- c) Diamagnetism
- d) Ionization energy

35- the most polar bond is

- a) Br-H
- b) I-H
- c) **Cl-H**
- d) H-H




سؤال الإجابة (ب) كالمعتاد

Test bank chapter (14)

b → CO
c → CO₂

Choose the most correct answer

1. Which is the correct equilibrium constant expression for the following reaction?



- a) $K_c = [Fe_2O_3][H_2]^3/[Fe]^2[H_2O]^3$
- b) $K_c = [H_2]/[H_2O]$
- c) $K_c = [H_2O]^3/[H_2]^3$
- d) $K_c = [Fe]^2[H_2O]^3/[Fe_2O_3][H_2]^3$

$K_c = \frac{[H_2O]^3}{[H_2]^3}$

ملاحظة
المواد الصلبة والسائلة لا يكتب من معادلات الاتزان
فقط "g" فقط

2. The following reactions occur at 500 K. Arrange them in order of increasing tendency to proceed to completion (least → greatest tendency).

- 1. $2NOCl \rightleftharpoons 2NO + Cl_2$
- 2. $2SO_3 \rightleftharpoons 2SO_2 + O_2$
- 3. $2NO_2 \rightleftharpoons 2NO + O_2$

$K_p = 1.7 \times 10^{-2}$ ① 0.017
 $K_p = 1.3 \times 10^{-3}$ ② 0.00013
 $K_p = 5.9 \times 10^{-3}$ ③ 0.0059

كلما كانت قيمة K أكبر
كان ميل التفاعل لإتمامه
النواتج أكبر

- a) 2 < 1 < 3
- b) 1 < 2 < 3
- c) 2 < 3 < 1
- d) 3 < 2 < 1

1 > 3 > 2

3. Calculate K_p for the reaction $2NOCl(g) \leftrightarrow 2NO(g) + Cl_2(g)$ at 400°C if K_c at 400°C for this reaction is 2.1×10^{-2} .

- a) 2.1×10^{-2}
- b) 1.7×10^{-3}
- c) 0.70
- d) 1.2

$K_p = \frac{[Cl_2][NO]^2}{[NOCl]^2}$

$K_p = K_c (RT)^{\Delta n}$

$\Delta n = (2+1) - 2 = 3 - 2 = 1$
 $K_p = 2.1 \times 10^{-2} (0.082 \times 673.15)^1 = 1.16 \approx 1.2$

4. For the reaction $H_2(g) + I_2(g) \leftrightarrow 2HI(g)$, $K_c = 50.2$ at 445°C. If $[H_2] = [I_2] = [HI] = 1.75 \times 10^{-3} M$ at 445°C, which one of these statements is true?

- a) The system is at equilibrium, thus no concentration changes will occur.
- b) The concentrations of HI and I₂ will increase as the system approaches equilibrium.
- c) The concentration of HI will increase as the system approaches equilibrium.
- d) The concentrations of H₂ and HI will fall as the system moves toward equilibrium.


$Q_c = \frac{[HI]^2}{[H_2][I_2]}$
 $= \frac{(1.75 \times 10^{-3})^2}{(1.75 \times 10^{-3})(1.75 \times 10^{-3})}$
 $= 1$

5. For the following reaction at equilibrium, which choice gives a change that will shift the position of equilibrium to favor formation of more products?



- a) Increase the total pressure by decreasing the volume. → حيز حجوم أقل
- b) Add more NO. → تزييد الناتج
- c) Remove Br₂. → حيز حجوم أكبر
- d) Lower the temperature. → تنخفض الحرارة حيز حجوم أكبر

6 - For the following reaction at equilibrium in a reaction vessel, which one of these changes would cause the Br₂ concentration to decrease?



- a) Increase the temperature. → يزداد لليمين X
- b) Remove some NO. → يزداد لليسار X
- c) Add more NOBr. → يزداد لليسار X
- d) Compress the gas mixture into a smaller volume. (يضيء لليسار) ضغط الحجم يزداد لفرق الأجزاء

7. For the reaction at equilibrium $2\text{SO}_3 \leftrightarrow 2\text{SO}_2 + \text{O}_2$ ($\Delta H^\circ_{\text{rxn}} = 198 \text{ kJ/mol}$), if we increase the reaction temperature, the equilibrium will

endo متفاعلات
توزاد حجومها لليمين

- a) shift to the right.
- b) shift to the left.
- c) not shift.
- d) The question cannot be answered because the equilibrium constant is not given.

8. For the equilibrium reaction $2\text{SO}_2\text{(g)} + \text{O}_2\text{(g)} \leftrightarrow 2\text{SO}_3\text{(g)}$, $\Delta H^\circ_{\text{rxn}} = -198 \text{ kJ/mol}$. Which one of these factors would cause the equilibrium constant to increase?

- a) Decrease the temperature.
- b) Add SO₂ gas.
- c) Remove O₂ gas.
- d) Add a catalyst.

	exo	endo
T increase	de K	in K
T decrease	in K	de K

في النواتج EXO
ما هي صيغته يتكافئ
بتأثير الإلتزان بـ Kc
! لا الحرارة!

9. The reaction $2\text{SO}_3\text{(g)} \leftrightarrow 2\text{SO}_2\text{(g)} + \text{O}_2\text{(g)}$ is endothermic. If the temperature is increased

- a) more SO₃ will be produced.
- b) K_c will decrease.
- c) no change will occur in K_c.
- d) K_c will increase.

ال K يزداد

10. If a catalyst is added to a chemical reaction, the equilibrium yield of a product will be ... , and the time taken to come to equilibrium will be than before.

- a) higher; less
- b) lower; the same
- c) higher; the same
- d) the same; less

المحفزات لا تؤثر على الإلتزان
بسبب تزييد سرعة التفاعل فقط
وإنها قلصت أقل

11- For the reaction, $\text{N}_2\text{(g)} + 3 \text{H}_2\text{(g)} \rightleftharpoons 2 \text{NH}_3\text{(g)}$

K_c = 0.0600 at a certain temperature. In an equilibrium mixture of the three gases, [NH₃] = 0.242 M and [H₂] = 1.03 M. What is the concentration of N₂ in this system?

- a) 3.9 M
- b) 3.2 x 10⁻³ M
- c) 0.89 M
- d) 1.4 x 10⁻² M

$$K_c = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$$

$$0.0600 = \frac{(0.242)^2}{\text{N}_2(1.03)^3} \Rightarrow \text{N}_2 = \frac{(0.242)^2}{0.0600(1.03)^3}$$

2

$$\text{N}_2 = 0.893 \text{ M}$$

11. Consider the reaction $\text{NH}_4\text{Cl}(s) \rightleftharpoons \text{NH}_3(g) + \text{HCl}(g)$.

If an equilibrium mixture of these three substances is compressed, equilibrium will _____, because _____.

- a) shift to the right; higher pressure favors fewer moles of gas
- b) shift to the right; higher pressure favors more moles of gas
- c) **shift to the left; higher pressure favors fewer moles of gas**
- d) shift to the left; higher pressure favors more moles of gas

الضغط يزداد يقل الحجم
يرجع لعدد المولات الأقل
يعني لليسار

12- Consider the equilibrium system; $\text{C}(s) + \text{CO}_2(g) \rightleftharpoons 2\text{CO}(g)$.

If more $\text{C}(s)$ is added, the equilibrium will _____; if CO is removed the equilibrium will _____.

- a) shift to the left; shift to the left
- b) shift to the right; shift to the right $\rightarrow \text{CO}$
- c) **shift to the right; shift to the left** $\rightarrow \text{CO}_2$
- d) be unchanged; shift to the left

- زدنا المتفاعلات
- راج اليمين
- قلنا CO (نواتج)
- راج اليمين

8. Consider the exothermic reaction at equilibrium: $2\text{SO}_2(g) + \text{O}_2(g) \rightleftharpoons 2\text{SO}_3(g)$, If the system is cooled, the equilibrium will _____, because _____.

- a) shift to the left; decreased temperature favors an exothermic reaction
- b) **shift to the right; decreased temperature favors an exothermic reaction**
- c) shift to the right; decreased temperature favors an endothermic reaction
- d) shift to the left; decreased temperature favors an endothermic reaction

Exothermic \rightarrow نواتج
لو زدنا الحرارة حيرج اليسار
لو خفضنا الحرارة حيرج اليمين


9. A large value of the equilibrium constant indicates that when the reaction reaches equilibrium, mostly _____ will be present.

- a) reactants
- b) **products**
- c) catalysts
- d) shrapnel

$K > 1 \rightarrow \text{product}$
 $K < 1 \rightarrow \text{reactant}$


10. When equilibrium is achieved

- a) $Q > K$
- b) $Q < K$
- c) $Q = K$
- d) $Q_2 = K$




If all species are gases and H₂ is added, the amount of CO present at equilibrium will:

- a) increase.
- b) decrease.
- c) remain unchanged.
- d) disappear.



If all species are gases and H₂O is added, the amount of CO present at equilibrium will:


- a) increase.
- b) decrease.
- c) remain unchanged.
- d) disappear.



If the reaction is endothermic and the temperature is raised, the amount of CO present will:

- a) increase.
- b) decrease.
- c) remain unchanged.
- d) disappear.

Handwritten note: "زاد يبقى للمين" (CO increases).




If all species are gases and the container is compressed, the amount of CO present will:

- a) increase.
- b) decrease.
- c) remain unchanged.
- d) disappear.

Handwritten note: "زاد الضغط يقل الحجم" (pressure increases, volume decreases) with an arrow pointing to the reaction.

$$n = 2 - 2 = 0$$

Handwritten note: "كله متعادل ما يتأثر" (all are balanced, no effect).



If all species are gases and the container is compressed, the amount of CO present will:

- a) increase.
- b) decrease.
- c) remain unchanged.
- d) disappear.

Handwritten note: "زودنا ضغطه حجم يقل" (we increase pressure, volume decreases) with an arrow pointing to the reaction.

Handwritten note: "ما حقيقاً نزل عدد المولات (متساوي)" (not really, number of moles is equal).

16. What is K_p in terms of K_c for the following reaction? $2\text{NO}(g) + \text{O}_2(g) \leftrightarrow 2\text{NO}_2(g)$

- a) $K_p = K_cRT$
- b) $K_p = K_c/RT$
- c) $K_p = K_cR/T$
- d) $K_p = K_c/(RT)^2$

$$K_p = K_c (RT)^{\Delta n}$$

$$K_p = K_c (RT)^{-1}$$

Handwritten note: "الاسي السالبي ينزل له مقام" (negative exponent goes to denominator).

$$\Delta n = 2 - (1+2) = 2 - 3 = -1$$

$$K_p = \frac{K_c}{RT}$$

17. What is the correct equilibrium constant expression for the reaction: **ANS. 3**



- | | |
|--------------------------------------|--|
| 1. $\frac{[PCl_3]^4}{[P_4][Cl_2]^6}$ | 4. $\frac{[Cl_2]^6}{[PCl_3]^4}$ |
| 2. $\frac{[PCl_3]^4}{[Cl_2]^6}$ | 5. $\frac{[4 PCl_3]^4}{[P_4][6 Cl_2]^6}$ |
| 3. $\frac{1}{[Cl_2]^6}$ | |

18. The equation relating K_p and K_c is


- a) $K_p = K_c (RT)^{\Delta n}$
 b) $K_p = K_c RT^{\Delta n}$
 c) $K_c = K_p RT^{\Delta n}$
 d) $K_c = K_p (RT)^{\Delta n}$

(الاقواس تعرف في الحد)
 (القانون ثابت)

19. K_p will be equal to K_c if

- a) $\Delta n = 1$
 b) $\Delta n = 0$
 c) $RT = 0$
 d) $\Delta n = -1$

20. Consider the reversible reaction at equilibrium at 392 C:




The partial pressures are found to be: A: 6.70 atm, B: 10.1 atm, C: 3.60 atm. Evaluate K_p for this reaction.

- a) 7.94×10^{-3}
 b) 0.146
 c) 0.0532
 d) 54.5

$$K_p = \frac{P_C}{P_A^2 P_B} = \frac{3.60}{(6.70)^2 (10.1)} = 7.94 \times 10^{-3}$$

20. Which of the following will result in an equilibrium shift to the right?




- a) Increase temperature/increase volume
 b) Increase temperature/decrease volume
 c) Decrease temperature/increase volume
 d) Decrease temperature/decrease volume

↑ الحرارة
 بالنواتج

× ←
 × ←
 ✓ ←
 →
 توزيع لعدد المولات المتساوي
 → اليسار

21. Which accurately reflects the changes in concentration that will occur if O_2 is added to disturb the equilibrium?



	[NO]	[O ₂]	[NO ₂]
a)	Increase	Increase	Increase
b)	Increase	Increase	Decrease
c)	Decrease	Decrease	Decrease
d)	Decrease	Increase	Increase

Test bank chapter (15)

شماره الجواب (40) صحیح
لغوی 2.08×10^{-13}

Choose the most correct answer

1-What is the concentration of H^+ in a 2.5 M HCl solution?

- a) 0
b) 1.3 M
c) 2.5 M
d) 5.0 M

هو نفس ما كانا (نفس)

2. What is the OH^- ion concentration in a 5.2×10^{-4} M HNO_3 solution?

- a) 1.9×10^{-11} M
b) 1.0×10^{-7} M
c) 5.2×10^{-4} M
d) Zero

$[H^+]$

القانون $K_w = [OH^-][H^+]$

$$[OH^-] = \frac{1 \times 10^{-14}}{5.2 \times 10^{-4}} = 1.9 \times 10^{-11}$$

3. Calculate the H^+ ion concentration in lemon juice having a pH of 2.4.

- a) 4.0×10^{-2} M
b) 250 M
c) 0.38 M
d) 4.0×10^{-3} M

$$[H^+] = 10^{-pH}$$

$$[H^+] = 10^{-2.4}$$

$$= 3.9 \times 10^{-3} \approx 4 \times 10^{-3} \text{ M}$$

4. Calculate the pH of a 6.71×10^{-2} M NaOH solution.

- a) 12.83
b) 2.17
c) 11.82
d) 6.71

$[OH^-]$

$K_w = [OH^-][H^+]$

$$[H^+] = \frac{1 \times 10^{-14}}{6.71 \times 10^{-2}} = 1.49 \times 10^{-13}$$

$$pH = -\log[H^+] \Rightarrow pH = -\log[1.49 \times 10^{-13}] = 12.83$$

طريقة ثانية:

$$pOH = -\log(6.71 \times 10^{-2})$$

$$= 1.17$$

$$pH + pOH = 14 \Rightarrow pH = 14 - 1.17 = 12.83$$

5. What is the pH of 0.0200 M aqueous solution of HBr ?

- a) 1.00
b) 1.70
c) 2.30
d) 12.30

$[H^+]$

$$pH = -\log[H^+] \Rightarrow pH = -\log(0.0200) = 1.699 \approx 1.70$$

6. The pOH of a solution of NaOH is 11.30. What is the $[H^+]$ for this solution?

- a) 2.0×10^{-3}
b) 2.5×10^{-3}
c) 5.0×10^{-12}
d) 4.0×10^{-12}

pOH

$$pOH + pH = 14$$

$$pH = 14 - pOH$$

$$pH = 14 - 11.30 = 2.7$$

$$[H^+] = 10^{-pH}$$

$$[H^+] = 10^{-2.7} = 1.995 \times 10^{-3}$$

$$\approx 2 \times 10^{-3}$$

7. What is the pH of a 0.0400 M aqueous solution of KOH?

- a) 12.60
b) 10.30
c) 4.00
d) 1.40

$$K_w = [H^+][OH^-]$$

$$[H^+] = \frac{1 \times 10^{-14}}{0.0400} = 2.5 \times 10^{-13}$$

$$pH = -\log[H^+] = -\log(2.5 \times 10^{-13}) = 12.60$$

طريقة اخرى (7)

$$pOH = -\log(0.04) = 1.38$$

$$pH + pOH = 14 \Rightarrow pH = 14 - 1.38 = 12.62$$

8. What is the approximate pH of a solution labeled 6×10^{-5} M HBr?

- a) 4.2
b) 4.5
c) 5.8
d) 9.8

$$pH = -\log[H^+]$$

$$pH = -\log(6 \times 10^{-5}) = 4.2$$

9. If the pH = 2 for an HNO₃ solution, what is the concentration of HNO₃?

- a) 0.10
b) 0.20
c) 0.010
d) 0.020

$$[H^+] = 10^{-pH}$$

$$= 10^{-2} = 0.01$$

10. A solution in which $[H^+] = 10^{-8}$ M has a pH of 8 and is basic

- a) 8, acidic
b) 6, basic
c) -6, basic
d) 8, basic

$$pH = -\log[H^+] \Rightarrow [H^+] = 10^{-pH}$$

$$= -\log(10^{-8}) = 8 > 7 \rightarrow \text{basic}$$

11. Which of the following solutions has the lowest pH at 25°C? (No calculations required.)

- a) 0.2 M NaOH → قاعدية
b) 0.2 M NH₃ → قاعدية
c) 0.2 M HCl → حمضية
d) pure water → متعادلة

قاعدة
PH > 7

حمضية
PH < 7

متعادلة
7

12. Calculate the pH of a 3.5×10^{-3} M HNO₃ solution.

- a) -2.46
b) 0.54
c) 2.46
d) 3.00

$$pH = -\log[H^+]$$

$$= -\log(3.5 \times 10^{-3}) = 2.46$$

13. The pH of 2.6×10^{-2} M KOH is

- a) 12.41
b) 15.59
c) 2.06
d) 7.00

$$pOH = -\log(2.6 \times 10^{-2}) = 1.585$$

$$pH + pOH = 14$$

$$pH = 14 - 1.59 = 12.41$$

طريقة اخرى

$$K_w = [H^+][OH^-]$$

$$[H^+] = \frac{K_w}{[OH^-]}$$

$$= \frac{1 \times 10^{-14}}{2.6 \times 10^{-2}} = 3.846 \times 10^{-13}$$

$$pH = -\log[H^+]$$

$$= -\log(3.846 \times 10^{-13}) = 12.41$$

14. What is the $[H^+]$ ion in a 4.8×10^{-2} M KOH solution?

- a) 2.8×10^{-13} M
- b) 1×10^{-7} M
- c) 4.8×10^{-11} M
- d) 4.8×10^{-2} M

$[OH^-]$

$$K_w = [H^+][OH^-]$$
$$[H^+] = \frac{1 \times 10^{-14}}{4.8 \times 10^{-2}} = 2.083 \times 10^{-13}$$

15. What is the $[OH^-]$ ion in a 5.2×10^{-4} M HNO₃ solution?

- a) 1.9×10^{-11} M
- b) 1.0×10^{-7} M
- c) 5.2×10^{-4} M
- d) zero

$[H^+]$




$$K_w = [H^+][OH^-]$$
$$[OH^-] = \frac{1 \times 10^{-14}}{5.2 \times 10^{-4}} = 1.9 \times 10^{-11} \text{ M}$$

Test bank chapters (24 & 25)

Choose the most correct answer

1. $C_{10}H_{22}$ is the formula of an

- a) alkane.
- b) alkene.
- c) alkyne.
- d) aromatic hydrocarbon.





2. A molecule with the formula C_3H_8 is a

- a) hexane
- b) propane
- c) decane
- d) butane

بروبان

3. Which compound below does not have geometric isomers (cis-trans isomers)?


- a) 1-butene \rightarrow بدون الارتباط
- b) 2-butene \rightarrow $\text{التي لا يوجد كيزومرات}$
- c) 2-pentene
- d) 3-hexene



4. The hybridization of carbon atoms in alkanes is

- a) sp
- b) sp^2
- c) sp^3
- d) sp^3d

5. Select the correct IUPAC name for



- a) 1,1,3-trimethylpentane
- b) 1-ethyl-1,3-dimethylbutane
- c) 2,4-dimethylhexane
- d) 3,5-dimethylhexane.

2,4-dimethylhexane

6 - An alkane with seven carbon atoms in a linear configuration is called a

- a) hexene
- b) heptene
- c) heptylane
- d) **heptane**




7. Which type of functional group does not include a carbonyl group in its structure?


- a) carboxylic acid \rightarrow $\begin{matrix} O \\ || \\ C-OH \end{matrix}$
- b) ether $R-O-R$
- c) ketone \leftarrow $\begin{matrix} O \\ || \\ C-R \end{matrix}$
- d) aldehyde \rightarrow $\begin{matrix} O \\ || \\ C-H \end{matrix}$



8. Vanillin is used as a flavoring agent. Identify the functional group circled.



- a) aldehyde
- b) ketone
- c) carboxylic acid
- d) Alcohol




9. The formula $(\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH})$ represents:

- a) an alcohol
- b) an alkene
- c) an alkyne
- d) an unsaturated hydrocarbon

10. Which of the following is a ketone?

- a) $\text{CH}_3\text{CH}_2\text{COCH}_3$
- b) $\text{CH}_3\text{CH}_2\text{CHO}$
- c) CH_3OCH_3
- d) $\text{CH}_3\text{CH}_2\text{COOH}$




11. The general formula for alkenes is

- a) $\text{C}_n\text{H}_{2n+2}$
- b) C_nH_{2n}
- c) C_nH_{n+2}
- d) C_nH_{2n}




12- Which of these molecules is unsaturated?

- كثير مشبع يعني مشبع *single bond*, *single bond*
 (عملياً العامة formula)
 a) C_3H_8
 b) CH_3OH
 c) C_5H_{10}
 d) CH_4 → ايثان




8. Which of these species is an aromatic compound?

- a) C_2H_2
- b) C_6H_{12}
- c) $\text{C}_6\text{H}_4\text{Br}_2$
- d) C_5H_{10}




هو هيدروكربون عطري
 - الصيغة العامة البنزين
 C_6H_6
 اهم شيء ستة كربونات
 وترتبطه بستة ذرات هيدروجين

9. Which of these species are structural isomers of C_6H_{14} ?




7C X




6C ✓

نفس عدد ذرات آکسیژن



7C X




6C ✓

- a) I and II
- b) I and III
- c) II and III
- d) II and IV


10. The compound that has a triple bond between one pair of carbon atoms is called

- a) an alkane.
- b) a chlorofluorocarbon.
- c) an alkyne
- d) an alkene.


11. The correct structure for 2,3,3-trimethylpentane is




A.




B.








C.



D.




12. Which one of these structures represents an ester functional group?



13. The functional group (RCOR) is characteristic of organic _____

- a) ketones
- b) acids
- c) aldehydes
- d) esters




14. Which of the following hydrocarbons does not have isomers?

- a) C₇H₁₆
- b) C₆H₁₄
- c) C₅H₁₀
- d) C₃H₈

*يوجدان وصلات وايضا
لا زلهم هيفار مقدر
احط سيج بالام*

15. Which of the following does NOT exhibit geometric isomerism? (Hint: draw them!)

- a) 4-octene
- b) 2-pentene
- c) 3-hexene
- d) 1-hexene




16. For which of the compounds below are cis-trans isomers possible?


*في الرابطة
الثنائية
في الطرف*

- 1) CH₃CH=CH₂ ×
- 2) CH₃CH=CHCH₂CH₃
- 3) CH₃CH=CHCH₃

- a) only 2
- b) both 1 and 2
- c) **both 2 and 3**
- d) all three



17. Select the correct IUPAC name for



2, 4 - dimethylhexane


- a) 1,1,3-trimethylpentane
- b) 1-ethyl-1,3-dimethylbutane
- c) 2,4-dimethylhexane
- d) 3,5-dimethylhexane

18. A protein is:

- a) a polymer of amino acids
- b) a fatty acid ester of glycerol
- c) a polysaccharide
- d) an addition polymer


19. A peptide bond (also called an amide bond) joins two amino acids together. What atoms are linked by this bond?

- a) C—O
- b) C—H
- c) C—N
- d) N—S



20. An amino acid is a compound that contains at least

- a) one amino group and one amide group.
- b) two amino groups and one carboxylic acid group.
- c) one hydroxyl group and one methyl group.
- d) one carboxylic acid group and one amino group







21. The functional group found in proteins is called a (an)

رابطة ببتيدية (amide)

- a) amide.
- b) carboxylic acid.
- c) amine.
- d) amino acid.

22. Which one of these choices is the general structural formula of an amino acid?



Test bank chapter (1)

Choose the correct answer

- The SI unit of time is the
 - hour
 - second**
 - minute
 - ampere
- The diameter of an atom is approximately 1×10^{-7} mm. What is this diameter when expressed in nanometers?
 - 1×10^{-18} nm
 - 1×10^{-15} nm
 - 1×10^{-9} nm
 - 1×10^{-1} nm**
- 6.0 km is how many micrometers?
 - 6.0×10^6 μm
 - 1.7×10^{-7} μm
 - 6.0×10^9 μm**
 - 1.7×10^{-4} μm
- The SI prefixes *giga* and *micro* represent, respectively:
 - 10^{-9} and 10^{-6} .
 - 10^6 and 10^{-3} .
 - 10^3 and 10^{-3} .
 - 10^9 and 10^{-6} .**
- Which of these quantities represents the largest mass?
 - 2.0×10^2 mg
 - 0.0010 kg
 - 1.0×10^5 μg
 - 2.0×10^2 cg**
- How many cubic centimeters are there in exactly one cubic meter?
 - 1×10^{-6} cm^3
 - 1×10^{-3} cm^3
 - 1×10^{-2} cm^3
 - 1×10^6 cm^3**

7. Ammonia boils at -33.4°C . What temperature is this in $^{\circ}\text{F}$?

- a) -60.1°F
- b) -92.1°F
- c) -28.1°F
- d) $+13.5^{\circ}\text{F}$

8. Which of the following is not an SI base unit?

- a) **Kilometer**
- b) Kilogram
- c) Second
- d) Kelvin

9. Which of the following SI base units is not commonly used in chemistry?

- a) kilogram
- b) kelvin
- c) **candela**
- d) mole

10. Which of the following prefixes means $1/1000$?

- a) kilo
- b) deci
- c) centi
- d) **milli**

11. Which of the following prefixes means 1000?

- a) **kilo**
- b) deci
- c) centi
- d) milli

12. Convert -77°F to kalvin ?

- a) **212.6 K**
- b) -212.6 K
- c) -28.1 K
- d) $+13.5\text{ K}$

13. The number 0.0005678 expressed in scientific notation is:

- a) 5.678×10^4
- b) 5.67×10^{-7}
- c) 5.678×10^{-4}
- d) 5.678×10^{-3}

Explanation: Since this number is less than one start moving the decimal point to the right until there is ONE non-zero number to the left of the decimal point. Write the rest of the number as is. Write the exponent as the number of places the decimal point was moved.

14. Which of the following is the smallest distance?

- a) 21 m
- b) 2.1×10^2 cm
- c) 21 mm
- d) 2.1×10^4 pm

Explanation: Even though 2.1×10^4 is the largest number in this question, the units of pm (picometers) are the smallest units here, making it the smallest distance.

15. What temperature is 95 °F when converted to degrees Celsius?

- a) 63 °C
- b) **35 °C**
- c) 127 °C
- d) 15 °C

16. What temperature is 37 °C when converted to kelvin?

- a) **310.15**
- b) 99 k
- c) 236 k
- d) 67.15

17. What temperature is 77 K when converted to degrees Celsius?

- a) -296°C
- b) 105°C
- c) **-196°C**
- d) 25°C

18. Express 75 Tg as pg

- a) 0.75 pg
- b) **75×10^{24} pg**
- c) 0.75 pg
- d) 75×10^{-24} pg

19. The SI prefixes *Tera* and *nano* represent, respectively:

- a) 10^6 and 10^{-9}
- b) 10^3 and 10^{-6}
- c) 10^{12} and 10^{-3}
- d) 10^9 and 10^{-12}

20. Which of these quantities represents the smallest mass?

- a) 2.0×10^2 mg
- b) 0.0010 kg
- c) 1×10^5 μ g
- d) 2.0×10^2 cg

21. Express 7.5 ng as Tg

- a) 7.5×10^{-21} Tg
- b) 75×10^{24} Tg
- c) 0.75 Tg
- d) 7.5×10^{21} Tg

22. At what temperature does the numerical reading on a Fahrenheit thermometer equal that on a Celsius thermometer?

- a) 0 °F
- b) -40°F
- c) 100 °F
- d) -32 °F

Explanation: since the temperature reading is the same so that mean °F = °C

$$? F = [^{\circ}C \times 9/5] + 32 \text{ } ^{\circ}F$$

Let temperature = t

$$t = [t \times 9/5] + 32 \text{ } ^{\circ}F$$

$$t - 9/5 t = 32 \text{ } ^{\circ}F$$

$$-4/5 t = 32 \text{ } ^{\circ}F$$

$$t = - 40 \text{ } ^{\circ}F = -40 \text{ } ^{\circ}C$$

Test bank chapter (2)

Choose the correct answer

NOTE: A periodic table is required to work many of the problems in this chapter.

- Which of these elements is most likely to be a good conductor of electricity?
 - N
 - S
 - He
 - Fe**
- An atom of the isotope sulfur-31 consists of how many protons, neutrons, and electrons?
(p = proton, n = neutron, e = electron)
 - 15 p, 16 n, 15 e
 - 16 p, 15 n, 16 e**
 - 16 p, 31 n, 16 e
 - 32 p, 31 n, 32 e
- A magnesium ion, Mg^{2+} , has
 - 12 protons and 13 electrons.
 - 24 protons and 26 electrons.
 - 12 protons and 10 electrons.**
 - 24 protons and 22 electrons.
- Which of these pairs of elements would be most likely to form an ionic compound?
 - P and Br
 - Cu and K
 - C and O
 - O and Zn**
- The elements in a column of the periodic table are known as
 - metalloids.
 - a period.
 - noble gases.
 - a group.**

6. Which is the correct formula for copper (II) phosphate?
- a) Cu_2PO_4
 - b) $\text{Cu}_3(\text{PO}_4)_2$
 - c) Cu_2PO_3
 - d) $\text{Cu}(\text{PO}_4)_2$
7. The correct name for NH_4NO_3 is
- a) ammonium nitrate.
 - b) ammonium nitrogen trioxide.
 - c) ammonia nitrogen oxide.
 - d) hydrogen nitrogen oxide.
8. What is the formula for the ionic compound formed by calcium ions and nitrate ions?
- a) Ca_3N_2
 - b) $\text{Ca}(\text{NO}_3)_2$
 - c) Ca_2NO_3
 - d) Ca_2NO_2
9. The Stock system name for Mn_2O_7 is
- a) dimanganese heptaoxide.
 - b) magnesium oxide.
 - c) manganese(VII) oxide.
 - d) manganese(II) oxide.
10. Which of these elements is chemically similar to oxygen?
- a) sulfur
 - b) calcium
 - c) iron
 - d) nickel
11. The formula of stannic oxide is SnO_2 . The valence of Sn is:
- a) +1
 - b) +2
 - c) +3
 - d) +4

Explanation: To know the charge on Sn atom, make this simple calculation remember that the charge on oxygen atom is -2, let X is the charge on Sn atom

$X + (-2 \text{ (charge on O)} \times 2 \text{ (number of O atoms)}) = 0$ (equal zero because the compound is neutral)

$X - 4 = 0 \gg \gg \gg \gg \gg x = + 4$

12. Which pair of atoms constitutes a pair of isotopes of the same element?

- | | | |
|------|------------------------|---------------------|
| (a). | ${}^1_6\text{X}$ | ${}^1_7\text{X}$ |
| (b). | ${}^{14}_6\text{X}$ | ${}^{12}_6\text{X}$ |
| (c). | ${}^{17}_9\text{X}$ | ${}^{17}_8\text{X}$ |
| (d). | ${}^{19}_{10}\text{X}$ | ${}^{19}_9\text{X}$ |

Explanation: Isotopes of an element are atoms of the same element with same number of protons but different number of neutrons. Only choice (b) has 2 atoms of X with 6 protons and 8 and 6 neutrons respectively.

13. Elements in Group 8A are known as the_____.

- a) chalcogens
- b) alkali metals
- c) noble gases**
- d) alkaline earth metals

14. _____typically forms ions with a +2 charge.

- a) Transition metals
- b) Halogens
- c) Alkaline earth metals**
- d) Alkali metals

Explanation: The alkaline earth metals are in group 2A of the periodic table and lose 2 electrons to form cations with 2 positive charges.

15. An *anion* is defined as

- a) a charged atom or group of atoms with a net negative charge.**
- b) a stable atom.
- c) a group of stable atoms.
- d) an atom or group of atoms with a net positive charge.

16. A cation is defined as

- a) a charged atom or group of atoms with a net negative charge.
- b) a stable atom.
- c) a group of stable atoms.
- d) an atom or group of atoms with a net positive charge.**

17. Atoms of the same element with different mass numbers (or number of neutrons) are called

- a) ions.
- b) neutrons.
- c) chemical families.
- d) isotopes.**

18. How many neutrons are there in an atom of lead ${}_{82}\text{Pb}$ whose mass number is 208?

- a) 82
- b) 126**
- c) 208
- d) 290

19. An atom of the isotope ${}^{16}\text{S}$ -31 consists of how many protons, neutrons, and electrons?

- a) 15 p, 16 n, 15 e
- b) 16 p, 15 n, 16 e**
- c) 16 p, 31 n, 16 e
- d) 32 p, 31 n, 32 e

20. A magnesium ion, ${}_{20}\text{Ca}^{2+}$, has

- a) 20 protons and 22 electrons.
- b) 20 protons and 20 electrons.
- c) 20 protons and 18 electrons.**
- d) 22 protons and 20 electrons.

21. A sulfide ion, ${}_{16}\text{S}^{2-}$, has:

- a) 16 protons and 16 electrons
- b) 32 protons and 16 electrons
- c) 16 protons and 14 electrons
- d) 16 protons and 18 electrons**

22. Which of these pairs of elements would be most likely to form a molecular compound?

- a) Na and Br
- b) Ca and O
- c) C and O**
- d) Zn and O

23. What is the formula for the ionic compound formed by calcium ions and nitrate ions?

- a) Ca_3N_2
- b) $\text{Ca}(\text{NO}_3)_2$**
- c) Ca_2NO_3
- d) Ca_2NO_2

24. Which is the correct formula for copper(II) phosphate?

- a) Cu_2PO_4
- b) $\text{Cu}_3(\text{PO}_4)_2$**
- c) Cu PO_4
- d) $\text{Cu}(\text{PO}_4)_2$

25. The correct name for NH_4NO_3 is

- a) ammonium nitrate.**
- b) ammonium nitrogen trioxide.
- c) ammonia nitrogen oxide.
- d) hydrogen nitrogen oxide.

26. The correct name for PCl_5 is

- a) monophosphate pentachloride
- b) phosphorus chloride
- c) monophosphate tetrachloride
- d) Phosphorus pentachloride**

27. Which of the following expressions represents two molecules of water?

- a) H_2O
- b) H_2O_2
- c) $2 \text{H}_2\text{O}$**
- d) 2HO_2

28. The empirical formula of a compound with molecules containing 12 carbon atoms, 14 hydrogen atoms, and 6 oxygen atoms is _____.

- a) $\text{C}_{12}\text{H}_{14}\text{O}_6$
- b) $\text{C}_2\text{H}_4\text{O}$
- c) CH_2O
- d) $\text{C}_6\text{H}_7\text{O}_3$**

Explanation: The empirical formula is always the simplest possible whole number ratio between the atoms of the molecules.

29. The charge on the manganese in the salt MnF_3 is _____.

- a) +1
- a) -1
- c) +3**
- d) -2

Explanation: Since every F has one negative charge, the Mn can have only 3 positive charges.

30. Magnesium reacts with a certain element to form a compound with the general formula MgX . What would the most likely formula be for the compound formed between potassium and element X?

- a) KX
- b) K_2X_2
- c) K_2X_3
- d) **None of the above**

Explanation: In the compound MgX , X must have 2 negative charges since Mg will always have 2 positive charges. The element K will always form an ion with 1 positive charge and hence the only combination of K and X could be K_2X , which is not one of the options.

31. Barium forms an ion with a charge of_____.

- a) +1
- b) -2
- c) +3
- d) **None of the above.**

Explanation: Barium is in group 2A of the periodic table and forms ions with only 2 positive charges.

31. Aluminum forms an ion with a charge of_____.

- a) +2
- b) -3
- c) **+3**
- d) +1

32. Iodine forms an ion with a charge of_____.

- a) -7
- b) +1
- c) **-1**
- d) +2

33. The chemical symbol for the ion with 11 protons and 10 electrons.

- a) Na
- b) F^-
- c) Ne
- d) **Na^+**

34. Which of these compounds is a binary compound?

- a) **NaCl**
- b) MgSO₄
- c) NaOH
- d) HCN

35. Atoms with the same number of electrons and number of protons are called...

- a) ions
- b) isotopes
- c) **neutral atoms**
- d) different atoms

36. Atoms which have different number of electrons are called...

- a) **ions**
- b) isotopes
- c) neutral atoms
- d) different atoms

37. Use the following table and choose which of the species are positively charged?

Atom or ion element	I	II	III	IV	V	VI
Atom or ion electrons (e)	6	10	18	10	28	7
Atom or ion protons (p)	6	8	17	11	30	7
Atom or ion neutrons (n)	6	8	18	11	36	6

- A. III and V
- B. **IV and V**

- C. II and III
- D. I and VI


38. Which isotope has 45 neutrons?

- (a). ⁸⁰₃₆Kr
- (b). ⁷⁸₃₄Se
- (c). ⁸⁰₃₅**Br**
- (d). ³⁴₁₇Cl

39. In the periodic table, the elements are arranged in_____.

- a) alphabetical order
- b) **order of increasing atomic number**
- c) order of increasing metallic properties
- d) order of increasing neutron content

40. An element in the upper right corner of the periodic table is _____.
- either a metal or metalloid
 - a metal
 - a non-metal**
 - either a metalloid or a non-metal
41. An element that appears in the lower left corner of a periodic table is _____.
- either a metal or metalloid
 - a metal**
 - either a metalloid or a non-metal
 - a non-metal
42. A molecular formula always indicates _____.
- how many of each atom are in a molecule**
 - the simplest whole-number ratio of different atoms in a compound
 - which atoms are attached to which in a molecule
 - the isotope of each element in a compound
 -
43. An empirical formula always indicates _____.
- which atoms are attached to which in a molecule
 - how many of each atom are in a molecule
 - the simplest whole-number ratio of different atoms in a compound**
 - the geometry of a molecule
44. There are _____ protons, _____ neutrons, and _____ electrons in $^{131}\text{I}^-$.
- 131, 53, and 54
 - 131, 53 and 52
 - 53, 78, and 54**
 - 53, 131, and 52
45. Which species has 48 electrons?
- $^{118}_{50}\text{Sn}^{+2}$
 - $^{116}_{50}\text{Sn}^{+4}$
 - $^{112}_{48}\text{Cd}^{+2}$
 - $^{68}_{31}\text{Ga}$



Test bank chapter (3)

Choose the correct answer


1. What is the mass, in grams, of one copper atom?
 - a) 1.055×10^{-22} g
 - b) 63.55 g
 - c) 20 g
 - d) 1.66×10^{-24} g
2. Determine the number of moles of aluminum in 96.7 g of Al?
 - a) 0.279 mol
 - b) 3.58 mol
 - c) 7.43 mol
 - d) 4.21 mol
3. Which of the following samples contains the greatest number of atoms?
 - a) 100 g of Pb
 - b) 2.0 mole of Ar
 - c) 0.1 mole of Fe
 - d) 5 g of He
4. How many molecules are there in 0.11 g of formaldehyde (CH₂O)?
 - a) 6.1×10^{-27}
 - b) 3.7×10^{-3}
 - c) 4×10^{22}
 - d) 2.2×10^{21}
5. How many sulfur atoms are present in 25.6 g of Al₂(S₂O₃)₃?
 - a) 0.393
 - b) 6×10^{-5}
 - c) 3.95×10^{22}
 - d) 2.37×10^{23}
6. The percent composition by mass of a compound is 76.0% C, 12.8% H, and 11.2% O. The molar mass of this compound is 284.5 g/mol. What is the molecular formula of the compound?
 - a) C₁₀H₆O
 - b) C₉H₁₈O
 - c) C₁₆H₂₈O₄
 - d) C₁₈H₃₆O₂

Mass
relationships in
chemical
reactions


7. What is the coefficient of H_2O when the following equation is properly balanced with the smallest set of whole numbers?



- a) 3
b) 4
c) 6
d) 12
8. When 22.0 g NaCl and 21.0 g H_2SO_4 are mixed and react according to the equation below, which is the limiting reagent?



- a) H_2SO_4
b) Na_2SO_4
c) HCl
d) NaCl
9. What are the coefficients, when the following equation is balanced?



- a) 1, 1, 1, 1
b) 2, 3, 2, 3
c) 4, 7, 4, 6
d) 1, 3, 1, 2
10. How many moles of carbon atoms are in 4 moles of dimethyl sulfoxide ($\text{C}_2\text{H}_6\text{SO}$)?
- a) 2
b) 6
c) 8
d) 4

Explanation: This is based on reading the formula and correctly extracting information from it. The formula $\text{C}_2\text{H}_6\text{SO}$ indicates that every mole of this compound has 2 moles of carbon atoms in it. Thus 4 moles of the compound would have $4 \times 2 = 8$ moles of C atoms.

11. How many sulfur atoms are in 25 molecules of $\text{C}_4\text{H}_4\text{S}_2$?

- a) 20
b) 4.8×10^{25}
c) 3.0×10^{23}
d) 50

Explanation: The molecular formula indicates that every molecule of $\text{C}_4\text{H}_4\text{S}_2$ has 2 sulfur atoms per molecule and hence 25 molecules of this compound will have $25 \times 2 = 50$ atoms of sulfur.

12. Calculate hydrogen atoms in 25 molecules of $C_4H_4S_2$.

- a) 25
- b) 3.8×10^{24}
- c) 6.0×10^{25}
- d) 100

Explanation: The formula of $C_4H_4S_2$ indicates that there are 4 hydrogen atoms per molecule and hence 100 hydrogen atoms in 25 molecules of $C_4H_4S_2$.

13. How many grams of oxygen are in 65.0 g of $C_2H_2O_2$?

- a) 18
- b) 29
- c) 9.5
- d) 35.8

Explanation: This question uses the mole to mole ratio between oxygen and $C_2H_2O_2$ and needs the following steps.

$$\frac{65.0 \text{ g } C_2H_2O_2}{58.0 \text{ g} \cdot \text{mol}^{-1}} \times \frac{2 \text{ moles O}}{1 \text{ mole } C_2H_2O_2} \times \frac{15.99 \text{ g O}}{1 \text{ mole of O}} = 35.8 \text{ g of O}$$

14. How many moles of carbon dioxide are there in 52.06 g of carbon dioxide?

- a) 0.8452
- b) 1.183
- c) 1.183×10^{23}
- d) 8.648×10^2

Explanation: This is a straight-forward conversion from grams to moles of CO_2 which is done as follows:

$$52.06 \text{ g } CO_2 \times \frac{1 \text{ mole } CO_2}{43.99 \text{ g } CO_2} = 1.183 \text{ moles of } CO_2$$

15. How many moles of magnesium nitrate, $Mg(NO_3)_2$, are in a 2.35 g of this compound?

- a) 38.4
- b) 65.8
- c) 0.0158
- d) 0.0261

Explanation: This is a straight-forward conversion from grams to moles of $Mg(NO_3)_2$ which is done as follows:

$$2.35 \text{ g } Mg(NO_3)_2 \times \frac{1 \text{ mole } Mg(NO_3)_2}{148.3148 \text{ g}} = 0.0158 \text{ moles}$$


16. How many moles of ammonium ions are there in 25.5 g of ammonium carbonate?

- a) 0.468
- b) 0.288
- c) 0.531
- d) 2.00

Explanation: Realize that the formula for ammonium carbonate is $(\text{NH}_4)_2\text{CO}_3$ and calculate the molar mass (96.0856 g/mol). Convert grams to moles and then using the stoichiometric ratio find the # of moles of ammonium ions.

$$25.5 \text{ g } (\text{NH}_4)_2\text{CO}_3 \times \frac{1 \text{ mol } (\text{NH}_4)_2\text{CO}_3}{96.0856 \text{ g}} \times \frac{2 \text{ moles NH}_4^+}{1 \text{ mol } (\text{NH}_4)_2\text{CO}_3} = 0.531 \text{ moles}$$

17. Magnesium and nitrogen react in a combination reaction to produce magnesium nitride:



In a particular experiment, 5.47 g sample of N_2 reacts completely. How many grams of Mg are needed for this reaction?

- a) 14.2 g
- b) 24.1 g
- c) 16.1 g
- d) 0.92 g

Explanation: Ensure that the equation is balanced. The grams of N_2 must be converted to moles of N_2 and then using the stoichiometric ratio between the Mg and N_2 , the grams of Mg can be calculated.

$$5.47 \text{ g N}_2 \times \frac{1 \text{ mole N}_2}{28.0134 \text{ g}} \times \frac{3 \text{ mole Mg}}{1 \text{ mole N}_2} \times \frac{24.3050 \text{ g Mg}}{1 \text{ mole Mg}} = 14.2 \text{ g Mg}$$

18. What information would you need to calculate the average atomic mass of an element?

- a) The number of neutrons in the element.
- b) The atomic number of the element.
- c) The mass and abundance of each isotope of the element.
- d) The position in the periodic table of the element.

19. The atomic masses of ^{35}Cl (75.53 %) and ^{37}Cl (24.47 %) are 34.968 amu and 36.956 amu, respectively. Calculate the average atomic mass of chlorine.

- a) 35.96 amu
- b) 35.45 amu
- c) 36.47 amu
- d) 71.92 amu

Mass relationships in chemical reactions

20. How many atoms are there in 5.10 moles of sulfur (${}_{16}\text{S} = 32 \text{ amu}$)?

- a) 3.07×10^{24}
- b) 9.59×10^{22}
- c) 6.02×10^{23}
- d) 9.82×10^{25}

21. Iodine has two isotopes ${}^{126}\text{I}$ and ${}^{127}\text{I}$, with the equal abundance. Calculate the average atomic mass of Iodine (${}_{53}\text{I}$).

- a) 126.5 amu
- b) 35.45 amu
- c) 1.265 amu
- d) 71.92 amu


22. The atomic masses of ${}^6\text{Li}$ and ${}^7\text{Li}$ are 6.0151 amu and 7.0160 amu, respectively. Calculate the natural abundance of these two isotopes. The average atomic mass of Lithium ($\text{Li} = 6.941 \text{ amu}$).

- a) ${}^6\text{Li} = 7.49\%$, ${}^7\text{Li} = 92.51\%$
- b) ${}^7\text{Li} = 7.49\%$, ${}^6\text{Li} = 92.51\%$
- c) ${}^6\text{Li} = 8.49\%$, ${}^7\text{Li} = 95.51\%$
- d) ${}^7\text{Li} = 7.22\%$, ${}^6\text{Li} = 82.51\%$

23. How many atoms are present in 3.14 g of copper (Cu)?


- a) 2.98×10^{22}
- b) 1.92×10^{23}
- c) 1.89×10^{24}
- d) 6.02×10^{23}

24. How many moles of NO_2 can be produced by the reaction of 0.886 mole of NO with 0.503 mole of O_2 according to the following chemical equation? (Note: First determine which is the limiting reagent).



- a) 0.886 mol
- b) 0.503 mol
- c) 1.01 mol
- d) 1.77 mol


25. How many kilograms of NH_3 are needed to produce $1.00 \times 10^5 \text{ kg}$ of $(\text{NH}_4)_2\text{SO}_4$ according to the following chemical equation?



- a) $1.70 \times 10^4 \text{ kg}$
- b) $3.22 \times 10^3 \text{ kg}$
- c) $2.58 \times 10^4 \text{ kg}$
- d) $7.42 \times 10^4 \text{ kg}$

**Mass
relationships in
chemical
reactions**

26. When 3.60 moles of CO mixed with excess oxygen gas and CO₂ is formed. Calculate no. of moles of CO₂ produced.




- a) 7.20 mol
- b) 44.0 mol
- c) 3.60 mol
- d) 1.80 mol

27. How many grams of N₂O are formed if 0.46 mole of NH₄NO₃ is used in the following chemical reaction?




- a) 2.0 g
- b) 3.7×10^1 g
- c) 2.0×10^1 g
- d) 4.6×10^{-1} g

28. What is the theoretical yield of chromium that can be produced by the reaction of 40.0 g of Cr₂O₃ with 8.00 g of aluminum according to the chemical equation below?



- a) 7.7 g
- b) 15.4 g
- c) 27.3 g
- d) 30.8 g


29. What is the percent yield of HF that can be produced by the reaction of 6.00 kg of CaF₂ with an excess of H₂SO₄ which yield 2.86 kg of HF according to the following chemical equation?



- a) 93.0 %
- b) 95.3 %
- c) 47.6 %
- d) 62.5 %

Mass
relationships in
chemical
reactions

30. Hydrochloric acid can be prepared by the following reaction:



How many grams of HCl can be prepared from 2.00 moles of H_2SO_4 and 150 g of NaCl?

- a) 7.30 g
- b) 93.5 g
- c) 146 g
- d) 150 g

31. Calculate the molar mass of Li_2CO_3 .

- a) 73.89 g
- b) 66.95 g
- c) 41.89 g
- d) 96.02 g

32. How many molecules of ethane (C_2H_6) are present in 0.334 g of C_2H_6 ?

- a) 2.01×10^{23}
- b) 6.69×10^{21}
- c) 4.96×10^{22}
- d) 8.89×10^{20}

33. Out of these, which is the richest source of nitrogen on a mass percentage basis?

- a) Urea, $(\text{NH}_2)_2\text{CO}$
- b) Ammonium nitrate, NH_4NO_3
- c) Guanidine, $\text{HNC}(\text{NH}_2)_2$
- d) Ammonia, NH_3

34. An analysis of Allicin (molar mass ≈ 162 g/mol) gives C: 44.4 percent; H: 6.21 percent; S: 39.5 percent; O: 9.86 percent. What is its molecular formula?

- a) $\text{C}_{12}\text{H}_{20}\text{S}_4\text{O}_2$
- b) $\text{C}_7\text{H}_{14}\text{SO}$
- c) $\text{C}_6\text{H}_{10}\text{S}_2\text{O}$
- d) $\text{C}_5\text{H}_{12}\text{S}_2\text{O}_2$

Mass relationships in chemical reactions

35. How many moles of Fe are present in 24.6 g of Fe_2O_3 ?
- a) 2.13 mol
 - b) 0.456 mol
 - c) 0.154 mol
 - d) **0.308 mol**
36. How many grams of sulfur (S) are needed to react completely with 246 g of mercury (Hg) to form HgS ?
- a) **39.3 g**
 - b) 24.6 g
 - c) 9.66×10^3 g
 - d) 201 g
37. What is the mass of F (fluoride) in 24.6 g of Tin (II) fluoride (SnF_2)?
- a) 18.6 g
 - b) 24.3 g
 - c) **5.97 g**
 - d) 75.7 g
38. What is the empirical formula of the compound with the following composition? 2.1 percent H, 65.3 percent O, 32.6 percent S.
- a) **H_2SO_4**
 - b) H_2SO_3
 - c) $\text{H}_2\text{S}_2\text{O}_3$
 - d) HSO_3
39. Which of the following equations is balanced?
- a) $2\text{C} + \text{O}_2 \rightarrow \text{CO}$
 - b) **$2\text{CO} + \text{O}_2 \rightarrow 2\text{CO}_2$**
 - c) $\text{H}_2 + \text{Br}_2 \rightarrow \text{HBr}$
 - d) $2\text{K} + \text{H}_2\text{O} \rightarrow 2\text{KOH} + \text{H}_2$



Test bank chapter (4)

Choose the correct answer

1. A 50.0 mL sample of 0.436 M NH_4NO_3 is diluted with water to a total volume of 250.0 mL. What is the ammonium nitrate concentration in the resulting solution?
 - a) 21.8 M
 - b) 0.459 M
 - c) 2.18×10^{-2} M
 - d) 8.72×10^{-2} M
2. How many milliliters would you need to prepare 60.0 mL of 0.200 M HNO_3 from a stock solution of 4.00 M HNO_3 ?
 - a) 3 mL
 - b) 240 mL
 - c) 24 mL
 - d) 1000 mL
3. What is the concentration (M) of CH_3OH in a solution prepared by dissolving 11.7 g of CH_3OH in sufficient water to give exactly 230 mL of solution?
 - a) 11.7
 - b) 2.30×10^{-2}
 - c) 0.0841
 - d) 1.59

Explanation: Need to convert the grams of CH_3OH to moles and then find the molarity of the solution by using the molarity formula. Do not forget to convert the ml to L.

$$11.7 \text{ g CH}_3\text{OH} \times \frac{1 \text{ mole CH}_3\text{OH}}{32.042 \text{ g}} \times \frac{1}{0.230 \text{ L}} = 1.59 \text{ M}$$

4. How many grams of H_3PO_4 are in 35.1 mL of a 2.75 M solution of H_3PO_4 ?
 - a) 0.61
 - b) 9.46
 - c) 20
 - d) 4.9

Explanation: Need to convert the ml of H_3PO_4 to liters and then find the # of moles of phosphoric acid. The moles of phosphoric acid can then be converted to grams of phosphoric acid.

$$35.1 \text{ ml} \times \frac{1 \text{ L}}{1000 \text{ ml}} \times 2.75 \text{ M} \times \frac{97.99 \text{ g H}_3\text{PO}_4}{1 \text{ mole}} = 9.46 \text{ g H}_3\text{PO}_4$$

5. What is the concentration (M) of a Na_2SO_4 solution prepared by dissolving 5.35 g of Na_2SO_4 in sufficient water to give 330 mL of solution?

- a) 1.14×10^2
- b) 0.016
- c) 61.7
- d) **0.114**

Explanation: Convert grams of Na_2SO_4 to moles of Na_2SO_4 , ml of water to liters of water and then find the molarity of the solution by using the molarity formula.

$$5.35 \text{ g Na}_2\text{SO}_4 \times \frac{1 \text{ mole Na}_2\text{SO}_4}{142.035 \text{ g Na}_2\text{SO}_4} \times \frac{1}{0.330 \text{ L}} = 0.114 \text{ M Na}_2\text{SO}_4$$

6. How many grams of LiOH are there in 750.0 mL of a 0.0158 M LiOH

solution?

- a) 2.11×10^{-5}
- b) 11.3
- c) **0.284**
- d) 3.50

Explanation: Calculate the number of moles of LiOH present in this solution using the molarity formula and then convert the number of moles to grams of LiOH .

$$7.50 \times 10^{-1} \text{ L} \times 0.0158 \text{ M} \times \frac{23.948 \text{ g}}{1 \text{ mole LiOH}} = 0.284 \text{ g LiOH}$$

7. A 50.0 mL sample of 0.436 M NH_4NO_3 is diluted with water to a total volume of 250.0 mL. What is the ammonium nitrate concentration in the resulting

solution?

- a) 21.8 M
- b) 0.459 M
- c) 2.18×10^{-2} M
- d) **8.72×10^{-2} M**

8. A 3.682 g sample of potassium chlorate KClO_3 is dissolved in enough water to give 375 mL of solution. What is the chlorate ion concentration in this solution?

- a) 3.00 M
- b) 4.41×10^{-2} M
- c) 0.118 M
- d) **8.01×10^{-2} M**

Gases

Test bank chapter (5)

Choose the most correct answer

1. A sample of oxygen occupies 47.2 liters under a pressure of 1240 torr at 25°C. What volume would it occupy at 25°C if the pressure were decreased to 730 torr?

- a) 27.8 L
- b) 29.3 L
- c) 32.3 L
- d) **80.2 L**

2. Under conditions of fixed temperature and amount of gas, Boyle's law requires that

- I. $P_1V_1 = P_2V_2$
- II. $PV = \text{constant}$
- III. $P_1/P_2 = V_2/V_1$

- a) I only
- b) II only
- c) III only
- d) **I, II, and III**

3. The volume of a sample of nitrogen is 6.00 liters at 35°C and 740 torr. What volume will it occupy at STP?

- a) 6.59 L
- b) 5.46 L
- c) 6.95 L
- d) **5.18 L**

4. The density of chlorine gas at STP, in grams per liter, is approximately:

- a) 6.2
- b) **3.2**
- c) 3.9
- d) 4.5

Explanation: $d = \text{molar mass} \times p / RT = 70 \times 1 / 0.082 \times 273 = 3.17 \text{ g/L}$

5. What pressure (in atm) would be exerted by 76 g of fluorine gas in a 1.50 liter vessel at -37°C?

- a) **26 atm**
- b) 4.1 atm
- c) 19,600 atm
- d) 84 atm

6. What is the density of ammonia gas at 2.00 atm pressure and a temperature of 25.0°C?

- a) 0.720 g/L
- b) 0.980 g/L
- c) **1.39 g/L**
- d) 16.6 g/L

7. Convert 2.0 atm to mmHg

- a) 150 mmHg
- b) 0.27 mmHg
- c) 150 mmHg
- d) **1520 mmHg**

Gases

8. A container with volume 71.9 mL contains water vapor at a pressure of 10.4 atm and a temperature of 465°C. How many grams of the gas are in the container?

- a) 0.421 g
- b) **0.222 g**
- c) 0.183 g
- d) 0.129 g

Explanation: $n = PV/RT = 0.0719 \times 10.4 / (0.0821 \times (465 + 273)) = 0.012$ mole

$$\text{Mass} = n \times \text{molar mass} = 0.012 \times 18 = 0.222 \text{ g}$$

9. What is the molar mass of a pure gaseous compound having a density of 4.95 g/L at -35 °C and 1020 torr?

- a) 24 g/mole
- b) 11 g/mole
- c) **72 g/mole**
- d) 120 g/mole

10. A 0.580 g sample of a compound containing only carbon and hydrogen contains 0.480 g of carbon and 0.100 g of hydrogen. At STP, 33.6 mL of the gas has a mass of 0.087 g. What is the molecular (true) formula for the compound?

- a) CH₃
- b) C₂H₆
- c) C₂H₅
- d) **C₄H₁₀**

11. Gas occupy 6L at 37°C what will be its volume when its temperature is doubled?

- a) **12 L**
- b) 6L
- c) 3.2 L
- d) 2L

12. A mixture of 90.0 grams of CH₄ and 10.0 grams of argon has a pressure of 250 torr under conditions of constant temperature and volume. The partial pressure of CH₄ in torr is:

- (a) 143
- (b) 100
- (c) 10.7
- (d) **239**

Explanation: from Dalton law $\gg \gg P_{\text{CH}_4} = X_{\text{CH}_4} P_{\text{total}}$, $n_{\text{CH}_4} = 90 / 16 = 5.625$ mole, $n_{\text{Ar}} = 10 / 39.95 =$

0.250 mole $X_{\text{CH}_4} = n_{\text{CH}_4} / (n_{\text{CH}_4} + n_{\text{Ar}}) = 5.625 / (5.625 + 0.250) = 0.96 \gg \gg \gg P_{\text{CH}_4} = 0.96 \times 250 = 239.3$ torr

13. What pressure (in atm) would be exerted by a mixture of 1.4 g of nitrogen gas and 4.8 g of oxygen gas in a 200 mL container at 57°C?

- a) 4.7
- b) 34
- c) 47
- d) **27**

Explanation: $P = n_{\text{total}} RT/V$, $n_{\text{N}_2} = 1.4 / (2 \times 14) = 0.05$ mole, $n_{\text{O}_2} = 4.8 / (2 \times 16) = 0.15$ mole
 $P = (0.05 + 0.15) (0.0821 \times (57 + 273)) / 0.2 = 27$ atm

Gases

14. A sample of hydrogen gas collected by displacement of water occupied 30.0 mL at 24°C and pressure 736 torr. What volume would the hydrogen occupy if it were dry and at STP? The vapor pressure of water at 24.0°C is 22.4 torr.


- a) 32.4 mL
- b) 21.6 mL
- c) 36.8 mL
- d) 25.9 mL**

Explanation: from Dalton law $\gg \gg P_{H_2} = P_{total} - P_{H_2O}$, $P_{H_2} = 736 - 22.4 = 713.6$ torr

$$n = PV/RT \gg \gg \gg n = (713.6/760) \times 0.03 / 0.0821 \times (24+273) = 0.00115 \text{ mole}$$

$$\text{at STP} \gg \gg \gg V = nRT/P = 0.00115 \times 0.0821 \times 273 / 1 = 0.026 \text{ L} \times 1000 = 25.9 \text{ mL}$$

15. Ammonia burns in oxygen gas to form nitric oxide (NO) and water vapor. How many volumes of NO are obtained from one volume of ammonia at the same temperature and pressure?



- a) One**
- b) Two
- c) Three
- d) Four

16. The pressure of 6.0 L of an ideal gas in a flexible container is decreased to one-third of its original value, and its absolute temperature is decreased by one-half. What is the final volume of the gas?

- a) 9.0 L**
- b) 6.0 L
- c) 4.0 L
- d) 1 L

Explanation: let $V_1 = 6$ & $V_2 = ?$, $T_1 = T$ & $T_2 = \frac{1}{2} T$, $P_1 = P$ & $P_2 = \frac{1}{3} P$

$$\text{From combined gas law } P_1 V_1 / T_1 = P_2 V_2 / T_2 \gg \gg \gg \frac{P \times 6}{T} = \frac{(\frac{1}{3})P \times V_2}{(\frac{1}{2})T} \gg \gg V_2 = \frac{P \times 6 \times T \times 3}{T \times 2 \times P}$$

17. Gas A is at 30°C and gas B is at 20°C. Both gases are at 1 atmosphere. What is the ratio of the volume of 1 mole gas A to 1 mole of gas B

- a) 606:303
- b) 3 : 2
- c) 2 : 3
- d) 303 : 293**


$$\text{Explanation: } \frac{V_A}{T_A} = \frac{V_B}{T_B} \gg \gg \gg \frac{V_A}{30+273} = \frac{V_B}{20+273} \gg \gg \gg \frac{V_A}{303} = \frac{V_B}{293}$$

18. The sample of argon occupies 50L at standard temperature. Assuming constant pressure, what volume with the gas occupy if the temperature is doubled.

- a) 25L
- b) 50L
- c) 100L**
- d) 100 mL

Gases

19. What total gas volume (in liters) at 520°C and 880 torr would result from the decomposition of 33 g of potassium bicarbonate according to the equation:



- (a) 56 L
- (b) 37 L
- (c) 10 L
- (d) 19 L**

20. Calculate the weight of KClO_3 that would be required to produce 29.5 L of oxygen measured at 127°C and 760 torr.



- (a) 73.5 g**
- (b) 12.2 g
- (c) 14.6 g
- (d) 24.4 g

21. The ideal gas law predicts that the molar volume (volume of one mole) of gas equals:

- (a) mRT/PV
- (b) $(MM)P/RT$
- (c) $1/2ms^{-2}$
- (d) RT/P**

22. For a gas, which pair of variables are inversely proportional to each other (if all other conditions remain constant)?

- a) P, V**
- b) V, T
- c) n, V
- d) n, P

23. Convert 562 mmHg to atm

- a) 0.739 atm**
- b) 4.27×10^5 atm
- c) 1.05 atm
- d) 0.562 atm

24. What is the volume of one mole of an ideal gas at STP?

- a) 24.5 L
- b) 22.4 L**
- c) 1.0 L
- d) 10.0 L

25. What are standard temperature and pressure (STP)?

- a) 0 °C, 1 torr
- b) 25 °C, 1 torr
- c) 0 °C, 1 atm**
- d) 25 °C, 1 atm

26. What is the unit of mole fraction

- a) mol
- b) mol^{-1}
- c) unitless**
- d) mol^2

Gases

27. Refer to Dalton's law of partial pressures explain what mole fraction is
- The number of moles of one component
 - The ratio of the number of moles of one component to the number of moles of all components present.**
 - The number of moles of one component divided by 100
 - The ratio of the number of moles of all components present to the number of moles of one component.
28. Write the ideal gas equation. Give the units for each term in the equation
- $PV = nRT$; P in torr, V in L, n in mol, R in Latm/Kmol, T in °C.
 - $PV = nRT$; P in torr, V in L, n in mol, R in Latm/Kmol, T in K.
 - (c) $PV = nRT$; P in atm, V in L, n in mol, R in Latm/Kmol, T in K.**
 - $PV = nRT$; P in atm, V in L, n in mol, R in Latm/Kmol, T in °C .
29. What is the difference between a gas and a vapor?
- A gas is a substance normally in the gaseous state at normal atmospheric conditions (25C, 1 atm); a vapor is the gaseous form of any substance that is a liquid or a solid at normal temperatures and pressures.**
 - A gas is the gaseous form of any substance; a vapor refers to a gas over a water surface.
 - A gas is a substance normally in the gaseous state at normal atmospheric conditions (25C, 1 atm); a vapor is a gas over a water surface.
 - A gas and a vapor are two interchangeable nomenclatures; they are identical.
30. What volume is occupied by 19.6 g of methane (CH₄) at 27°C and 1.59 atm?
- 1.71 L
 - 18.9 L**
 - 27.7 L
 - 302 L
31. A 4.37 gram sample of a certain diatomic gas occupies a volume of 3.00 L at 1.00 atm and a temperature of 45°C. Identify this gas.
- F₂**
 - N₂
 - H₂
 - O₂

Explanation: $MM = mRT/PV \gggg MM = 4.37 \times 0.0821 \times (45+273) / 1 \times 3 = 37.77/2 = 18.88 \text{ g/mole} \sim \text{F}_2$


32. A sample of hydrogen gas was collected over water at 21°C and 685 mmHg. The volume of the container was 7.80 L. Calculate the mass of H₂(g) collected. (Vapor pressure of water = 18.6 mmHg at 21°C.)
- 0.283 g
 - 0.571 g**
 - 0.589 g
 - 7.14 g
33. Which of the following is/are characteristic(s) of gases?
- High compressibility
 - Relatively large distances between molecules
 - Formation of homogeneous mixtures regardless of the nature of gases
 - High compressibility, relatively large distances between molecules AND formation of homogeneous mixtures regardless of the nature of gases**

Gases

34. A small bubble rises from the bottom of a lake, where the temperature and pressure are 4°C and 3.0 atm , to the water's surface, where the temperature is 25°C and the pressure is 0.95 atm . Calculate the final volume of the bubble if its initial volume was 2.1 mL .
- 0.72 mL
 - 6.2 mL
 - 41.4 mL
 - 7.1 mL**
35. Calculate the mass, in grams, of 2.74 L of CO gas measured at 33°C and 945 mmHg .
- 0.263 g
 - 2.46 g
 - 3.80 g**
 - 35.2 g
36. Which of the following gases will have the greatest density at the same specified temperature and pressure?
- H_2
 - CClF_3**
 - CO_2
 - C_2H_6
37. Determine the molar mass of chloroform gas if a sample weighing 0.389 g is collected in a flask with a volume of 102 cm^3 at 97°C . The pressure of the chloroform is 728 mmHg .
- 187 g/mol
 - 121 g/mol**
 - 112 g/mol
 - 31.6 g/mol
38. What is the molar mass of Freon-11 gas if its density is 6.13 g/L at STP?
- 0.274 g/mol
 - 3.64 g/mol
 - 78.2 g/mol
 - 137 g/mol**
40. A mixture of three gases has a total pressure of $1,380\text{ mmHg}$ at 298 K . The mixture is analyzed and is found to contain 1.27 mol CO_2 , 3.04 mol CO , and 1.50 mol Ar . What is the partial pressure of Ar ?
- 0.258 atm
 - 301 mmHg
 - 356 mmHg**
 - $5,345\text{ mmHg}$

Gases

41. A sample of hydrogen gas was collected over water at 21°C and 685 mmHg. The volume of the container was 7.80 L. Calculate the mass of H₂(g) collected. (Vapor pressure of water = 18.6 mmHg at 21°C.)
- a) 0.283 g
 - b) 0.572 g
 - c) 0.589 g
 - d) 7.14 g
42. A 0.271 g sample of an unknown vapor occupies 294 mL at 140°C and 847 mmHg. The empirical formula of the compound is CH₂. What is the molecular formula of the compound?
- a) CH₂
 - b) C₂H₄
 - c) C₃H₆
 - d) C₄H₈
43. How many liters of chlorine gas at 25°C and 0.950 atm can be produced by the reaction of 12.0 g of MnO₂? $\text{MnO}_2(\text{s}) + 4\text{HCl}(\text{aq}) \rightarrow \text{MnCl}_2(\text{aq}) + 2\text{H}_2\text{O}(\text{l}) + \text{Cl}_2(\text{g})$
- a) 5.36×10^{-3} L
 - b) 0.138 L
 - c) 0.282 L
 - d) 3.55 L




Test bank chapter (7)

Choose the most correct answer

- The lowest energy state of an atom is referred to as its
 - bottom state.
 - ground state.**
 - fundamental state.
 - original state.
- All s orbitals are
 - shaped like four-leaf clovers.
 - dumbbell-shaped.
 - spherical.**
 - triangular.
- $[\text{He}]2s^22p^2$ is the electron configuration of which element?
 - Beryllium Be
 - Boron B
 - carbon C**
 - nitrogen N
- What are the valence electrons of vanadium (V)?
 - $4s^2$
 - $3d^3$
 - $4s^23d^3$**
 - $3d^5$
- What are the valence electrons of gallium Ga?
 - $4s^2$
 - $3d^3$
 - $4s^24p^1$**
 - $3d^5$
- The electron configuration of a neutral atom is $[\text{Ne}] 3s^23p^1$. The four quantum numbers of the last electron are:
 - (2, 1, -1, +1/2)
 - (3, 3, -1, +1/2)
 - (3, 0, -1, +1/2)
 - (3, 1, -1, +1/2)**
- How many unpaired electrons does chromium (Cr) have?
 - 0
 - 2
 - 4
 - 6**
- How many unpaired electrons does selenium (Se) have?
 - 0
 - 2**
 - 4
 - 6

9. What is the maximum number of orbitals described by the quantum numbers: $n = 3$ $l = 2$
- 1
 - 3
 - 5**
 - 9
10. What is the maximum number of orbitals described by the quantum numbers: $n = 4$
- 7
 - 14
 - 16**
 - 48
11. The maximum number of electrons that can occupy an energy level described by the principal quantum number, n , is
- $n + 1$
 - $2n$
 - $2n^2$**
 - n^2
12. A possible set of quantum numbers for the last electron added to complete an atom of sodium Na in its ground state is
- $n = 3, l = 1, m_l = 0, m_s = \frac{1}{2}$
 - $n = 3, l = 0, m_l = 0, m_s = \frac{1}{2}$**
 - $n = 2, l = 1, m_l = -1, m_s = \frac{1}{2}$
 - $n = 2, l = 0, m_l = -1, m_s = \frac{1}{2}$
13. The ground-state electron configuration of a calcium Ca atom is
- [Ne]3s²
 - [Ne]3s²3p⁶
 - [Ar]4s¹3d¹
 - [Ar]4s²**
14. Which one of the following sets of quantum numbers is not possible?
- | | n | l | m_l | m_s |
|-------|-----|-----|-------|-------|
| Row 1 | 4 | 3 | -2 | +1/2 |
| Row 2 | 3 | 2 | -3 | -1/2 |
| Row 3 | 3 | 0 | 0 | +1/2 |
| Row 4 | 4 | 1 | 1 | -1/2 |
| Row 5 | 2 | 0 | 0 | +1/2 |
- Row 1
 - Row 2**
 - Row 3
 - Row 4
15. The number of orbitals in a d subshell is
- 1
 - 3
 - 5**
 - 7

16. Which ground-state atom has an electron configuration described by the following *orbital diagram*?



- a) phosphorus
- b) germanium
- c) selenium
- d) tellurium

17. A ground-state atom of nickel has ____ unpaired electrons and is ____.

- a) 0, diamagnetic
- b) 6, diamagnetic
- c) 3, paramagnetic
- d) 2, paramagnetic

18. What is the frequency (s^{-1}) of electromagnetic radiation that has a wavelength of 0.53 m?

- a) 5.7×10^8
- b) 1.8×10^{-9}
- c) 1.6×10^8
- d) 1.3×10^{-33}

Explanation: The frequency and wavelength of electromagnetic radiation are related by the equation $c = \lambda\nu$, where c is the speed of light ($=3.00 \times 10^8 \text{ m/s}$), λ is the wavelength in m and ν is the frequency in s^{-1} or Hz. The frequency can be calculated by rearranging the above formula to get $\nu = c/\lambda = 3 \times 10^8 / 0.53 = 5.7 \times 10^8 \text{ s}^{-1}$


19. The energy of a photon of light is ____ proportional to its frequency and ____ proportional to its wavelength.

- a) directly, directly
- b) inversely, inversely
- c) inversely, directly
- d) directly, inversely

20. The wavelength of a photon of energy $5.25 \times 10^{-19} \text{ J}$ is _____ m.

- a) 2.64×10^6
- b) 3.79×10^{-7}
- c) 2.38×10^{23}
- d) 4.21×10^{-24}

Explanation: The wavelength and energy are related by the formula $E = hc/\lambda$, where h ($6.626 \times 10^{-34} \text{ Js}$) is Planck's constant, c is the speed of light ($3.00 \times 10^8 \text{ m/s}$) and λ is the wavelength in meters. The wavelength can then be calculated by rearranging the above formula as follows: $\lambda = hc/E = 6.63 \times 10^{-34} \times 3 \times 10^8 / 5.25 \times 10^{-19} = 3.79 \times 10^{-7} \text{ m}$



21. What is the frequency (s^{-1}) of a photon of energy $4.38 \times 10^{-18} \text{J}$?

- a) 438
- b) 1.45×10^{-16}
- c) **6.61×10^{15}**
- d) 2.30×10^7

Explanation: The frequency ν of this photon can be calculated by rearranging the equation $E = h \nu$ where E is the energy, h = Planck's constant and ν = frequency in s^{-1} . $\nu = E/h = 4.38 \times 10^{-18} / 6.63 \times 10^{-34} = 6.61 \times 10^{15}$

22. An electron in a Bohr hydrogen atom has energy of $-1.362 \times 10^{-19} \text{J}$. The value of n for this electron is _____.

- a) 1
- b) 2
- c) 3
- d) **4**

Explanation: The energy of an electron in a particular energy state in the hydrogen atom can be calculated by using the formula $E = -R_H/n^2 = (-2.18 \times 10^{-18} \text{J})/n^2$, where n is the principal quantum number for the energy state. The value of n can be found by rearranging the above formula as follows:

$$n = \sqrt{\frac{-2.18 \times 10^{-18} \text{J}}{-1.362 \times 10^{-19} \text{J}}} = 4$$

19. The $n = 2$ to $n = 6$ transition in the Bohr hydrogen atom corresponds to the ___ of a photon with a wavelength of ___ nm.

- a) emission, 411
- b) **absorption, 411**
- c) absorption, 657
- d) emission, 389

Explanation: There are 2 parts to this question. Since the electron is moving from a smaller value of n (n_i) to a larger value of n (n_f), it must be absorbing energy. The wavelength responsible for this transition can be calculated by using the formula: $E = R_H (1/n_i^2 - 1/n_f^2)$ & $E = hc/\lambda$, R_H is (Rydberg constant) = $2.18 \times 10^{-18} \text{J}$

20. How many quantum numbers are necessary to designate a particular electron in an atom _____?

- a) 3
- b) **4**
- c) 2
- d) 1

21. The _____ quantum number defines the shape of an orbital.

- a) spin
- b) magnetic
- c) principal
- d) **angular**

22. There are _____ orbitals in the third shell

- a) 25
- b) 4
- c) **9**
- d) 16

Explanation: The number of orbitals in a shell is easily calculated by the formula # of orbitals = n^2 where n = principal quantum number, which is 3 in this case.

23. The angular quantum number is 2 in _____ orbitals.

- a) s
- b) p
- c) **d**
- d) f

24. The $n = 1$ shell contains _____ p orbitals. All the other shells contain _____ p orbitals.

- a) 3, 6
- b) **0, 3**
- c) 6, 2
- d) 3, 3

Explanation: If $n = 1$, then the only possible value of ℓ is 0 which means that $n = 1$ can contain only s orbitals. When $n > 1$, the value of $\ell = 1$ is possible making the existence of 3 p orbitals possible.

25. The principal quantum number of the first d subshell is _____.

- a) 1
- b) 2
- c) **3**
- d) 4

26. The total number of orbitals in a shell is given by _____.

- a) L^2
- b) **n^2**
- c) $2n$
- d) $2n + 1$

28. Each p-subshell can accommodate a maximum of _____ electrons.

- a) **6**
- b) 2
- c) 10
- d) 3

Explanation: There are 3 different p orbitals: p_x , p_y and p_z . Each of these can contain 2 electrons leading to the maximum number of electrons as 6.

29. Each p-subshell can accommodate a maximum of _____ electrons.

- a) **6**
- b) 2
- c) 10
- d) 3

Explanation: There are 3 different p orbitals: p_x , p_y and p_z . Each of these can contain 2 electrons leading to the maximum number of electrons as 6.

30. The 3p subshell in the ground state of atomic xenon contains _____ electrons.

- a) 2
- b) **6**
- c) 36
- d) 10

Explanation: Since Xe is a noble gas, its subshells will be completely filled regardless of their principal quantum number. Thus, the 3p subshell will contain 6 electrons.

31. $[\text{Ar}]4s^23d^{10}4p^3$ is the electron configuration of a(n) _____ atom.

- a) **As**
- b) V
- c) P
- d) Sb

Explanation: The easiest way to answer this question is to count the total number of electrons and find which element that number corresponds to. The total number of electrons is = 18 (for the Ar) + 2 + 10 + 3 = 33 which corresponds to As.


32. The principal quantum number for the outermost electrons in a Br atom in the ground state is _____.

- a) 2
- b) 3
- c) **4**
- d) 5

Explanation: The electronic configuration of bromine is $[\text{Ar}]3d^{10}4s^24p^5$ shows that the outermost electrons are in the s and p orbitals in the 4th energy level making the principal quantum number = 4.

33. All of the _____ have a valence shell electron configuration ns^1 .

- a) noble gases
- b) halogens
- c) chalcogens
- d) **alkali metals**



34. Which one of the following is correct?

- a) $v + \lambda = c$
- b) $v/\lambda = c$
- c) $\lambda = cv$
- d) **$v\lambda = c$**

35. In the Bohr model of the atom,_____.

- a) electrons travel in circular paths called orbitals
- b) electrons can have any energy
- c) **electron energies are quantized**
- d) electron paths are controlled by probability

36. Which one of the following is not a valid value for the magnetic quantum number of an electron in a 5d subshell?

- a) 2
- b) **3**
- c) 0
- d) 1

Explanation: For an electron in the 5d subshell the value of $\ell = 2$ and the magnetic quantum number m_ℓ can have values from $-1, \dots, 0, \dots, +1$, meaning m_ℓ could not have a value = 3.

37. Which of the subshells below do not exist due to the constraints upon the angular quantum number?

- a) 2s
- b) **2d**
- c) 2p
- d) none of the above

Explanation: The values of the azimuthal quantum number “l” are decided by the values of the principal quantum number “n”. The values of l will only be from $0 \dots n - 1$. Thus, for $n = 2$, only the values of 0 and 1 will be possible for ℓ , which means that only the 2s and 2p orbitals will be possible.

38. An electron cannot have the quantum numbers $n =$ _____, $\ell =$ _____, $m_\ell =$ _____.





- a) 2, 0, 0
- b) 2, 1, -1
- c) 3, 1, -1
- d) **1, 1, 1**

Explanation: The values of 1, 1, 1 would be impossible since if $n = 1$, the only value of ℓ would be = 0.

39. Which quantum number determines the energy of an electron in a hydrogen atom?

- a) **n**
- b) n and ℓ
- c) m_ℓ
- d) ℓ

39. Which electron configuration represents a violation of the Pauli exclusion principle?



Explanation: According to the Pauli Exclusion Principle no two electrons in an atom cannot have the same 4 quantum numbers. The 2 electrons in the 2s orbital have the same value for their m_s which is not allowed. **(d)**

40. Which of the following is a valid set of four quantum numbers? (n , ℓ , m_ℓ , m_s)

- a) **2, 0, 0, + 1/2**
- b) 2, 2, 1, - 1/2
- c) 1, 0, 1, + 1/2
- d) 2, 1, +2, + 1/2

Explanation: Here is why only option (a) is the correct answer: In option (b), $\ell = 2$ which is not allowed, in (c) $m_\ell \neq 1$ since $l = 0$ and in (d) $m_\ell > 1$ which are all not allowed.

41. Which of the following is not a valid set of four quantum numbers? (n , ℓ , m_ℓ , m_s)

- a) 2, 0, 0, + 1/2
- b) 2, 1, 0, - 1/2
- c) **1, 1, 0, + 1/2**
- d) 1, 0, 0, + 1/2

Explanation: Since n can never be equal to ℓ , option c is the only set that is not valid

Test bank chapter (8)

Choose the correct answer

1. Elements in the modern version of the periodic table are arranged in order of increasing_____.
- oxidation number
 - atomic mass
 - average atomic mass
 - atomic number**

Explanation: The older version of the periodic table had the elements arranged in order of increasing atomic mass, but the modern version of the periodic table is based on the increasing order of atomic number.

2. The first ionization energies of the elements_____as you go from left to right across a period of the periodic table, and_____as you go from the bottom to the top of a group in the table.
- increase, increase**
 - increase, decrease
 - decrease, increase
 - decrease, decrease

Explanation: The ionization energies (IE s) of elements increase to the right in a row since larger amounts of energy need to be supplied to remove an electron. The elements become more non-metallic making it harder to remove an electron.

3. The_____have the most negative electron affinities
- alkaline earth metals
 - alkali metals
 - halogens**
 - transition metals

Explanation: The electron affinity of an element is defined as the energy change that occurs when an electron is added to a gaseous atom. The halogens have the most negative electron affinities indicating that they are most comfortable accepting an electron. The formation of an anion essentially gives the halogen atom the electron configuration of the nearest noble gas. The negative sign here indicates that the addition of an electron to the halogens results in energy being released by the halogen atom.

4. Na reacts with element X to form an ionic compound with the formula Na_3X . Ca will react with X to form .
- CaX_2
 - CaX
 - Ca_2X_3
 - Ca_3X_2**

Explanation: The element X must have 3 negative charges for it to form the compound Na_3X , since each Na has 1 positive charge, the formula of the compound formed by the reaction of Ca and X would have to be Ca_3X_2 .

5. Atomic radius generally increases as we move_____.
- down a group and from right to left across a period**
 - up a group and from left to right across a period
 - down a group and from left to right across a period
 - up a group and from right to left across a period

6. The atomic radius of main-group elements generally increases down a group because_____.
- effective nuclear charge increases down a group
 - effective nuclear charge decreases down a group
 - both effective nuclear charge increases down a group and the principal quantum number of the valence orbitals increases
 - the principal quantum number of the valence orbitals increases**

7. Which of the following correctly lists the five atoms in order of increasing size (smallest to largest)?

- $O < F < S < Mg < Ba$
- $F < O < S < Mg < Ba$**
- $F < O < S < Ba < Mg$
- $O < F < S < Ba < Mg$

Explanation: Fluorine and oxygen are in the same period (#2) and next to each other with F being the smallest of these 5 atoms. Ba is in group 2A and in row 6 (farthest “down” a group) and is the largest of the atoms. Mg is in group 2A and in the third period and hence will be bigger than F, O and S. Even though S is in the same period as Mg it is in group 6A making it smaller than Mg.

8. Which of the following correctly lists the five atoms in order of increasing size (smallest to largest)?

- $F < K < Ge < Br < Rb$
- $F < Ge < Br < K < Rb$
- $F < K < Br < Ge < Rb$
- $F < Br < Ge < K < Rb$**

Explanation: Fluorine is in group 7A and period 2 making it the smallest of the 5 atoms here. Br is also in group 7A but is in period 4 making it larger than F, Ge is in group 4A and also in period 4 but is to the left of Br making it larger than Br. K and Rb are both in group 1A but K is in period 4 and Rb is in period 5, making the Rb atom the largest of all the 5 atoms.

9. Of the following atoms, which has the largest first ionization energy?

- Br
- O**
- C
- P

Explanation: The ionization energy (IE) typically increases from left to right in a period and decreases from top to bottom in a group. Thus for C and O which are in the same period, O will have the larger IE. Br is in period #4 and will have the lowest IE out of these 4 elements.

10. Of the following elements, which has the largest first ionization energy?

- Na
- Al
- Se
- Cl**

Explanation: The ionization energy (IE) typically increases from left to right in a period and decreases from top to bottom in a group. Na, Al and Cl are all in period 3 with the chlorine atom to the farthest right and will have the highest IE.

11. Which ion below has the largest radius?

- a) Cl^-
- b) K^+
- c) Br^-
- d) F^-

Explanation: Typically cations are smaller than their parent atoms while anions are larger than the parent atoms. Of the atoms here, the Br atom would be the largest as it is farthest down the group and hence its anion also will be the largest ion.

12. The ion with the smallest radius is _____.

- a) Br^-
- b) Cl^-
- c) O^{2-}
- d) F^-

Explanation: Typically anions are larger than the parent atoms. Of the atoms here, the F atom would be the smallest as it is farthest up the group and hence its anion also will be the smallest ion

13. Which of the following is an isoelectronic series?

- a) B^{5-} , Si^{4+} , As^{3-} , Te^{2-}
- b) O^{2-} , F^- , Ne , Na^+
- c) S, Cl, Ar, K

Explanation: Isoelectronic series contain a combination of atoms and ions or only ions with the same number of electrons. Here the series containing the O^{2-} , F^- , Ne and Na^+ is the only one where all atoms/ions contain 10 electrons.

14. _____ is isoelectronic with argon and _____ is isoelectronic with neon.

- a) Cl^- , F^-
- b) Cl^- , Cl^+
- c) F^+ , F^-
- d) Ne , Kr^+

Explanation: The Cl^- ion has 18 electrons and is isoelectronic with argon while the F^- ion has 10 electrons making it isoelectronic with neon.

15. Chlorine is much more ability to exist as an anion than is sodium. This is because _____.

- a) chlorine is bigger than sodium
- b) chlorine has a greater ionization energy than sodium does
- c) **chlorine has a greater electron affinity than sodium does**
- d) chlorine is a gas and sodium is a solid

16. The alkaline earth metals are found in _____ of the periodic table.

- a) Group 1A
- b) **Group 2A**
- c) Group 7A
- d) Group 8A

17. How many *valence electrons* does a boron atom (B) have?

- a) 1
- b) 3
- c) 5
- d) 7

18. Which ion is *isoelectronic* with Ar?

- a) Ni^{2+}
- b) F^-
- c) Br^-
- d) K^+

19. Which of these choices is the electron configuration of the iron (III) ion (Fe^{3+})?

- a) $[\text{Ar}]4s^23d^3$
- b) $[\text{Ar}]4s^13d^5$
- c) $[\text{Ar}]3d^5$
- d) $[\text{Ar}]3d^6$

20. In what group of the periodic table is the element with the electron configuration $[\text{Ar}]4s^23d^{10}4p^1$?

- a) 1A
- b) 2A
- c) 3A
- d) 4A

21. How many *valence electrons* does a tin (Sn) atom have?

- a) 2
- b) 4
- c) 14
- d) 36

22. Which of these ground-state ions has unpaired electrons?

- a) P^{3-}
- b) V^{5+}
- c) S^{2-}
- d) Sc^{2+}

23. Consider the element with the electron configuration $[\text{Xe}]6s^24f^7$. This element is

- a) a representative element.
- b) a lanthanide element.
- c) a nonmetal.
- d) an actinide element

24. If the radius of atom X is greater than the radius of atom Y, then it is also likely that

- a) X has a larger electron affinity than Y does.
- b) X has a larger first ionization energy than Y does.
- c) X has greater metallic character than Y does.

25. Arrange these ions in order of increasing ionic radius: K^+ , P^{3-} , S^{2-} , Cl^- .

Increasing radius \rightarrow

Row 1 $K^+ < Cl^- < S^{2-} < P^{3-}$

Row 2 $K^+ < P^{3-} < S^{2-} < Cl^-$

Row 3 $P^{3-} < S^{2-} < Cl^- < K^+$

Row 4 $Cl^- < S^{2-} < P^{3-} < K^+$

Row 5 $Cl^- < S^{2-} < K^+ < P^{3-}$

- a) Row 1
- b) Row 2
- c) Row 3
- d) Row 4

26. Selenium (${}_{34}Se$) element is

- a) a nonmetal
- b) found in group 6A
- c) found in period 2
- d) **both a and b**

28. The outer electron configuration of the noble gases is

- a) **$ns^2 np^6$**
- b) $ns^2 nd^{10}$
- c) $ns^2 np^4$
- d) $ns^2 np^8$

29. Which of the following species is isoelectronic with Cl^-

- a) **K^+**
- b) Na^+
- c) O^{2-}
- d) F^-

30. Gallium (Ga) element is found in the periodic table in

- a) period 3, group 1B
- b) period 3, group 4 A
- c) period 4, group 1A
- d) **period 4, group 3A**

31. Titanium (Ti) element is found in the periodic table in

- a) s-block
- b) p-block
- c) **d-block**
- d) f-block

32. Write the electronic configuration for Co^{+2}

- a) $[Ar] 4s^2 3d^5$
- b) **$[Ar] 3d^7$**
- c) $[Ar] 4s^1 3d^6$



33. Select the correct order of radius of the two ions

- a) $A^+ > A^-$
- b) $A^- < A$
- c) $A^{2+} > A^+$
- d) $A^{2+} < A^+$**

34. Two ions are referred to as isoelectronic if they have the same number of

- a) electrons.**
- b) protons.
- c) atoms.
- d) neutrons.

35. The energy required to remove an electron from an atom in its ground state is known as the

- a) potential energy.
- b) activation energy.
- c) electron affinity.
- d) ionization energy.**

36. Which will have the highest ionization energy?

- a) C
- b) N**
- c) O
- d) B

37. Order the following (N^{3-} , Li^+ , C, O^{2-}) according to increasing atomic/ionic radius.

- a) $\text{C} < \text{Li}^+ < \text{O}^{2-} < \text{N}^{3-}$
- b) $\text{N}^{3-} < \text{O}^{2-} < \text{C} < \text{Li}^+$
- c) $\text{Li}^+ < \text{C} < \text{N}^{3-} < \text{O}^{2-}$
- d) $\text{Li}^+ < \text{C} < \text{O}^{2-} < \text{N}^{3-}$**

Test bank chapter (9)

Choose the most correct answer

1. What are the two types of chemical bonds commonly found in compounds ?
 - a) ionic and covalent.
 - b) ionic and electrolytic.
 - c) **ionic and covalent.**
 - d) electrolytic and compound.

2. Which type of electrons are used by atoms to form chemical bonds ?
 - a) core electrons.
 - b) **valence electrons.**
 - c) lone pair electrons.
 - d) unpaired electrons.

3. What is the statement of “Atoms tend to gain, lose, or share electrons until they are surrounded by eight valence electrons” called ?
 - a) the rule of octaves.
 - b) the double quartet rule.
 - c) the eight electron rule.
 - d) **the octet rule.**

4. What type of orbital will lose the electron first when a transition metal atom becomes a +ve ion?
 - a) p
 - b) f
 - c) d
 - d) **s**

5. What type of bonds does a molecule of CS₂ contain ?
 - a) two single bonds.
 - b) **two double bonds.**
 - c) one single bond and one double bond.
 - d) one single bond and one triple bond.

6. What is the correct electron-dot structure of an atom with atomic number Z= 5 in the ground state? **ANS. B**

- (A) $\cdot\overset{\cdot}{\underset{\cdot}{\text{X}}}\cdot$
- (B) $\overset{\cdot}{\underset{\cdot}{\text{X}}}\cdot$
- (C) $\cdot\overset{\cdot}{\underset{\cdot}{\text{X}}}\cdot$
- (D) $\overset{\cdot}{\underset{\cdot}{\text{X}}}\cdot$

7. Which compound below contains an atom that is surrounded by more than an octet of electrons?

- a) **PF₅**
- b) CH₄
- c) NBr₃
- d) OF₂

8. Which choice below correctly lists the elements in order of increasing electronegativity?

- a) **C < N < O < F**
- b) N < C < O < F
- c) N < C < F < O
- d) C < N < F < O

9. Which atom sometimes violates the octet rule?

- a) C
- b) N
- c) O
- d) **S**

10. How many resonance structures can be drawn for NO₃⁻?

- a) 1
- b) 2
- c) **3**
- d) 4

11. Considering formal charge, what is the preferred Lewis structure of NCO⁻? **ANS.3**



12. What is the correct formal charge on sulfur (S) in Lewis structure of (SO₄²⁻) that satisfy the octet rule?

- a) **+2**
- b) -2
- c) +1
- d) 0

13. What is the correct formal charge on sulfur (S) in the favorable Lewis structure of (SO₄²⁻) ?

- a) +2
- b) -2
- c) +1
- d) **0**

Chemical Bonding I: Basic Concepts

14. Which of these pairs of elements would be most likely to form an ionic compound?

- a) Cl and I
- b) Al and K
- c) Cl and Mg
- d) C and S

15. Which of these covalent bonds is the most polar (i.e., highest percent ionic character)?

- a) Al—I
- b) Si—I
- c) Al—Cl
- d) Si—Cl

15. which of these structures is the correct Lewis structure for CS₂ ? ANS.c

- a) $\overset{\cdot\cdot}{\text{C}}=\overset{\cdot\cdot}{\text{S}}-\overset{\cdot\cdot}{\text{S}}$
- b) $\overset{\cdot\cdot}{\text{S}}-\overset{\cdot\cdot}{\text{C}}-\overset{\cdot\cdot}{\text{S}}$
- c) $\overset{\cdot\cdot}{\text{S}}=\text{C}=\overset{\cdot\cdot}{\text{S}}$
- d) $\overset{\cdot\cdot}{\text{S}}=\overset{\cdot\cdot}{\text{C}}-\overset{\cdot\cdot}{\text{S}}$

16. How many lone pairs in the N₂ molecule are there?

- a) 1
- b) 2
- c) 3
- d) 4

17. Classify the O-H bond in CH₃OH as ionic, polar covalent, or nonpolar covalent?

- a) Ionic
- b) polar covalent
- c) nonpolar covalent

18. How many single bond(s), double bond(s) and lone pair(s) are there in Lewis structure for a chlorate ion, ClO₃⁻?

- a) 2, 1, 10
- b) 3, 0, 9
- c) 2, 1, 8
- d) 3, 0, 10

19. How many resonance structures are there for the sulfur dioxide molecule that satisfy the octet rule ?

- a) 1
- b) 2
- c) 3
- d) none of these.

20. What is the formal charge on the oxygen atom in N₂O (the atomic order is N-N-O)?
- 0
 - +1
 - 1**
 - 2
21. Which of these substances will display an incomplete octet in its Lewis structure?
- CO₂
 - Cl₂
 - ICl
 - NO**
22. How many paired and unpaired electrons are there in the Lewis symbol for a phosphorus atom?
- 4, 2
 - 2, 4
 - 4, 3
 - 2, 3**

Explanation: Read the question carefully here, you are being asked for how many valence electrons are paired and how many are unpaired. The abbreviated electron configuration of the P atom is given by [Ne] 3s²3p³. The outermost electrons would be arranged as 2 electrons paired and 3 electrons unpaired as shown below:



23. What is the most likely forms of magnesium ion based on the octet rule?
- Mg²⁺**
 - Mg²⁻
 - Mg⁶⁺
 - Mg⁶⁻

Explanation: According to the octet rule the Mg atom will achieve an octet by losing its 2 outermost electrons and thus gaining 2+ charges. Since Mg is located in the alkali metal group it will lose electrons rather than gaining them.

24. What is most likely forms of phosphorus ion based on the octet rule?
- P³⁺
 - P⁵⁻
 - P⁵⁺
 - P³⁻**

Explanation: According to the octet rule the phosphorus atom should gain 3 electrons, thus gaining 3 negative charges and forming the phosphide ion.

25- What is the only noble gas without eight valence electrons ?

- a) Ar
- b) Ne
- c) He
- d) Kr

Explanation: The noble gases are characterized by the presence of eight electrons in their outermost shell with one notable exception of Helium. Since He has only 2 electrons it can never have 8 in its outermost shell.

26- What is the maximum number of double bonds that a hydrogen atom can form?

- a) 0
- b) 1
- c) 2
- d) 3

Explanation: Each hydrogen atom has a single electron in its valence shell and as a result can form only one bond. It cannot form a double bond as it does not have the necessary electrons to share.

28. What is the maximum number of double bonds that a carbon atom can form?

- a) 4
- b) 1
- c) 2
- d) 0

Explanation: Each carbon atom has 4 valence electrons that it can share with other atoms. Since each double bond corresponds to a pair of electrons, the carbon atom can form only 2 double bonds.

29. Given the electronegativities below, which covalent single bond is most polar?

Atom	H	C	N	O
Electronegativity	2.1	2.5	3.0	3.5

- a) C-H
- b) N-H
- c) O-H
- d) O-N

Explanation: Bond polarity can be judged based on the differences between the electronegativities of the atoms involved. Of the available choices, the bond between O and H will have the largest electronegativity difference making it the most polar bond in this group.

30. How many valence electrons does the ICl_4^- ion have?

- a) 34
- b) 36
- c) 35
- d) 28

Explanation: valence electrons $A = (7 \times 1) + (7 \times 1) + 1 = 36$


Chemical Bonding I: Basic Concepts

- 31-What is the trend of the electronegativity from left to right within a period and from top to bottom within a group?
- a) decreases, increases
 - b) increases, increases
 - c) stays the same, increases
 - d) **increases, decreases**
32. How many nonbonding and bonding electron pairs in the central phosphorus atom are there in Lewis structure of PF_3 ?
- a) 2, 2
 - b) **1, 3**
 - c) 3, 1
 - d) 1, 2
33. Which of the following molecules contains both ionic and covalent bonds?
- a) C_5H_{12}
 - b) **NaClO_4**
 - c) CaCl_2
 - d) H_2O
34. What is the term of the ability of an atom in a molecule to attract electron density to itself ?
- a) **Electronegativity**
 - b) Electron affinity
 - c) Diamagnetism
 - d) Ionization energy
- 35-Which one of the below is the most polar bond ?
- a) Br-H
 - b) I-H
 - c) **Cl-H**
 - d) H-H

Test bank chapter (14)


Choose the most correct answer

1. Which is the correct equilibrium constant expression for the following reaction?



- a) $K_c = [\text{Fe}_2\text{O}_3] [\text{H}_2]^3 / [\text{Fe}]^2 [\text{H}_2\text{O}]^3$
- b) $K_c = [\text{H}_2] / [\text{H}_2\text{O}]$
- c) $K_c = [\text{H}_2\text{O}]^3 / [\text{H}_2]^3$
- d) $K_c = [\text{Fe}]^2 [\text{H}_2\text{O}]^3 / [\text{Fe}_2\text{O}_3] [\text{H}_2]^3$

2. The following reactions occur at 500 K. Arrange them in order of increasing tendency to proceed to completion (least → greatest tendency).



- a) $2 < 1 < 3$
- b) $1 < 2 < 3$
- c) $2 < 3 < 1$
- d) $3 < 2 < 1$


3. Calculate K_p for the below reaction if K_c at for this reaction is 2.1×10^{-2} at 400°C .



- a) 0.689
- b) 0.115
- c) 0.137
- d) 1.2

Chemical Equilibrium

4. For the following reaction:



$K_c = 50.2$ at $445\text{ }^\circ\text{C}$. If $[\text{H}_2] = [\text{I}_2] = [\text{HI}] = 1.75 \times 10^{-3}\text{M}$ at $445\text{ }^\circ\text{C}$, which one of these statements is true?


- a) The system is not at equilibrium; thus, no concentration changes will occur.
- b) The concentrations of HI and I_2 will increase as the system approaches equilibrium.
- c) The concentration of HI will increase as the system approaches equilibrium.
- d) The concentrations of H_2 and HI will decrease as the system moves toward equilibrium.

5. For the below reaction at equilibrium, which choice gives a change that will shift the position of equilibrium to favor formation of more products?



- a) Increase the total pressure by decreasing the volume.
- b) Add more NO.
- c) Remove Br_2 .
- d) Lower the temperature.

6 - For the following reaction at equilibrium in a reaction vessel, which one of these changes would cause the Br_2 concentration to decrease?



- a) Increase the temperature.
- b) Remove some NO.
- c) Add more NOBr.
- d) Compress the gas mixture into a smaller volume.


7. For the below reaction at equilibrium, if we increase the reaction temperature, the equilibrium will:



- a) shift to the right.
- b) shift to the left.
- c) not shift.
- d) The question cannot be answered because the equilibrium constant is not given.

Chemical Equilibrium

8. For the equilibrium reaction:



Which one of these factors would cause the equilibrium constant to increase?

- a) Decrease the temperature.
- b) Increase the temperature.
- c) Add SO_2 gas.
- d) Remove O_2 gas.

9. The reaction below is endothermic. If the temperature is increased,




- a) more SO_3 will be produced.
- b) K_c will increase.
- c) K_c will decrease.
- d) no change will occur in K_c .

10. If a catalyst is added to a chemical reaction, the equilibrium yield of a product will be, and the time taken to come to equilibrium will bethan before.

- a) higher; less
- b) lower; the same
- c) higher; the same
- d) the same; less

11- For the reaction:



$K_c = 0.0600$ at a certain temperature. In an equilibrium mixture of the three gases, $[\text{NH}_3] = 0.242 \text{ M}$ and $[\text{H}_2] = 1.03 \text{ M}$. What is the concentration of N_2 in this system?

- a) 3.9 M
- b) 0.003 M
- c) 0.89 M
- d) 1.12 M

Chemical Equilibrium


12. Consider the reaction,



If an equilibrium mixture of these three substances is compressed, equilibrium will, because

- a) shift to the right; higher pressure favors fewer moles of gas
- b) shift top the right; higher pressure favors more moles of gas
- c) shift to the left; higher pressure favors fewer moles of gas
- d) shift to the left; higher pressure favors more moles of gas


13- Consider the equilibrium system:



If more C(s) is added, the equilibrium will; if CO is removed the equilibrium will

- a) shift to the left; shift to the left
- b) shift to the right; shift to the left
- c) be unchanged; shift to the right
- d) be unchanged; shift to the left

14. Consider the exothermic reaction at equilibrium:



If the system is cooled, the equilibrium will, because

- a) shift to the left; decreased temperature favors an exothermic reaction
- b) shift to the right; decreased temperature favors an exothermic reaction
- c) shift to the right; decreased temperature favors an endothermic reaction
- d) shift to the left; decreased temperature favors an endothermic reaction

15. A large value of the equilibrium constant indicates that when the reaction reaches equilibrium, mostly _____ will be present.


- a) reactants
- b) products
- c) catalysts
- d) shrapnel

Chemical Equilibrium

16. When equilibrium is achieved?

- a) $Q > K$
- b) $Q < K$
- c) $Q = K$
- d) $Q^2 = K$


17. for the following reaction:



If all species are gases and H_2 is added, the amount of CO present at equilibrium will:

- a) increase.
- b) decrease.
- c) remain unchanged.
- d) disappear.


18. For the reaction:



If all species are gases and H_2O is added, the amount of CO present at equilibrium will:

- a) increase.
- b) decrease.
- c) remain unchanged.
- d) disappear.


19. For the reaction:



If the reaction is endothermic and the temperature is raised, the amount of CO present will:

- a) increase.
- b) decrease.
- c) remain unchanged.
- d) disappear.


20. For the reaction:



If all species are gases and the container is compressed, the amount of CO present will:


- a) increase.
- b) decrease.
- c) remain unchanged.
- d) disappear.

21. What is K_p in terms of K_c for the following reaction?



- a) $K_p = K_c RT$
- b) $K_p = K_c / RT$
- c) $K_p = K_c R/T$
- d) $K_p = K_c / (RT)^2$

22. What is the correct equilibrium constant expression for the reaction:



a.
$$\frac{[\text{PCl}_3]^4}{[\text{P}_4][\text{Cl}_2]^6}$$

c.
$$\frac{[\text{PCl}_3]^4}{[\text{Cl}_2]^6}$$

b.
$$\frac{1}{[\text{Cl}_2]^6}$$

d.
$$\frac{[\text{PCl}_3]^4}{[\text{P}][6\text{Cl}_3]}$$

23. The equation relating K_p and K_c is:


- a) $K_p = k_c (RT)^{\Delta n}$
- b) $K_p = K_c RT^{\Delta n}$
- c) $K_c = K_p RT^{\Delta n}$
- d) $K_c = K_p (RT)^{\Delta n}$

Chemical Equilibrium

24. K_p will be equal to K_c if:

- a) $\Delta n = 1$
- b) $\Delta n = 0$
- c) $\Delta n = -1$
- d) $RT = 0$

25. Consider the reversible reaction at equilibrium at 392 °C:



The partial pressures are found to be: A: 6.70 atm, B: 10.1 at, C: 3.60 atm. Evaluate K_p for this reaction.


- a) 7.94×10^{-3}
- b) 0.0794
- c) 0.794
- d) 7.94

26. Which of the following will result in an equilibrium shift to the right?



- a) Increase temperature/increase volume
- b) Increase temperature/decrease volume
- c) decrease temperature/increase volume
- d) decrease temperature/decrease volume

27. Which accurately reflects the change in concentration that will occur if O_2 is added to disturb the equilibrium?



	[NO]	[O ₂]	[NO ₂]
a)	Increase	Increase	Increase
b)	Increase	Increase	Decrease
c)	Decrease	Decrease	Decrease
d)	Decrease	Increase	Increase

Test Bank Chapter (15)

Choose the most correct answer:

1- What is the concentration of H^+ in a 2.5 M HCl solution?

- a) 0
- b) 1.3 M
- c) 2.5 M
- d) 5.0 M

2. What is the OH^- ion concentration in a 5.2×10^{-4} M HNO_3 solution?

- a) 1.9×10^{-11} M
- b) 1.0×10^{-7} M
- c) 5.2×10^{-4} M
- d) Zero

3. Calculate the H^+ ion concentration in lemon juice having a pH of 2.4

- a) 4.0×10^{-2} M
- b) 250 M
- c) 0.38 M
- d) 4.0×10^{-3} M

4. Calculate the pH of a 6.71×10^{-2} M NaOH solution.

- a) 12.83
- b) 2.17
- c) 11.82
- d) 6.71

5. What is the pH of 0.0200 M aqueous solution of HBr?

- a) 1.00
- b) 1.70
- c) 2.30
- d) 12.30

6. The pOH of a solution of NaOH is 11.30, what is the $[H^+]$ for this solution?

- a) 2.0×10^{-3}
- b) 2.5×10^{-3}
- c) 5.0×10^{-12}
- d) 4.0×10^{-12}

7. What is the pH of a 0.04 M aqueous solution of KOH?
- a) 12.60
 - b) 10.30
 - c) 4.00
 - d) 1.40
8. What is the approximate pH of a solution labeled 6×10^{-5} M HBr?
- a) 4.2
 - b) 4.5
 - c) 5.8
 - d) 9.8
9. If the pH = 2 for an HNO₃ solution, what is the concentration of HNO₃?
- a) 0.10
 - b) 0.20
 - c) 0.010
 - d) 0.020
10. A solution in which $[H^+] = 10^{-8}$ M has a pH of ____ and is ____.
- a) 8, acidic
 - b) 6, basic
 - c) -6, basic
 - d) 8, basic
11. Which of the following solutions has the lowest pH at 25°C? (No calculations required.)
- a) 0.2 M NaOH
 - b) 0.2 M NH₃
 - c) 0.2 M HCl
 - d) pure water
12. Calculate the pH of a 3.5×10^{-3} M HNO₃ solution.
- a) -2.46
 - b) 0.54
 - c) 2.46
 - d) 3.00
13. The pH of 2.6×10^{-2} M KOH is
- a) 12.41
 - b) 15.59
 - c) 2.06
 - d) 7.00

14. What is the $[\text{H}^+]$ ion in a 4.8×10^{-2} M KOH solution?

- a) 2.08×10^{-13} M
- b) 1×10^{-7} M
- c) 4.8×10^{-11} M
- d) 4.8×10^{-2} M

15. What is the $[\text{OH}^-]$ ion in a 5.2×10^{-4} M HNO_3 solution?

- a) 1.9×10^{-11} M
- b) 1.0×10^{-7} M
- c) 5.2×10^{-4} M
- d) zero

Test bank chapters (24 & 25)Choose the correct answer

1. C₁₀H₂₂ is the formula of an

- a) **alkane.**
- b) alkene.
- c) alkyne.
- d) aromatic hydrocarbon.

2. A molecule with the formula C₃H₈ is a

- a) hexane
- b) **propane**
- c) decane
- d) butane


3. Which compound below does not have geometric isomers (cis-trans isomers)?

- a) **1-butene**
- b) 2-butene
- c) 2-pentene
- d) 3-hexene

4. The hybridization of carbon atoms in alkanes is

- a) sp
- b) sp²
- c) **sp³**
- d) sp³d

5. Select the correct IUPAC name for



- a) 1,1,3-trimethylpentane
- b) 1-ethyl-1,3-dimethylbutane
- c) **2,4-dimethylhexane**
- d) 3,5-dimethylhexane.


6 - An alkane with seven carbon atoms in a linear configuration is called a

- a) hexene
- b) heptene
- c) heptylpane
- d) **heptane**

7. Which type of functional group does not include a carbonyl group in its structure?

- a) carboxylic acid
- b) ether
- c) ketone
- d) aldehyde

8. Vanillin is used as a flavoring agent. Identify the functional group circled.



- a) aldehyde
- b) ketone
- c) carboxylic acid
- d) Alcohol

9. The formula ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$) represents:

- a) an alcohol
- b) an alkene
- c) an alkyne
- d) an unsaturated hydrocarbon

10. Which of the following is a ketone?

- a) $\text{CH}_3\text{CH}_2\text{COCH}_3$
- b) $\text{CH}_3\text{CH}_2\text{CHO}$
- c) CH_3OCH_3
- d) $\text{CH}_3\text{CH}_2\text{COOH}$

11. The general formula for *alkenes* is

- a) $\text{C}_n\text{H}_{2n+2}$
- b) $\text{C}_{2n}\text{H}_{2n}$
- c) C_nH_{n+2}
- d) C_nH_{2n}





12. Which of these molecules is *unsaturated*?

- a) C_3H_8
- b) CH_3OH
- c) C_5H_{10}
- d) CH_4

8. Which of these species is an aromatic compound?

- a) C_2H_2
- b) C_6H_{12}
- c) $C_6H_4Br_2$
- d) C_5H_{10}

9. Which of these species are *structural isomers* of C_6H_{14} ?







- a) I and II
- b) I and III
- c) II and III
- d) II and IV





10. The compound that has a triple bond between one pair of carbon atoms is called

- a) an alkane.
- b) a chlorofluorocarbon.
- c) an alkyne.
- d) an alkene.

11. The correct structure for 2,3,3-trimethylpentane is



12. Which one of these structures represents an *ester* functional group?

- a) 
- b) 
- c) 
- d) 

13. The functional group (RCOR) is characteristic of organic _____

- a) ketones
b) acids
c) aldehydes
d) esters

14. Which of the following hydrocarbons does not have isomers?

- a) C_7H_{16}
b) C_6H_{14}
c) C_5H_{10}
d) C_3H_8

15. Which of the following does NOT exhibit geometric isomerism? (Hint: draw them!)

- a) 4-octene
b) 2-pentene
c) 3-hexene
d) 1-hexene

16. A protein is:

- a) a polymer of amino acids
- b) a fatty acid ester of glycerol
- c) a polysaccharide
- d) an addition polymer


17. A peptide bond (also called an amide bond) joins two amino acids together. What atoms are linked by this bond?

- a) C — O
- b) C — H
- c) C — N
- d) N — S

18. An amino acid is a compound that contains at least




- a) one amino group and one amide group.
- b) two amino groups and one carboxylic acid group.
- c) one hydroxyl group and one methyl group.
- d) one carboxylic acid group and one amino group

19. The functional group found in proteins is called a (an)



- a) amide.
- b) carboxylic acid.
- c) amine.
- d) amino acid.

20. Which one of these choices is the general structural formula of an amino acid?



21. Which one of these structures represents a ketone functional group?

