

# MATH203 Calculus

Dr. Bandar Al-Mohsin

School of Mathematics, KSU

2/3/14


# Double Integrals

## Riemann Sum

Let  $f$  be a function of two variables defined on region  $R$ , and Let  $P = \{R_k\}$  be an inner partition of  $R$ . A Riemann sum of  $f$  for  $P$  is any sum of the form

$$\sum_k f(u_k, v_k) \Delta A_k \quad (1)$$

where  $u_k, v_k$  is a point in  $R_k$  and  $\Delta A_k$  is the area of  $R_k$



# Double Integrals

## Remarks

1- The summation (1) extends over all the subregions  $R_1, R_2, \dots, R_n$  of  $P$ .

2-  $\lim_{\|P\| \rightarrow 0} \sum_k f(u_k, v_k) = C \quad C \in \mathbb{R},$  if  $f$  is continuous on  $R$ .

## Double Integrals of $f$ over $R$

If  $f$  is a function of two variables that is defined on a region  $R$ . The double integral of  $f$  over  $R$  is

$$\iint_R f(x, y) dA = \lim_{\|P\| \rightarrow 0} \sum_k f(x_k, y_k) \Delta A_k \quad (2)$$


provide the limit exists.

# Double Integrals

## Remarks

1- If  $f(x, y) \geq 0$  and continuous throughout the region  $R$ , then the double integral  $\iint_R f(x, y) dA$  may be used to find the Volume  $V$  of the solid  $Q$  that lies under the graph of  $z = f(x, y)$  and over  $R$ , i.e.

$$V = \iint_R f(x, y) dA, \quad f(x, y) \geq 0 \text{ on } R$$



2- If the region  $R$  describes the base of a mountain and  $f(x, y)$  is the height at point  $(x, y)$ , then the double integral  $\iint_R f(x, y) dA$  is the

Volume of the mountain.

3- If the region  $R$  describes the surface of a lake and  $f(x, y)$  is the depth of the water at point  $(x, y)$ , then the double integral  $\iint_R f(x, y) dA$  is the


Volume of the water in the lake.

4- If  $f(x, y) \leq 0$  and continuous throughout the region  $R$ , then the double integral  $\iint_R f(x, y) dA$  is the negative of the Volume  $V$  of the solid  $Q$  that lies over the graph of  $z = f(x, y)$  and under  $R$ .

# Properties of double integrals

- $\iint_R c f(x, y) dA = c \iint_R f(x, y) dA$  for every real number  $c$ .
- $\iint_R [f(x, y) \pm g(x, y)] dA = \iint_R f(x, y) dA \pm \iint_R g(x, y) dA$ .
- If  $Q$  is the union of two non-overlapping regions  $R$  and  $S$ ,  

$$\iint_Q f(x, y) dA = \iint_R f(x, y) dA + \iint_S f(x, y) dA$$



- If  $f(x, y) \leq 0$  throughout the region  $R$ , then  $\iint_R f(x, y) dA \leq 0$


## Evaluation theorem (1) Rectangular Regions

Let  $f$  be continuous function on a closed rectangular region  $R$ , then  $\iint_R f(x, y) dA$  can be evaluated by using an **iterated integral** of the following type

$$\int_a^b \left[ \int_c^d f(x, y) dy \right] dx = \int_a^b \int_c^d f(x, y) dy dx$$

or

$$\int_c^d \left[ \int_a^b f(x, y) dx \right] dy = \int_c^d \int_a^b f(x, y) dx dy$$



# Double Integrals

## Remarks

- 1-  $\int_c^d f(x, y)dy$  is partial integration w.r.t  $y$ , regarding  $x$  as a constant.
- 2-  $\int_a^b f(x, y)dx$  is partial integration w.r.t  $x$ , regarding  $y$  as a constant.

## Examples

Evaluate the following integrals:

$$(1) \int_1^2 \int_{-1}^2 (12xy^2 - 8x^3)dydx.$$

$$(2) \int_{-1}^2 \int_1^2 (12xy^2 - 8x^3)dx dy.$$

$$(3) \int_1^3 \int_2^4 (40 - 2xy)dx dy.$$

$$(4) \int_1^2 \int_{1-x}^{\sqrt{x}} x^2 y dx dy.$$




# Double Integrals

## Non-Rectangular Regions

**CASE 1** An iterated integral may be defined over the region  $R_x$  as shown below

$$\int_a^b \left[ \int_{g(x)}^{h(x)} f(x, y) dy \right] dx = \int_a^b \int_{g(x)}^{h(x)} f(x, y) dy dx$$




# Double Integrals

## Non-Rectangular Regions


**CASE 2** An iterated integral may be defined over the region  $R_y$  as shown below

$$\int_c^d \left[ \int_{p(y)}^{q(y)} f(x, y) dx \right] dy = \int_c^d \int_{p(y)}^{q(y)} f(x, y) dx dy$$




# Double Integrals


## Some important graphs




(a)  $y = \sqrt{x}$




(b)  $y = x^3$




(c)  $y = \sqrt{1-x^2}$



(d)  $y = e^x$



(e)  $y = \ln(x)$



(f)  $y = \tan x$

Figure: Some important graphs

# Double Integrals

## Examples

Sketch the region bounded by the graphs of :

(1)  $y = \sqrt{x}$  and  $y = x^3$ .

(2)  $y = \sqrt{1-x^2}$  and  $y = 0$ .

for  $f(x, y) = x - y$ .